



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines


Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search


Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>





R441f

W. G. FARLOW







FLORAE SCANDINAVIAE

PRODROMUS

ENUMERANS

P L A N T A S

SVECIAE, LAPPONIAE, FINLANDIAE ET
POMERANIAE

AC

DANIAE, NORVEGIAE, HOLSATIAE
ISLANDIAE GROENLANDIAEQUE

A U C T O R E

ANDREA JAHANNE RETZIO

PH. MAG. HIST. NAT. ET OECON. PROF. R. O. SOC. PHYSIOGR. LUND.
SECR. R. ACAD. SCIENT. SVEC. R. SOC. PATR. HGLM. IMP. PETROP.
OECON. RR. SOC. SCIENT. ET MED. HAFN. SOC. LINNEAN. LOND.
ACAD. SCIENT. PATAV. ET MANTUAN. SOC. NAT. CURIOS.
BEROL. SCIENT. ET RE. LL. GOTHOB. EDUCAT. HOLM. PATRIOT.
HASSO-HOMB. MEMBRI, ET R. ACAD. SCIENT. TAURIN.
AC SOC. OECON. LIPS. CORRESP.

EDITIO ALTERA.

LIPSIAE,

APUD SIEGFRIED LEBRECHT CRUSIUS, 1795.

Handwritten note:

Handwritten text, mostly illegible due to extreme fading and noise. The text appears to be organized into several lines or paragraphs, but the specific words and sentences cannot be discerned.

**SOCIETATI
LINNEANA E
LONDINENSIS.**

D. D. D.

CINERIS LINNEI CULTOR.

THE JOURNAL OF THE
ROYAL ANTHROPOLOGICAL INSTITUTE

Vol. 10, Part 1, 1900

L. B.

Ultra sexennium est, quod in Bibliopoliis nostris defuerunt Exemplaria Florae Svecicae Linneanae. Inaudivi equidem Beat. Salvium novam meditatam esse editionem, sed praematura ejus mors forte in causa fuit, cur illam adhuc desideremus. Apoplexia tandem lethali correptus quoque fuit immortalis Florae Auctor, quae impedivit, quominus laboriosissimus antea Senex manum postea operibus emendandis admoveret. Aliis itaque hanc reliquit operam, ipse cum summa animi tranquillitate altiora petens, contentusque educasse tot artis Filios, relicta olim affecuturos.

Elegantia concinnitasque Florae Linneanae omnium oculis patet. Exempli instar omnium Florarum, etiam olim edendarum inserviit, inservietque, quo nihil egregius, nihil magis completum.

Verum Scriptorum Linneanorum pretium, ego sane, si ullus, dudum expertus sum, qui plane nulla in Botanicis Zoologicisve usus sum informatione viva.

Scripta illius me docuerunt naturam rimari, et viginti ad summum horarum cum incomparabili Viro consortium mihi postea magis profuit, quam duorum annorum Collegia potuissent.

Ingenue equidem fateor, mihi commune cum plerisque, si non omnibus in Scientiis juvenibus fuisse fatum; quod scilicet Methodus Linneana variis in locis defectu laborare videbatur, quem supplere non erubuit tumida clanculum et audax juvenilis mens. Quantumne vero animum deprefferunt pauca, sed auro cariora pretiosissimaque illa momenta, quibus licuit cum Illustri hoc Sene colloqui. Didici me oscitanter judicasse, ipsas perfectiones pro naevis male habuisse, measque emendationes ad confusionem, quam adeo caute evitare studuit oculatissimus Vir, ducere. Utinam alii ingeniis suis non indulgerent, sed quae non recte intelligant, eorum prius resolutionem expectarent, quam emendanda cogitarent! Sane et sibi et Scientiae melius consulerent. Qui aliorum errores detegere methodosque recoquere tantum curae sibi habet, nunquam artis magister erit, sed tenebris obvolvitur, quae alii magno cum labore explicaverunt.

Genuini in arte Magistri criteria sunt, Tironum consulere simplici et titubanti conceptui, aliorum errores non venari, nisi fontes illorum detegere valebit, taciteque si quos deprehenderit corrigere, nisi contrarium poscant Scientiae progressus. Ejus fuit indolis jam Beatus A LINNE, illumque in his sequi debent omnes veri illius sectatores.

Nec

Nec Scientiis prohi. dolor! magis quam illustri nomine claris Familiis scandala defunt, qui omnem, ut aliorum merita obfuscent, operam impendant. Hisce vero indulgendum, exempla quoniam ingenii superbientis et amoris proprii delirantis, quos fuge! nobis offerunt.

Quod autem multo deterius est, variant Scientiarum status. Periodice enim quasi coluntur, protegentur, negliguntur, oblivione plane traduntur. En inconstantiae generis humani luculenta vestigia! Non est quod periodicae utilitati haec tribuas, potius Maecenatum gustui, nam omnis Scientia omni tempore utilis, licet non aequae adamanta et exulta. Ejusmodi variatio praecaveri et debet et potest. Debet: nam in Republica Scientiarum nulla negligenda; potest: si ad informationem juventutis tendentes res sollicito curentur, periti producantur, et doctrina atque experientia magis quam aetas, divitiae generisque nobilitas aestimentur.

Quando apta atque concinna Enchiridia defunt, necesse claudicet juvenum institutio, cum inepta, obscura diffusaque taedium apud Doctores et nauseam apud Discipulos excitent. Facile erit multa adducere exempla, laudatorum vero Virorum nominibus parcere fas est. Eo quo jam vivimus saeculo defectum nonnullorum, ineptitudinem aliorum hujus farinae scriptorum bene notum habemus, illis autem succurrere sat negligenter studemus. E nostris, qui in hoc studio maxime sane meruit, est Cel. Dn. C. D. GJÖRWELL, Assess. et Biblioth. Reg. Hic enim egregius vir non solum ma-

Scripta illius me docuerunt naturam rimari, et viginti ad summum horarum cum incomparabili Viro consortium mihi postea magis profuit, quam duorum annorum Collegia potuissent.

Ingenue equidem fateor, mihi commune cum plebisque, si non omnibus in Scientiis juvenibus fuisse fatum; quod scilicet Methodus Linneana variis in locis defectu laborare videbatur, quem supplere non erubuit tumida clanculum et audax juvenilis mens. Quantumne vero animum deprefferunt pauca, sed auro cariora pretiosissimaque illa momenta, quibus licuit cum Illustri hoc Sene colloqui. Didici me oscitanter judicasse, ipsas perfectiones pro naevis male habuisse, measque emendationes ad confusionem, quam adeo caute evitare studuit oculatissimus Vir, ducere. Utinam alii ingeniis suis non indulgerent, sed quae non recte intelligant, eorum prius resolutionem expectarent, quam emendanda cogitarent! Sane et sibi et Scientiae melius consulerent. Qui aliorum errores detegere methodosque recoquere tantum curae sibi habet, nunquam artis magister erit, sed tenebris obvolvitur, quae alii magno cum labore explicaverunt.

Genuini in arte Magistri criteria sunt, Tironum consulere simplici et titubanti conceptui, aliorum errores non venari, nisi fontes illorum detegere valebit, taciteque si quos deprehenderit corrigere, nisi contrarium poscant Scientiae progressus. Ejus fuit indolis jam Beatus A LINNE, illumque in his sequi debent omnes veri illius sectatores.

Nec

Nec Scientiis prohi deor! magis quam illustri nomine claris Familiis scandala defunt, qui omnem, ut aliorum merita obfuscent, operam impendant. Hisce vero indulgendum, exempla quoniam ingenii superbientis et amoris proprii delirantis, quos fuge! nobis offerunt.

Quod autem multo deterius est, variant Scientiarum status. Periodice enim quasi coluntur, protegentur, negliguntur, oblivione plane traduntur. En inconstantiae generis humani luculenta vestigia! Non est quod periodicae utilitati haec tribuas, potius Maecenatum gustui, nam omnis Scientia omni tempore utilis, licet non aequae adamanta et exulta. Ejusmodi variatio praecaveri et debet et potest. Debet: nam in Republica Scientiarum nulla negligenda; potest: si ad informationem juventutis tendentes res sollicitè curentur, periti producantur, et doctrina atque experientia magis quam aetas, divitiae generisque nobilitas aestimentur.

Quando apta atque concinna Enchiridia defunt, necesse claudicet juvenum institutio, cum inepta, obscura diffusaque taedium apud Doctores et nauseam apud Discipulos excitent. Facile erit multa adducere exempla, laudatorum vero Virorum nominibus parcere fas est. Eo quo jam vivimus saeculo defectum nonnullorum, ineptitudinem aliorum hujus farinae scriptorum bene notum habemus, illis autem succurrere sat negligenter studemus. E nostris, qui in hoc studio maxime sane meruit, est Cel. Dn. C. D. GJÖRWELL, Assess. et Biblioth. Reg. Hic enim egregius vir non solum ma-

num operi ipse admovit, sed alios quoque excitavit, elegantiorum scriptorum translationis curam gessit, suisque impensis, ut nostrae juventuti inserviant, plurima edidit: verbo, laudes et gratias de publico non tantas reportavit, quantas revera meruit.

Neminem, cui vel parum cognita sit Historiae Naturalis dignitas et utilitas, negaturum puto, Juventutem etiam in hoc Scientiarum genere in Scholis ac Gymnasiis institui debere. Hoc vero quominus fiat, obstat aptorum enchiridiorum defectus: Sive Plurimum Reverendi D. NATH. THENSTEDT Concion. Auh anno MDCCLXXI Introductionem ad Zoologiam *) conscripsi; Regnum Vegetabile eodem tractare modo mihi quidem proposuissem, propositum vero quoque lubenter deserui, quum in se laborem non satis jucundum suscepit Clar. D. Mag. A. ZETTER. Opusculum jam ante quinquennium absolutum esse scio, quid autem, quominus in publicum prodierit, effeceris, ignoro.

Deficiunt, ut nuper monui, Florae Svecicae Exemplaria, et, si vel argento parabilis esset Liber, tamen simul cum Generibus Plantarum ILL. A LINNE, nimis pretiosus ac Volumine incommodus Tironibus. Compendium igitur utriusque concinnavi et pretio et magnitudine commodum, ut spero. Cum vero Sveciae Daniacque regiones easdem fere generent Plantas; in una
denique

*) Inledning til Djur-Riket etc. Stockholm 1772. 8. med. figurer.

denique sensim novae detegantur, quarum copia intra alterius fines tantum fuerit observata: ut Danis aequae ac Svecis utilis fieret Libellus, omnes hactenus intra utriusque Regni limites detectas plantas in uno collegi opusculo; quodque idcirco *Prodromum Florae Scandinaviae* appellavi. Neminem moveat vox *Scandinaviae*; optime quidem notum mihi habeo, veteris Scandinaviae fines cum nostrae Florae non easdem esse, sed magis aptum non cognovi titulum, et hocce fine haerese uti me posse credidi. *Prodromi* titulo ideo usus sum, quod, si Deus vitam sanitatemque largiatur, majorem Floram propediem edere, in qua singulas Plantas, ea, quam egregius POLICH in *Historia Plantarum* fere, vel immortalis A. LINNÉ in *Mantissa altera* adhibuerunt, methodo describendas, constitui.

Ne praesenti opusculo deesset et concinnitas et utilitas,

Superflua evitaui Synonyma;

Plerumque idoneam citavi iconem;

Cuicunque Classi praemissi Characteres compendiosos Generum primarios;

Cuicunque Genere, denique addidi Characterem habituales methodo Cel. D. GOUAN, ad Species in opusculo occurrentes, applicatum, quod Tironibus praepriis gratum fore spero;

In clave Generum Species anomalas notavi, addito loco vero, quo facilius a Tirone inveniantur;

Rarissime observationes quasdam addidi;

Plantas Florae Danicae et Norvegicae ad veras reduxi Species; Signo ¶ illas notavi, quas nemo, quantum scio, intra Sveciae fines adhuc legit;

x
 Durationes ac Stationes vocibus abruptis vel unica falcata
 littera exposui, sequenti modo :

<i>a.</i> arena.	<i>nem.</i> nemora.
<i>acer.</i> acerofae fylvae.	<i>pal.</i> paludes.
<i>agr.</i> agri.	<i>pafc.</i> pascua.
<i>alp.</i> alpes.	<i>pisc.</i> piscinae.
<i>apr.</i> aprica.	<i>pr.</i> prata.
<i>aqu.</i> aquae.	<i>putr.</i> putrida.
<i>arb.</i> arbores.	<i>qu.</i> aquae quietae.
<i>arg.</i> argilla.	<i>r.</i> rudera.
<i>arv.</i> arva.	<i>riv.</i> rivuli.
<i>c.</i> culta.	<i>f.</i> sepes.
<i>coem.</i> coemiteria.	<i>fab.</i> fabulosa.
<i>cort.</i> cortices.	<i>ficc.</i> sicca.
<i>er.</i> ericeta.	<i>ster.</i> steriles.
<i>fluv.</i> fluvii.	<i>sydv.</i> fylvae.
<i>font.</i> fontes.	<i>terr.</i> terra.
<i>glar.</i> glareosa.	<i>tect.</i> tecta.
<i>hort.</i> horti.	<i>ul.</i> uliginosa.
<i>hum.</i> humida.	<i>umbr.</i> umbrosa.
<i>inund.</i> inundata.	<i>v.</i> viae.
<i>l.</i> littora.	<i>v. agr.</i> versurae agrorum.
<i>lac.</i> lacus.	<i>A.</i> annua planta.
<i>lap.</i> lapides.	<i>B.</i> biennis.
<i>lign.</i> ligna.	<i>P.</i> perennis.
<i>m.</i> mare.	<i>F.</i> frutex.
<i>mont.</i> montes.	<i>S.</i> suffrutex.
<i>mur.</i> muri.	<i>Arb.</i> arbor.

Hunc itaque Libellum non illis tantum, quibus
 volupe est ad Scholas et Universitates Sveciae Daniaeque
 Botanice incumbere, utile enchiridion, sed etiam Bo-
 tanicis exteris acceptum fore spero, cum illius ope,
 facil-

facillimo negotio Rev. GUNNERI *Flora Norvegica* et primis undecim Fasciculis *Florae Danicae* uti possint.

Optassem equidem hasce pagellas ut perlegisset antequam typis mandatae fuissent Magnus semper A LINNE. Multo emendatius sane, multo accuratius, forsitan etiam auctius prodidisset tunc opusculum. Hoc vero quominus fieret, cum recusaverit mors nunquam intermorituræ memoriae Herois, Amicos et nostros et Scientiae, ut errores forte commissos benigne non tantum excusent, sed etiam quo emendentur, indicent, implorare fas est.

Vale L. B. et Auctori fave. *Londini Gothorum*
die 1. Maji 1779.

A. J. R.

Quod plus quam tribus cœnuriis (362) Plantarum aucta prodeat haec Flora non mihi solum debetur, sed multa Amicis. Inter hoc praecipue Celeberrimis Viris ACHARIO Medico Provinciali in Ostrogothia, AFZELIO Demonstratori Upsaliensi, EHRHART Botanico Regio Hannoverano, HELLWIG Pharmacopoeo Stralsundensi, KÜLFIN Professore Stettinensi, SMITH Praesidi Societ. Linneanae, VAHL Professore Hafniensi, WESTRING Medico Regio laetus gratusque tribuo.

Studiose laboravi ut correctior evaderet libellus. Ea propter omnes, quos mihi videre licuit aliorum circa plantas enumeratas, observationes contuli, examinavi illisque pro viribus usus sum, propriasque addidi.

Chara-

Characterem secundarium vel habitualem Essentiali subjunxi, quum aptius uno in loco utrumque inveniri crediderim.

Systemati Linneano immutato quidem additus sum, at non coeca erga Auctorem veneratione, hinc in nonnullis ab eo recessi, et Muscos lubentissime secundum Methodum Hedwigianam ordinallem, si modo omnes enumeratos de novo examinare licuisset sicque suis generibus inferere. At Tironibus utile duxi appendicis loco elenchum addere Muscorum Hedwigianum, ab ipso egregio Auctore benevole revisum, ut reliquae species sensim generibus suis adjungi queat, atque sic haec Florae pars etiam perfici possit.

Lundae Gothorum 1795.

AUCTORES CITATI.

- A. H. X. C. FRIIS ROTTBÖLL Afhandling om en del rare Planter etc. in Vol. X. Actorum Scient. Hafniens. .
A. Holm. Kongl. Svenska Vetenskaps Academiens Handlingar.
A. Soc. Linn. Transactions of the Linnean Society V. I. London 1791. 4.
Act. Nid. Det Kongl. Norske Videnskabs Selskabs Skrifter. T. I—V. Hafniae 1761. etc. 8.
Berl. Besch. N. F. Fr. Beschäftigungen der Berlinischen Gesellschaft Naturforschender Freunde. Berlin 1775. etc. 8.
Cavan. Diss. A. J. CAVANTILLES Monadelphiae Classis Dissertationes X. Mattiti 1790. 4.
Clus. Hist. C. CLUSII Rariorum Plantarum Historia. Amstelredamiae 1601. fol.
Dickf. J. DICKSON Fasciculus Plantarum Cryptogamicarum Britanniae I, II. Londini 1785. 8.
Dill. Musc. J. J. DILLENII Historia Muscorum. Oxonii 1741. 4.
Ehrh. Beytr. FR. EHRHART Beyträge zur Naturkunde etc. Band 1—7. Hannover 1787. etc. 8.
Ehrh. Cal. ejusdem Calamariae, Gramina et Tetrapetaloidae. fol. Dec. I—X.
Ell. Cor. F. ELLIS Natural History of the Corallines. London 1755. 4.
Fl. D. Icones Florae Danicae. Hafniae. fol.

- Fl. Isl. *Flora Islandica*. Appendix ad Eggert Olafsen og
Biarne Povellens Rejse igiennem Island. Sorø 1772. 4.
- Fl. L. C. A. LINNÉ *Flora Lapponica* Amstel. 1737. 8.
- Fl. N. J. E. GUNNERI *Flora Norvegica*. T. I. II. Hafniae.
fol.
- Fl. Sv. C. A. LINNÉ *Flora Svecica*, Holmiae 1755. 8. Ed. 2.
- Fn. Sv. — — — *Fauna Svecica*. ibid. 1761. 8. Ed. 2.
- Gmel. Fuc. S. G. GMELIN *Historia Fucorum*. Petropoli
1768. 4.
- Gmel. Sib. F. G. GMELIN *Flora Sibirica*. T. I—IV. Petropoli
1747. etc. 4.
- H. Kew. W. AITON *Hortus Kewensis*. Vol. III. London
1789. 8.
- Hoffm. Lich. G. F. HOFFMANN *Descriptio et adumbratio Plan-
tarum e Classe Cryptogamica Linnei quæ Lichenes
dicuntur*. Vol. II. Lipsiae 1790. fol.
- Hoffm. Sal. *ejusdem* *Historia Salicum*. Lipsiae 1787. fol.
- Jacqu. Fl. Austr. N. J. JAQUIN *Florae Austriacae etc. icones*.
Vol. V. Viennae 1773. etc. fol.
- Jacqu. Vind. *ejusdem* *Enumeratio Stirpium etc. in agro Vin-
dobonensi etc. Vindobonae* 1762. 8.
- Leers Herb. F. D. LEERS *Flora Herbornensis* ed. alt. Berolini
1789. 8.
- Linn. Pl. rar. C. A. LINNÉ *Plantarum rariorum Horti Upsa-
liensis*. Fasc. I. Lipsiae 1767. fol.
- Morif. Hist. ROB. MORISONI *Historia Universalis Plantarum
Oxionensis*. T. III. fol.
- Murr. Pr. Gott. J. A. MURRAY *Prodromus Designationis Stir-
pium Gottingensium*. Gottingae 1770. 8.
- Müll. Fridr. O. F. MÜLLERS *Flora Fridrichsdalina*. Argente-
rati 1767. 8.
- M. Pl. C. A. LINNÉ *Mantissa Plantarum altera*. Holmiae
1771. 8.

- Mönch. Hall. C. MÖNCH Enumeratio Plantarum indigenarum
Hassiae praesertim inferioris. Cassel 1777. 8.
- N. Act. Svec. Konigl. Svenska Vetenskaps. Academiens Öf-
Handlingar. för år. 1780 etc. 8.
- Obf. Bot. Fasc. A. F. RETZII Observationes Botanicae sex
Fasciculis comprehensae. Lipsiae 1791. fol.
- Pluk. Phyt. L. PLUCKENETII Phytographia etc.
- Poll. J. A. POLLICH Historia Plantarum in Palatin. Elector.
nascentium. T. III. Manhemii 1776 etc. 8.
- Raji Angl.] J. RAJI Synopsis Methodica Stirpium. Londini
Raji Syn.] 1724. 8. Ed. 3.
- Ros. Obf. EB, ROSÉN Observationes Botanicae. Lond. Goth.
1749. 4.
- Roth. Germ. A. G. ROTHII Tentamen Florae Germanicae.
T. III. Lipsiae 1788. 8.
- Scheuchz. Agrost. J. SCHEUCHZERI Agrostographia. Tiguri
1719. 4.
- Seop. Carn. J. A. SCOPOLI Flora Carniolica. T. II. Ed. 2.
Viennae 1772. 8.
- Schreb. Gr. J. C. D. SCHREBERS Abbildung der Gräser. fol.
- Schreb. Lipf. — — — — Spicilegium Florae Lipsicae.
Lipsiae 1771. 8.
- Skr. Nat. Selfk. Skrivter af Natur Historie Selskabet. Kiöben-
havn 1790 etc. 8.
- Smith icon. J. E. SMITH Plantarum icones haecenus ineditae.
Fasc. III. Londini 1789. fol.
- S. N. C. A. LINNE' Systema Vegetabilium. Ed. XIV. Gottin-
gae 1784. 8.
- Sp. Pl. C. A. LINNE' Species Plantarum. T. II. Holmiae 1763.
Ed. 2. 8.
- St. Helv. ALB. A. HALLER Historia Stirpium indigenarum Hel-
vetiae T. I—III. Bernae 1768. fol.
- Uster ann. P. USTERI Annalen der Botanik. Stück I—7.
Zürich 1791 etc. 8.

Naill. Par. S. VAILLANT Botanicon Parisiense. Lugd. Bat.
1727. fol.

Web. spic. G. H. WERER Spicilegium Florae Gottingensis
Gothae 1778. 8.

Weig. Fl. C. E. WEIGEL Flora Pomerano-Rugica. Berolini
1769. 8.

Weig. Suppl. C. E. WEIGEL Supplementum Florae.

Weig. Obs. — — — 'Observationes Botanicae. Resp.
M. U. Wittich. Gryphiswaldiae 1772. 4.

Weis Crypt. F. G. WEIS Plantae Cryptogamicae Florae Got-
tingensis. Gottingae 1770. 8.

Willden Fl. C. L. WILLDENOW Florae Berolinensis Prodro-
mus. Berolini 1787. 8.

CORRIGENDA.

Pag. 1.	lin. 7	lege	HIPPURIS.
— 3	— 13	—	lacinia.
— 4	— 24	—	graminea.
— 11	— 7	—	longitudinaliter.
— 12	— 12	—	articulata.
— 19	— 15	—	disperſis.
— 20	— 11	—	repente.
—	— 16	—	capillari.
— 24	— 19	—	ſpiciformi.
— 34	— 24	—	geniculis.
—	— 30	—	In
— 35	— 15	—	retroſum
— 36	— 15	—	<i>Psyllium</i> .
— 41	— 73	—	tuberculata.
— 60	— 31	—	ut
— 63	— 32	—	plantas.
— 70	— 28	—	Holoſteo.
— 76	— 32	—	aufuga.
— 79	— 25	—	filiformis.
— 107	— 15	—	ſubbiſloris.
— 112	ſub <i>Agrimonia</i> adde		
	Rad. ramola. Caul. herbacea, ſtriſta.		
	Fol. pinnata, alterna, rugoſa.		
	Flor. ſpicata. Fulcr. ſtipulae.		
— 118	lin. 33	lege	Huc
— 120	— 34	—	indigena.
— 130	— 25	—	lanceolata.
— 133	— 2	—	emerſis.
— 144	— 3	—	<i>Criſta</i>
— 145	— 25	—	dubitanter.
— 150	— 2	—	ſtamina.
— 167	— 29	—	XXX annos docuit.
— 168	— 33	—	Penninis.
— 192	— 9	—	Linofyris.
— 195	— 31	—	Partille.
— 196	— 32	—	noſtram.
— 235	— 17	—	utrinque.
— 258	— 2	—	erectiusculis.
— 263	— 35	—	complicato
— 271	— 20	—	ſubtreſſellatis.
— 272	— 33	—	<i>Haematomma</i> .

Pag. 275	lin. 13	lège	infulis subrotundo-angulatis.
— 276	— 6	—	<i>Parella.</i>
— 284	— 23	—	lobatus.
— 300	— 29	—	tereti.
— 314	— 15	—	ovato.
— 340	— 43	—	<i>Holofchoenus.</i>
— 343	— 24	—	<i>Menyanthes.</i>
— 358	— 33	—	<i>Linofyris.</i>
— —	— 36	—	<i>Tanacetum.</i>
— 364	— 42	—	squamosa.
— 377	— 21	—	<i>Cynofurus.</i>

I. MONANDRIA.

I. MONOGYNIA.

SALICORNIA. *Cal.* 1-phyll. *Cor.* o. *Sem.* 1.

Rad. ramosa. *Caul.* articulata, succulenta, ramis oppositis. *Fol.* nulla. *Flor.* in articulis. *Fulcr.* o.

HIPPURIS. *Cal.* germini adnatus. *Cor.* o. *Stigma* simplex.

Rad. articulata, horizontalis. *Caul.* articulata, herbacea. *Fol.* verticillata. *Flor.* axillaris, sessilis. *Fulcr.* o.

Alchemilla Aphanes vide Tetrandria.

2. DIGYNIA.

CALLITRICHE. *Cal.* o. *Cor.* 2-pet. *Caps.* 2-locul. 4-sperma.

Rad. fibrosa. *Caul.* herbacea, filiformis, ramosa. *Fol.* sessilis, opposita, simplex. *Flor.* axillaris; pedunculata. *Fulcr.* o.

2 MONANDRIA. MONOGYNIA. Salicornia.

I. MONANDRIA.

I. M O N O G Y N I A.

SALICORNIA.

1. *S. herbacea*, herbacea patula, articulis apice compressis emarginato-bifidis. Fl. Sv. 1. Fl. D. 303. Fl. N. 274. A. l. m.
2. *S. fruticosa*, caule erecto fruticoso. S. N. 52. P. l. m. a).

HIPPURIS.

3. *H. vulgaris*, foliis octonis subulatis. S. N. 53. Fl. Sv. 2. Fl. N. 275. Fl. D. 87. P. riv. qu.
4. *H. maritima*, foliis lato-lanceolatis, inferioribus quaternis, superioribus quinis. *H. maritima*. Hell. Diff. c. fig. *H. lanceolata*. Obs. Bot. III. 7. t. 1. *H. tetraphylla*. Suppl. Pl. p. 81. P. mar. qu.

2. D I G Y N I A.

CALLITRICHE.

5. *C. verna*, foliis superioribus ovalibus, floribus androgynis. Fl. Sv. 3. Fl. N. 520. Fl. D. 129. A. aqu.
 6. *C. autumnalis*, foliis omnibus linearibus apice bifidis, floribus hermaphroditis. Fl. Sv. 4. Fl. N. 702. A. qu.
- Blitum Chenopodioides*. M. Pl. 2. p. 170. varietas tantum monstrosa *Bliti virgati* fuit.

- a) Crescit in Insula Hwen. Haec pedalis, dura, lignosa; illa vix semipedalis, succulenta, herbacea. Jungit *Hudson*; mihi diversa videtur, at vivam examinare non licuit. Legi Salicorniam tres antheras successive explicare; at ubi, oblitus; alios illam diandram observasse etiam vidi.

II. DIAN-

II. DIANDRIA.

I. MONOGYNIA.

LIGUSTRUM. *Cal* 4-dent. *Cor.* 4-fida. *Bacca* 4-sperma.
Rad. ramosa. *Caul.* fruticosa. *Fol.* simplex, opposita, integerrima. *Flor.* racemosa, terminalis. *Fulcr.* o.

CIRCAEA. *Cal.* 2-phyll. *Cor.* 2-pet. *Sem.* i. biloculare.
Rad. repens. *Caul.* herbacea, simplex. *Fol* simplex, opposita. *Flor.* racemosa, terminalis. *Fulcr.* o.

VERONICA. *Cal.* 4-5-partit. *Cor.* limbo 4-part. lacinia inferiore angustiore. *Capsf.* 2-locul.
Rad. fibrosa. *Caul.* ramosa, decumbens l. erecta. *Fol.* simplex, opposita l. terna, rarius alterna. *Flor.* varia. *Fulcr.* o.

GRATIOLA. *Cal.* 7-phyll. foliolis 2 exterior. patulis. *Cor.* irregul. resup. *Stam.* 2-steril. *Capsf.* 2-loc.
Rad. horizontalis, nodosa, fibrosa. *Caul.* angulata, herbacea. *Fol.* simplex, opposita, integra. *Flor.* solitaria, axillaris. *Fulcr.* o.

PINGUICULA. *Cal.* 2-labiat. 5-fid. *Cor.* ringens, calcarata. *Capsf.* 1-loc.
Rad. fibrosa. *Caul.* simplex, nuda. *Fol.* simplex, integra, radicalis, pinguis. *Flor.* solitaria, terminalis, nutans. *Fulcr.* o.

UTRICULARIA. *Cal.* 2-phyll. aequalis. *Cor.* ringens, calcarata. *Capsf.* 1-locul.
Rad. fibrosa. *Caul.* geniculata. *Fol* immersa, capillacea, multifida, vesiculis sparsis. *Flor.* racemosa, pedunculata. *Fulcr.* squamosa.

VERBENA. *Cal.* 5-dent. dente 1 truncato. *Cor.* infundib. subaequ. curva. *Stam.* 2-4. *Sem.* 2-4. *Rad.* ramosa. *Caul.* tetragona, ramosa. *Fol.* opposita, simplex. *Flor.* spicata, terminalis et axillaris. *Fulcr.* o.

LYCOPUS. *Cal.* 5-fidus. *Cor.* 4-fida. lacinia unica emarginata. *Stam.* distantia. *Sem.* 4 retusa. *Rad.* repens. *Caul.* tetragona, ramosa. *Fol.* opposita, simplex. *Flor.* axillaris, verticillata. *Fulcr.* o.

SALVIA. *Cal.* 2-lab. 5-dent. labio super. compresso. *Cor.* ringens. *Stam.* transverse pedicellata. *Sem.* 4. *Rad.* ramosa. *Caul.* ramosa, tetragona. *Fol.* opposita, simplex, rugosa, inferius petiolata, superior sessilis. *Fulcr.* bracteae, saepius coloratae.

Lepidium ruderale vide Tetradynamia Siliculosa.

2. DIGYNIA.

ANTHOXANTHUM. *Cal.* gluma 2-valv. 1-flor. *Cor.* gluma 2-valv. acuminata. *Sem.* 1. *Rad.* fibrosa. *Caul.* culmosa, articulata, simplex. *Fol.* gramminea. *Flor.* spicata, terminalis, nuda. *Fulcr.* o.

II. DIANDRIA.

I. MONOGYNIA.

LIGUSTRUM.

7. *L. vulgare*, foliis lanceolatis acutis paniculae pedicellis oppositis. S. N. 56. Fl. Sv. 5. F.

CIRCAEA.

8. *C. lutetiana*, caule erecto piloso, foliis ovatis subseratis. Obs. Bot. I. 9. Fl. Sv. 6. Fl. N. 320. Fl. D. 256. P. sylv.
 9. *C. alpina*, caule prostrato glabro, foliis cordatis dentatis. Obs. Bot. I. 9. Fl. Sv. 7. Fl. N. 436. Fl. D. 210. P. sylv. mont.

VERONICA.

* *Spicatae*.

10. *V. maritima*, spicis terminalibus, foliis ternis inaequaliter serratis. Fl. Sv. 8. Fl. N. 74. Fl. D. 374. Obs. Bot. I. 9. P. l. m.
 11. *V. longifolia*, spicis terminalibus, foliis oppositis lanceolatis serratis acuminatis. Fl. Sv. 9. P. pr. b).
 12. *V. spicata*, spicâ terminali, foliis oppositis crenatis obtusis, caule ascendente simplicissimo. Fl. Sv. 10. Fl. D. 52. P. a. c).
 13. *V. hybrida*, spicis terminalibus, foliis oppositis obtuse serratis scabris, caule erecto. Fl. Sv. 11. Raji Syn. 3. t. 11. f. 1. P.
 14. *V. officinalis*, spicis lateralibus pedunculatis, foliis oppositis, caule procumbente. Fl. Sv. 12. Fl. N. 46. Fl. D. 248. P. sylv.

A 3

** *Co-*

b) cfr. Afz. Obs. p. 23.

c) *Cel. Afzelius* epistola d. 22. Febr. 1782 ait, se hanc et spuriam ex eadem radice provenientes observasse. cfr. Diss. ejus de Veget. Svec.

* * *Corymboso-racemosae.*

- † 15. *V. fruticulosa*, corymbō terminali, foliis lanceolatis obtusiusculis crenatis, caulibus fruticulosis. S. N. 59. Fl. N. 829. Fl. D. 342. P. alp.
16. *V. alpina*, corymbō terminali, foliis oppositis, calycibus hispidis. Fl. Sv. 15. Fl. L. t. 9. f. 4. Fl. N. 45. Fl. D. 16. P. alp.
17. *V. serpyllifolia*, racemo terminali subspicato, foliis ovatis glabris crenatis. Fl. Sv. 16. Fl. N. 58. Fl. D. 492. Pr. pr.
18. *V. Beccabunga*, racemis lateralibus, foliis ovatis planis, caule repente. Fl. Sv. 14. Fl. N. 59. Fl. D. 511. P. qu. riv.
19. *V. Anagallis*, racemis lateralibus foliis lanceolatis serratis, caule erecto. Fl. Sv. 13. Fl. N. 1010. Fl. D. 903. A. qu.
20. *V. scutellata*, racemis lateralibus alternis, pedicellis pendulis, foliis linearibus integerrimis. Fl. Sv. 17. Fl. N. 658. Fl. D. 209. P. inund.
- † 21. *V. Teucrium*, racemis lateralibus longissimis, foliis ovatis rugosis dentatis obtusiusculis, caulibus procumbentibus. S. N. 59. P. umbr.
- † 22. *V. montana*, racemis lateralibus paucifloris, calycibus hirsutis, foliis ovatis rugosis crenatis petiolatis, caule debili. S. N. 59. P. umbr.
23. *V. Chamaedryf*, racemis lateralibus, foliis ovatis sessilibus rugosis dentatis, caule bifariam piloso. F. Sv. 18. Fl. N. 47. Fl. D. 448. P. pr.

* * * *Pedunculis unifloris.*

24. *V. agrestis*, floribus solitariis, foliis cordatis incisfis pedunculo brevioribus. Fl. Sv. 20. Fl. N. 954. Fl. D. 449. A. c.
25. *V. arvensis*, floribus solitariis, foliis cordatis incisfis pedunculo longioribus. Fl. Sv. 19. Fl. D. 515. A. c.

26. *V. hederaefolia*, floribus solitariis, foliis cordatis planis quinquelobis. Fl. Sv. 21. Fl. N. 1015. Fl. D. 428. A. c. r.
27. *V. triphyllus*, floribus solitariis, foliis digitato-partitis, pedunculis calyce longioribus. Fl. Sv. 22. Fl. D. 627. A. c. pr.
28. *V. verna*, floribus solitariis, foliis digitato-partitis, pedunculis calyce brevioribus. Fl. Sv. 23. Fl. N. 655. Fl. D. 252. A. ster.
29. *V. acinifolia*, floribus solitariis pedunculatis, foliis ovatis glabris crenatis, caule erecto subpiloso. S. N. 60. Ros. obs. p. 59. Vaill. Par. t. 33. f. 3. A. a.
30. *V. peregrina*, floribus solitariis sessilibus, foliis lanceolato-linearibus glabris obtusis subserratis, caule erecto. Fl. Sv. 24. Fl. D. 407. A. c.
- † 31. *marilandica*, floribus solitariis sessilibus, foliis linearibus, caulibus diffusis. S. N. 61. Fl. Isl. 3.

GRATIOLA.

- † 32. *Gr. officinalis*, foliis lanceolatis serratis, floribus pedunculatis. S. N. 64. Fl. D. 363. P. pr. pal.

PINGUICULA.

33. *P. vulgaris*, nectario cylindraceo longitudine petali. Fl. Sv. 25. Fl. N. 385. Fl. D. 93. P. ul.
34. *P. alpina*, nectario conico petalo brevior. Fl. Sv. 26. Fl. L. t. 12. f. 3. Fl. N. 640. t. 4. f. 4-6. Fl. D. 453. P. ul. alp.
35. *P. villosa*, scapo subviloso. Fl. Sv. 27. Fl. L. t. 12. f. 2. Fl. N. 641. P. ul. alp.

UTRICULARIA.

36. *U. vulgaris*, nectario conico, scapo paucifloro. Fl. Sv. 28. Fl. N. 683. Fl. D. 138. P. pal.
37. *U. minor*, nectario carinato. Fl. Sv. 29. Fl. D. 128. P. pal.

VERBENA.

38. *V. officinalis*, tetrandra, spicis filiformibus paniculatis, foliis multifido-laciniatis, caule solitario. Fl. Sv. 30. Fl. D. 628. A. r.

LYCOPUS.

39. *L. europaeus*, foliis sinuato-ferratis. Fl. Sv. 31. Fl. N. 173. P. riv.

SALVIA.

40. *S. pratensis*, foliis cordato-oblongis crenatis, summis amplexicaulibus, verticillis subnudis, corollis galea glutinosis. Fl. Sv. 32. P. pr.

2. D I G Y N I A.

ANTHOXANTHUM.

41. *A. odoratum*, spica oblonga ovata, flosculis subpedunculatis arista longioribus. Fl. Sv. 33. Fl. N. 5. Fl. D. 666. P. pr.

III. TRIANDRIA.

I. MONOGYNIA.

VALERIANA. *Cal.* o. *Cor.* 1-pet. 5-fida, basi hinc gibba. *Sem.* 1.

Rad. fibrosa. *Caul.* articulata, herbacea, fistulosa, simplex vel dichotoma. *Fol.* opposita, simplex vel composita. *Flor.* terminalis, corymbosa. *Fulcr.* o.

IRIS. *Cal.* spatha. *Cor.* 6-petaloidea alterne deflexa. *Styli* 3-petaloidei.

Rad. horizontalis, tereti-cárnosa, tuberosa. *Caul.* simplex, foliosa. *Fol.* ensiformis. *Flor.* sparsa, terminalis, spathacea. *Fulcr.* o.

SCHOENUS. *Cal.* glumae paleaceae univalves fasciculatim congestae. *Cor.* o. *Sem.* 1. subrotundum inter glumas.

Rad. fibrosa. *Caul.* culmosa, teres, farcta, foliosa. *Fol.* graminea. *Flor.* terminalis, spiculis capitatis. *Fulcr.* involucria subulata.

CYPERUS. *Cal.* glumae paleaceae distiche imbricatae. *Cor.* o. *Sem.* 1. nudum.

Rad. fibrosa. *Caul.* culmosa, simplex, farcta. *Fol.* graminea. *Flor.* umbellata spiculis. *Fulcr.* involucria laxa.

SCIRPUS. *Cal.* glumae paleaceae imbricatae. *Cor.* o. *Sem.* 1. setis brevibus paucioribus basi vel apice vel nullis.

Rad. fibrosa. *Caul.* culmosa, farcta, enodis. *Fol.* graminea vel cylindrica farcta. *Flor.* terminalis, simplex l. umbellata, spiculis teretibus. *Fulcr.* involucria.

ERIOPHORUM. *Cal.* glumae paleaceae undique imbricatae. *Cor.* o. *Sem.* 1. lana longissima cinctum.

Rad. fibrosa. *Caul.* culmosa, farcta. *Fol.* graminea. *Flor.* terminalis, spicata. *Fulcr.* involucrum.

NARDUS. *Cal.* o. *Cor.* 2-valv. *Sem.* tectum.

Rad. fibrosa. *Caul.* culmosa, foliosa, farcta, articulata. *Fol.* graminea. *Flor.* spicata, secunda. *Fulcr.* o.

2. DIGYNIA.

ASPRELLA. *Cal.* o. *Cor.* 2-valv. clausa. *Stam.* corolla breviora.

Rad. fibrosa. *Caul.* culmosa, articulata, foliosa. *Fol.* graminea. *Flor.* paniculata. *Fulcr.* o.

PHALARIS. *Cal.* 2-valv. carinatus, longitudine aequalis, *corollam* includens 1. florum.

Rad. fibrosa. *Caul.* culmosa, simplex, articulata, foliosa. *Fol.* graminea. *Flor.* paniculato-subspicata. *Fulcr.* o.

PANICUM. *Cal.* 3-valv. valvula tertia minima, 1-flora.

Rad. fibrosa. *Caul.* culmosa, simplex l. ramosa, articulata, foliosa. *Fol.* graminea. *Flor.* spicata, verticillata l. secunda. *Fulcr.* involuella fetosa.

PHLEUM. *Cal.* 2-valv. sessilis, linearis, truncatus apice 2-cuspidi, 1-flor. *Cor.* inclusa.

Rad. fibrosa l. bulbosa. *Caul.* culmosa, articulata, foliosa. *Fol.* graminea. *Flor.* spicata. *Fulcr.* o.

ALOPECURUS. *Cal.* 2-valv. 1-flor. *Cor.* 1-valv.

Rad. fibrosa. *Caul.* culmosa, articulata, foliosa. *Fol.* graminea. *Flor.* spicata. *Fulcr.* o.

MILIUM.

MILIUM. *Cal.* 2-valv. 1-flor. valvulis subaequalibus.
Cor. brevissima. *Stigm.* penicilliformis.

Rad. fibrosa. *Caul.* culmosa, articulata, foliosa.
Fol. graminea. *Flor.* paniculata. *Fulcr.* o.

AGROSTIS. *Cal.* 2-valv. 1-flor. corolla paulo minor.
Cor. 2-valv. *Stigm.* longitudinaliter hispida.

Rad. fibrosa. *Caul.* culmosa, articulata, foliosa.
Fol. graminea. *Flor.* paniculata. *Fulcr.* o.

AIRA. *Cal.* 2-valv. 2-flor. absque rudimento tertii
floris.

Rad. fibrosa. *Caul.* culmosa, articulata, foliosa.
Fol. graminea. *Flor.* paniculata. *Fulcr.* o.

MELICA. *Cal.* 2-valv. 1-2-florus, cum rudimento se-
cundi vel tertii.

Rad. fibrosa. *Caul.* culmosa, articulata. *Fol.*
graminea. *Flor.* secunda. *Fulcr.* ligulae va-
ginae.

POA. *Cal.* 2-valv. multiflorus. *Spiculae* ovatae, valvulis
acutiusculis margine scariosis.

Rad. fibrosa. *Caul.* culmosa, articulata, foliosa.
Fol. graminea. *Flor.* paniculata. *Fulcr.* o.

BRIZA. *Cal.* 2-valv. multiflorus. *Spiculae* distichae,
valvulis obtusis ventricosis.

Rad. fibrosa. *Caul.* culmosa, articulata. *Fol.*
graminea. *Flor.* paniculata. *Fulcr.* o.

DACTYLIS. *Cal.* 2-valv. multiflor. compressus, val-
vula altera carinata majore.

Rad. fibrosa. *Caul.* culmosa, articulata, foliosa.
Flor. paniculato-glomerata. *Fulcr.* o.

CYNOSURUS. *Cal.* 2-valv. multiflorus. *Involucrum*
1-phyll. laterale.

Rad. fibrosa. *Caul.* culmosa, articulata. *Fol.*
graminea. *Flor.* spicata, secunda. *Fulcr.*
bractea,

FESTUCA.

- FESTUCA.** *Cal.* 2-valv. multiflor. *Spiculae* oblongae, teretiusculae, glumis acuminatis.
Rad. fibrosa. *Caul.* culmosa; articulata. *Fol.* graminea. *Flor.* panicula varia. *Fulcr.* o.
- BROMUS.** *Cal.* 2-valv. multiflor. *Spiculae* oblongae, teretes, distichae, arista infra apicem.
Rad. fibrosa. *Caul.* culmosa, articulata. *Fol.* graminea. *Flor.* paniculata, spicata. *Fulcr.* o.
- STIPA.** *Cal.* 2-valv. 1-flor. *Cor.* valvula exterior arista terminali basi articulata.
Rad. fibrosa. *Caul.* culmosa, articuta. *Fol.* fetacea. *Flor.* paniculata. *Fulcr.* o.
- AVENA.** *Cal.* 2-valv. multiflor. arista dorsali contorta.
Rad. fibrosa. *Caul.* culmosa, articulata. *Fol.* graminea. *Flor.* paniculata. *Fulcr.* o.
- ARUNDO.** *Cal.* 2-valv. *Flosculi* congesti, basi lana cincti, mutici.
Rad. horizontalis, articulata. *Caul.* culmosa, articulata, inanis. *Fol.* graminea, rigida. *Flor.* paniculata. *Fulcr.* o.
- LOLIUM.** *Involucr.* 1-phyll. multiflor. fixum.
Rad. fibrosa. *Caul.* culmosa, articulata, foliosa. *Fol.* graminea. *Flor.* spicata, disticha. *Fulcr.* invol.
- ELYMUS.** *Involucr.* 2-phyll. sub quacunque spicula.
Rad. articulata. *Caul.* culmosa, articulata. *Fol.* graminea, rigida. *Flor.* spicata. *Fulcr.* invol.
- HORDEUM.** *Involucrum* 6-phyll. 3-flor.
Rad. fibrosa. *Caul.* culmosa, articulata. *Fol.* graminea. *Flor.* spicata. *Fulcr.* involucr.
- TRITICUM.** *Involucr.* 2-phyll. solitarium, multiflorum.
Rad. horizontalis, articulata. *Caul.* culmosa, articulata. *Fol.* graminea. *Flor.* spicata. *Fulcr.* invol.

3. TRIGYNIA.

MONTIA. *Cal.* 2-phyll. *Cor.* 1-pet. *Capsf.* 1-locul.
3-valv.

Rad. fibrosa. *Caul.* herbacea. *Fol.* opposita,
simplex. *Flor.* axillaris, terminalis, aggrega-
gata. *Fulcr.* bracteae.

HOLOSTEUM. *Cal.* 5-phyll. *Cor.* 5-pet. *Capsf.* 1-locul.
apice dehiscens.

Rad. ramosa. *Caul.* herbacea, geniculata. *Fol.*
opposita, simplex. *Flor.* umbellata, deflexa.
Fulcr. bracteae minutae.

KOENIGIA. *Cal.* 3-phyll. *Cor.* 0. *Sem.* 1. nudum.

Rad. simplex, perpendicularis. *Caul.* ramosa,
compressa, geniculata. *Fol.* simplex, integra,
varians. *Flor.* aggregata, terminalis. *Fulcr.*
stipulae vaginantes.

III. TRIANDRIA.

1. MONOGYNIA.

VALERIANA.

42. *V. officinalis*, floribus triandris, foliis omnibus pinnatis. Fl. Sv. 34. Fl. N. 157. Fl. D. 570. P. pr. hum.
43. *V. dioica*, floribus triandris dioicis foliis pinnatis integerrimis. Fl. Sv. 35. Fl. D. 687. Obs. Bot. I. 10. P. ul.
44. *V. locusta*, floribus triandris, caule dichotomo, foliis linearibus. Fl. Sv. 36. A.
 α . *olitoria*, fructu simplici. Fl. D. 738. c.
 β . *radiata*? involucro flores cingente. Act. Holm. 1769. p. 244. r.

IRIS.

- †45. *I. germanica*, barbata, foliis ensiformibus glabris falcatis brevioribus scapo multifloro. Syst. N. 89. Fl. N. 1032. P.
46. *I. pseudacorus*, imberbis, foliis ensiformibus, petalis alternis stylo minoribus. S. N. 90. Fl. Sv. 37. Fl. N. 264. Fl. D. 494. P. qu. fl.
47. *I. sibirica*, imberbis, foliis linearibus, scapo subtrifloro tereti, germinibus trigonis. S. N. 91. P.
- †48. *I. spuria*, imberbis, foliis linearibus, scapo subtrifloro tereti, germinibus hexagonis. S. N. 91. Fl. D. 91. P.

SCHOENUS.

49. *S. Mariscus*, culmo tereti, foliis margine dorsoque aculeatis, floribus subdiandris. Fl. Sv. 38. Fl. N. 1082. P. pal.
50. *S. nigricans*, culmo tereti nudo, capitulo ovato, involucri diphylli valvula altera subulata longa. Fl. Sv. 39. P. pal.

TRIANDRIA. MONOGYNIA. *Cyperus*. 15

51. *S. ferrugineus*, culmo tereti nudo, spica duplici, involucris valvula majore spicam aequante. Fl. Sv. 40. P. pal.
52. *S. fuscus*, culmo tereti folioso, spiculis subfasciculatis, foliis filiformibus canaliculatis. S. N. 95. P. pal.
53. *S. albus*, culmo subtriquetro folioso, floribus fasciculatis, foliis setaceis. Fl. Sv. 41. Fl. N. 824. Fl. D. 320. P. pal.
- Obs. Flores diandri Ehrh. B. IV. 154.

CYPERUS.

- † 54. *C. flavescent*, culmo triquetro nudo, umbella triphylla, pedunculis simplicibus inaequalibus, spicis confertis lanceolatis. S. N. 97. A. pr. pal.
- † 55. *C. fuscus*, culmo triquetro nudo, umbella trifida, pedunculis simplicibus inaequalibus, spicis confertis linearibus. S. N. 97. Fl. D. 179. Leers Herb. t. 1. f. 2. A. pr. hum.

SCIRPUS.

56. *S. palustris*, culmo tereti nudo, spica subovata terminali. Fl. Sv. 42. Fl. N. 546. Fl. D. 273. Leers Herb. t. 1. f. 3. P. l. qu. fl.
57. *S. Baeothrion*, culmo tereti striato nudo basi vaginato, vagina truncata, spica terminali pauciflora bivalvi: valvulis ovatis spica brevioribus margine scariosis: exteriore brevior. Suppl. Pl. 103. Ehrh. B. 2. 81. Fl. D. 167. *S. pauciflorus* Fl. Scot. 1078. P. pr. pal.
58. *S. caespitosus*, culmo striato nudo, spica bivalvi terminali longitudine calycis, radicibus squamula interstinctis. Fl. Sv. 43. Fl. N. 393. P. pal. ul.
59. *S. acicularis*, culmo tereti nudo setiformi, spicula ovata bivalvi, seminibus nudis. Fl. Sv. 44. Fl. N. 394. Fl. D. 387. P. inund.

† 60. *S.*

16 TRIANDRIA. MONOGYNIA. Eriophorum.

- † 60. *S. fluitans*, culmis teretibus nudis alternis, caule folioso flaccido. S. N. 98. Morif. III. p. 230. S. 8. t. 10. f. 31. d). P. inund.
- 61. *S. lacustris*, culmo tereti nudo, spicis ovatis pluribus pedunculatis terminalibus. Fl. Sv. 46. Fl. N. 948. P. fl. qu.
- † 62. *S. Holoschoenus*, culmo tereti nudo, spicis subglobosis glomeratis pedunculatis, involucro diphylo inaequali mucronato. S. N. 99. Fl. D. 454. P.
- 63. *S. setaceus*, culmo nudo setaceo, spica terminali sessili. Fl. Sv. 45. Fl. D. 311. Leers Herb. t. 1. f. 6. P. l. m.
- 64. *S. caricis*, culmo nudo subtriquetro, spica disticha, involucro monophyllo. Pr. Fl. Sc. 56. *Schoenus compressus*. S. N. 95. Obs. Bot. I. 10. N. Act. Holm. 1787. p. 243. Poll. 1. t. 1. f. 2. Leers Herb. t. 1. f. 1. P. ul. a. e).
- 65. *S. maritimus*, culmo triquetro, panicula conglobata foliacea, spicularum squamis trifidis: intermedia subulata. Fl. Sv. 47. Fl. N. 340. Fl. D. 937. P. l. m.
- 66. *S. sylvaticus*, culmo triquetro folioso, umbella foliacea, pedunculis nudis supradecompositis, spicis confertis. Fl. Sv. 48. Fl. N. 604. Fl. D. 707. Leers Herb. t. 1. f. 4. P. qu. sylv.

ERIOPHORUM.

- 67. *E. vaginatum*, culmis vaginatis teretibus, spica scariosa. Fl. Sv. 50. Fl. N. 659. Fl. D. 236. P. pal.

68. *E.*

d) Cel. Vahl.

e) Spica Caricis, culmus Schoeni, ad Pritzier Pomeraniae sesquipedalis.

68. *E. polytachyon*, culmis teretibus, foliis planis, spicis pedunculatis. Fl. Sv. 49. Fl. N. 174. Leers Herb. t. 1. f. 5. P. pal.
69. *E. alpinum*, culmis nudis triquetris, spica pappo brevior. Fl. Sv. 51. Fl. N. 711. Fl. D. 620. P. pal. alp.

NARDUS.

70. *N. stricta*, spica setacea recta secunda. Fl. Sv. 52. Fl. N. 221. Schr. Gr. t. 7. Leers Herb. t. 1. f. 7. P. pr. ster.

2. DIGYNIA.

ASPRELLA.

- †71. *A. oryzoides*. *Phalaris oryzoides*. S. N. 104. *Ehrhartia clandestina* Wigg. Hoff. 695. Schr. Gr. t. 22. P. pal.

PHALARIS.

72. *P. phleoides*, panicula cylindrica spiciformi glabra passim vivipara. Fl. Sv. 54. Fl. N. 465. Fl. D. 531. P. v. agr. ficc.
73. *P. arundinacea*, panicula oblonga ventricosa ampla. Fl. Sv. 53. Fl. N. 700. Fl. D. 259. Leers Herb. t. 7. f. 3. P. l. fl. qu.

PANICUM.

74. *P. verticillatum*, spica verticillata, racemulis quaternis, involucellis unifloris bifetis, culmis diffusis. S. N. 105. A. c.
- †75. *P. glaucum*, spica tereti, involucellis bifloris fasciculato-pilosis, seminibus undulato-rugosis. S. N. 105. Weig. Obs. p. 20. suppl. 873. A. c.

76. *P. viride*, spica tereti, involucellis bifloris fasciculato-pilosis, feminibus nervosis. S. N. 105. Leers Herb. t. 2. f. 2. *P. Crus Galli*. Fl. D. 852. A. c. a. f).
77. *P. Crus Galli*, spicis alternis conjugatisque, spiculis subdivisis, glumis aristatis hispidis, rachi quinquangulari. Fl. Sv. 55. Leers Herb. t. 2. f. 3. A. agr.
78. *P. sanguinale*, spicis digitatis basi interiore nodosis, flosculis geminis muticis, vaginis foliorum punctatis. S. N. 106. Fl. D. 388. Schr. Gr. t. 16. Leers Herb. t. 2. f. 6. A. agr.

PHLEUM.

79. *P. pratense*, spica cylindrica longissima ciliata, culmo erecto. Fl. Sv. 56. Fl. N. 60. Schr. Gr. t. 14. Leers Herb. t. 3. f. 2. P. pr.
80. *P. nodosum*, spica cylindrica, culmo adscendente, foliis obliquis, radice bulbosa. S. N. 108. Fl. N. 741. Fl. D. 380. Leers Herb. t. 3. f. 2. P. v. agr. g).
81. *P. alpinum*, spica ovato-cylindracea. Fl. Sv. 57. Fl. N. 150. Fl. D. 213. P. alp.
82. *P. arenarium*, spica ovata ciliata, culmo ramoso. Fl. Sv. 58. Fl. D. 915. A. a. h).

ALOPE-

- f) Hoc est gramen illud in Fl. Sv. sub nomine *Alopecuri panicæ* allatum, Alopecurus vero panicæ verus omnino e Scania exulat, et icon Fl. D. maxime dubia.
- g) Evanescit haud raro bulbosa radicum facies, at manet tamen ab antecedente distinctum omnino gramen, etiam cultum.
- h) Cum Phalaride phleoide jungitur in Horto Kewensi. Phalaridi potius quam Phleo annumerari forte debet, at species tum diversa.

ALOPECURUS.

83. *A. pratensis*, culmo spicato erecto, glumis villosis, corollis muticis. Fl. Sv. 59. Fl. N. 457. Schr. Gr. t. 19. Leers Herb. t. 2. f. 4. P. pr.
84. *A. agrestis*, culmo spicato erecto, glumis laevibus. S. N. 108. Weig. Suppl. 839. Fl. D. 697. Schr. Gr. t. 19. f. 2. Leers Herb. t. 2. f. 5. P. pr.
85. *A. geniculatus*, culmo spicato infracto, corollis muticis. Fl. Sv. 60. Fl. N. 605. Leers Herb. t. 2. f. 7. P. pr. inund.
- † 86. *A. paniceus*, panicula subspicata, glumis villosis, corollis aristatis. S. N. 108. Fl. D. 861. ? A. c.

MILIUM.

87. *M. effusum*, floribus paniculatis dispergis muticis. Fl. Sv. 61. Fl. N. 681. Leers Herb. t. 8. f. 7. P. sylv.

AGROSTIS.

* *aristatae*.

88. *A. spica venti*, petalo exteriori arista recta stricta longissima, panicula patula. Fl. Sv. 62. Fl. N. 916. Fl. D. 853. Leers Herb. t. 4. f. 1. A. agr.
89. *A. arundinacea*, panicula oblonga, petalo exteriori basi villosa, arista torta calyce longiore. Fl. Sv. 63. Fl. N. 1028. P. pr. ster.
90. *A. rubra*, paniculae parte florente patentissima, petalo exteriori glabro: arista terminali tortili recurva. Fl. Sv. 64. Fl. N. 712. P. pr. hum.
91. *A. canina*, calycibus elongatis, petalorum arista dorsali recurva, culmis prostratis subramosis. Fl. Sv. 65. Fl. N. 793. Leers Herb. t. 4. f. 2. P. pr. hum.
- β. *minor*, Fl. D. 161.

** *muticae*.

92. *A. stolonifera*, panicula ramulis patentibus muticis, culmo repente, calycibus aequalibus. Fl. Sv. 66. Fl. N. 1065. Fl. D. 564. P.
 α. erecta, foliis planis. arv.
 β. decumbens, foliis undulatis. a.
 γ. pumila, panicula mutica secunda, culmis fasciculatis erectis. S. N. III. i).
 †93. *A. alba*, panicula laxa, calycibus muticis aequalibus, culmo repetente. S. N. III. Leers Herb. t. 4. f. 5. P. pr.
 †94. *A. flava*, panicula patente, calycibus subulatis aequalibus hispidiusculis non coloratis, flosculis muticis dipetalis. Fl. D. 751. P. pr. apr. m.
 95. *A. capillaris*, panicula capillaris patente, calycibus subulatis aequalibus hispidiusculis coloratis, flosculis muticis monopetalis. S. N. III. N. Aët. Holm. 1787. p. 245. Fl. N. 458. Fl. D. 163. Leers Herb. t. 4. f. 3. P. pr. apr.

AIRA.

* *muticae*.

96. *A. aquatica*, panicula patente, floribus laevibus muticis calyce longioribus, foliis planis. Fl. Sv. 68. Fl. N. 742. Fl. D. 381. P. riv. fl. k).

** *aristatae*.

97. *A. subspicata*, foliis planis, panicula subspicata, flosculis medio aristatis: arista reflexa laxiore. Fl. Sv. 69. Fl. N. 422. Fl. D. 228. P. alp.

98. *A.*

- i) Est omnino ustilago Agr. stoloniferae, observante Cl. Ehrhart *Beyrr.* 4. 157.
 k) Similis omnino Poae, inprimis in ficcis, ubi calycibus 5-6 floris ludit.

98. *A. caespitosa*, foliis planis, panicula patente, petalis basi villosis aristatisque: arista recta brevi. Fl. Sv. 70. Fl. N. 262. Fl. D. 240. Leers Herb. t. 4. f. 8. P. pr.
99. *A. flexuosa*, foliis setaceis, culmis subnudis, panicula divaricata, pedunculis flexuosis. Fl. Sv. 71. Fl. N. 261. Fl. D. 157. Leers Herb. t. 5. f. 1. P. ster.
100. *A. montana*, foliis setaceis, panicula angustata; flosculis basi pilosis aristatis: arista tortili longiore. Fl. Sv. 72. Fl. N. 474. Leers Herb. t. 5. f. 2. P. mont.
101. *A. alpina*, foliis subulatis, panicula densa, flosculis basi pilosis aristatis: arista brevi. Fl. Sv. 73. Fl. N. 917. Fl. D. 961. P. alp.
102. *A. canescens*, foliis setaceis: summo spathaceo paniculam inferne obvolvente. Fl. Sv. 74. Fl. N. 918. B. a.
103. *A. praecox*, foliis setaceis: vaginis angulatis, floribus paniculato-spicatis, flosculis basi aristatis. Fl. Sv. 75. Fl. D. 383. A. a.
- † 104. *A. cariophyllaea*, foliis setaceis, panicula divaricata, floribus aristatis distantibus. S. N. 112. Fl. D. 382. Leers Herb. t. 5. f. 7. A. a.

MELICA.

105. *M. ciliata*, flosculi inferioris petalo exteriore ciliato. Fl. Sv. 77. P. ster.
106. *M. nutans*, petalis imberbibus, panicula nutante simplici, calycibus trifloris. Fl. Sv. 76. Fl. N. 463. Fl. D. 962. Schr. Gr. t. 6. Leers Herb. t. 3. f. 4. *M. montana*, Hudf. Fl. Angl. I. 37. P. fylv.
107. *M. uniflora*, panicula rara, calycibus bifloris: flosculo altero hermaphrodito: altero sterili. Obs. Bot. I. p. 10. *M. nutans*, Hudf. Fl. Angl. I. 37. P. fylv.

108. *M. coerulea*, panicula coarctata, floribus cylindricis. S. N. 113. *Aira coerulea*, Fl. Sv. 67. Fl. N. 259. Fl. D. 239. Leers Herb. t. 4. f. 7. P. ul.

POA.

109. *P. aquatica*, panicula diffusa, spiculis sexfloris linearibus. Fl. Sv. 78. Fl. N. 200. Fl. D. 920. Leers Herb. t. 5. f. 5. P. l. fl.
110. *P. alpina*, panicula diffusa ramosissima, spiculis sexfloris cordatis. Fl. S. 79. Fl. N. 600. P. alp.
 β. vivipara. Fl. D. 807.
111. *P. trivialis*, panicula diffusa, spiculis trifloris basi pubescentibus, culmo erecto tereti. Fl. Sv. 80. Fl. N. 476. Leers Herb. t. 6. f. 2. P. pr. pasc.
112. *P. angustifolia*, panicula diffusa, spiculis quadrifloris pubescentibus, culmo erecto tereti. Fl. Sv. 81. Fl. N. 507. Leers Herb. t. 6. f. 3. Pr. pr.
113. *P. pratensis*, panicula diffusa, spiculis quinquefloris glabris, culmo erecto tereti. Fl. Sv. 82. Fl. N. 456. Leers Herb. t. 6. f. 4. Pr. pr.
114. *P. annua*, paniculis diffusis angulis rectis, spiculis obtusis, culmo obliquo compresso. Fl. Sv. 83. Fl. N. 609. Leers Herb. t. 6. f. 1. A. pr.
115. *P. salina*, panicula diffusa, spiculis septemfloris linearibus, culmo basi subrepente, spicularum glumis omnibus obtusissimis. Poll. Pal. I. 92. Ehrhart Gram. Gr. 13. P. foss.
- † 116. *P. palustris*, panicula diffusa, spiculis subtrifloris pubescentibus, foliis subtus scabris. S. N. 113. Fl. D. 740. P. nem.
117. *P. compressa*, panicula coarctata secunda, culmo obliquo compresso. Fl. Sv. 84. Fl. N. 1004. Fl. D. 742. Vaill. t. 18. f. 8. Leers Herb. t. 5. f. 4. P. ficc.

118. *P. nemoralis*, panicula attenuata, spiculis subbifloris mucronatis scabris, culmo incurvo. Fl. Sv. 85. Fl. N. 1027. Fl. D. 749. Schèuchz. Agrost. t. 2. Leers Herb. t. 5. f. 3. P. sylv. nem.
119. *P. bulbosa*, panicula secunda patentiuscula, spiculis quadrifloris. S. N. 115. Fn. Sv. 557. A. vers. agr.
120. *P. arenaria*, paniculae subdiffusae spiculis sessilibus quadrifloris, culmis semiteretibus, foliis convolutis. *Poa maritima*, Fl. D. 251. *Aira aquatica*, β. Hudf. Angl. I. 34. *Poa cristata*, Fl. N. 780. exclusis synonymis praeter Raji. P. l. mar. V. ar.
121. *P. cristata*, panicula spicata, calycibus subpilosis subquadrifloris pedunculo longioribus, petalis aristatis. S. N. 115. Leers Herb. t. 5. f. 6. P. apr. mont.
- † 122. *P. glauca*, panicula patula secunda, spiculis subtrifloris, flosculis acuminatis, basi pubescentibus, foliis subulatis. Fl. D. 964. mont.

DACTYLIS.

123. *D. glomerata*, panicula secunda glomerata. Fl. Sv. 87. Fl. N. 478. Fl. D. 743. Schr. Gr. t. 8. f. 1. Leers Herb. t. 3. f. 3. P. pr. rud.

BRIZA.

124. *B. media*, spiculis ovatis, calyce flosculis (7) brevior. Fl. Sv. 86. Fl. N. 603. Fl. D. 258. B. pr.

CYNOSURUS.

125. *C. cristatus*, bracteis pinnatifidis. Fl. Sv. 88. Fl. N. 452. Fl. D. 238. Schr. Gr. t. 8. f. 2. Leers Herb. t. 7. f. 4. P. pr.
126. *C. coeruleus*, bracteis integris. Fl. Sv. 89. Fl. N. 610. *Sesteria coerulea* Ard. spec. 2. p. 18. t. 6. f. 3-5. P. pr. ster. ul.

FESTUCA.

127. *F. ovina*, panicula secunda coarctata aristata, culmo tetragono nudiusculo, foliis setaceis. Fl. Sv. 91. Fl. N. 453. Leers Herb. t. 8. f. 3. P. pr. ster.
 α. *mutica*, Leers Herb. t. 8, f. 3. *
 β. *vivipara*, Raji Syn. 3. t. 22. f. 1. alp.
128. *F. bromoides*, panicula secunda, spiculis erectis, calycis valvula altera minuta, altera acuminata. S. N. 118. Pluk. phyt. t. 33. f. 10. P. pr. ster.
129. *F. rubra*, panicula secunda scabra, spiculis sex-floris aristatis: flosculo ultimo mutico, culmo semitereti. Fl. Sv. 92. Fl. N. 581. Leers Herb. t. 8. f. 1. P. pr.
130. *F. duriuscula*, panicula oblonga secunda, spiculis oblongis laevibus, foliis setaceis. S. N. 118. Fl. D. 848. Leers Herb. t. 8. f. 2. P. pr. pasc. sicc.
- † 131. *F. dumetorum*, panicula spiciformis pubescente, foliis filiformibus. S. N. 118. Fl. D. 700. P. pr.
132. *F. elatior*, panicula secunda erecta, spiculis subaristatis: exterioribus teretibus. Fl. Sv. 94. Fl. N. 930. Schr. Gr. t. 2. Leers Herb. t. 8. f. 6. P. pr.
133. *F. decumbens*, panicula erecta, spiculis subovatis muticis, calyce flosculis majore, culmo decumbente. Fl. Sv. 93. Fl. N. 794. Fl. D. 162. Leers Herb. t. 7. f. 5. P. pasc. sicc.
134. *F. adscendens*, panicula erecta subramosa, spiculis teretibus angustis muticis, culmo adscendente, foliorum vaginis cylindricis. A. Holm. 1769. p. 245. B. pingv.
135. *F. fluitans*, panicula ramosa erecta, spiculis sessilibus teretibus muticis. Fl. Sv. 95. Fl. N. 409. Fl. D. 237. Schr. Gr. t. 3. Leers Herb. t. 8. f. 5. P. agr.

BROMUS.

136. *B. secalinus*, panicula nutante, spiculis ovato oblongis, aristis rectis, seminibus distinctis Fl. Sv. 96. Fl. N. 164. A. agr. ster.
α. vitiosus, Weig. Obf. p. 4. t. 1. f. 2. Leers Herb. t. 11, f. 2,
β. hordeaceus, Fl. Sv. 96. *β. Br. multiflorus*, Weig. Obf. p. 2. t. 1. f. 1.
137. *B. mollis*, panicula erectiuscula, spiculis ovatis pubescentibus: aristis rectis, foliis mollissime villosis. S. N. 119. Weig. Obf. p. 7. t. 1. f. 4. Schr. Gr. t. 6. Leers Herb. t. 2. f. 1. B. v. agr. sicc.
β. B. nanus, Weig. Obf. p. 8. t. 1. f. 9. in ficcis vulgatissimus, certe huic propior quam *Br. hordeaceo*, nec distincta species. An *mollis* a *hordeaceo* vere distinctus?
- † 138. *B. inermis*, panicula erecta, spiculis subteretibus subulatis nudis submuticis. S. N. 119. Schr. Gr. t. 3. P. 1).
139. *B. asper*, panicula nutante scabra, spiculis (subsexflores) villosis aristatis, foliis scabris. S. N. 119. *Bromus montanus*, Prodr. Fl. Scand. p. 19. Obf. Bot. Fasc. 2. p. 7. P. sylv.
140. *B. littoreus*, panicula erectiuscula scabra, spiculis oblongis subquadrifloris (4-6) glabris muticis. Prodr. Fl. Scand. p. 19. Weig. Suppl. 848. Gmel. Fl. Sibir. I. p. 110. t. XXI. et p. 111. ? St. Helv. 1511 m). P. lit. mar.

B 5

131. *B.*

1) Kölpin.

m) Maximus omnium. Figura ut et descriptio Gmelini optime quadrat, praeter villositatem. In Pomerania Cl. Weigel, in Scania Gen. á Engeström invenerunt.

141. *B. sterilis*, panicula patula, spiculis oblongis distichis, glumis subulato-aristatis. S. N. 120. Leers Herb. t. 11. f. 4. *Bromus grandiflorus*. Weig. Obsf. p. 9. t. 1. f. 6. B. ster.
142. *B. arvensis*, panicula nutante, spiculis ovato-oblongis. Fl. Sv. 97. Fl. N. 812. Fl. D. 293. Weig. Obsf. p. 6. t. 1. f. 3. Leers Herb. t. 11. f. 3. A. v. agr.
143. *B. tectorum*, panicula nutante, spiculis linearibus. Fl. Sv. 98. Fl. N. 586. Leers Herb. t. 10. f. 2. *Bromus grandiflorus*. Weig. Obsf. p. 10. β. A. ster.
144. *B. giganteus*, panicula nutante, spiculis quadrifloris aristis brevioribus. Fl. Sv. 99. Weig. Obsf. p. 11. t. 1. f. 5. Schr. Gr. t. 11. Fasc. Obsf. Bot. I. p. 11. Leers Herb. t. 10. f. 1. *Bromus triflorus*. Fl. D. 440. P. sylv.
- † 145. *B. triflorus*, panicula patente, floribus subtrifloris. S. N. 120. *Bromus patentissimus*. Weig. Obsf. p. 12. t. 1. f. 7. A. nem.
146. *B. pinnatus*, culmo indiviso, spiculis alternis subsessilibus teretibus subaristatis. Fl. Sv. 100. Fl. D. 164. Leers Herb. t. 10. f. 3. *Bromus sylvaticus*. Poll. I. 118. *Br. gracilis*. Weig. Obsf. p. 15. t. 1. f. 11. β. *Br. pinnatus*. Weig. Obsf. p. 14. t. 1. f. 10. Poll. I. 117. P. sylv.
- † 147. *B. cristatus*, spiculis distiche imbricatis sessilibus depressis. S. N. 120. Fl. Isl. 4. Gmel. Fl. Sib. I. t. 23. *Triticum cristatum*, Schr. Gr. t. 23. f. 2. P.

STIPA.

148. *S. pennata*, aristis lanatis. S. N. 121. Fn. Svec. 557. P. ster.

AVENA.

AVENA.

149. *A. elatior*, paniculata, calycibus bifloris, flosculo hermaphrodito mutico, masculo aristato. Fl. Sv. 102. Fl. N. 492. Fl. D. 165. Schr. Gr. t. 1. Leers Herb. t. 10. f. 4. P. nem.
150. *A. strigosa*, paniculata, calycibus bifloris, flosculis pedicellatis apice biaristatis, dorso arista geniculata. Schr. Lipf. p. 52. Obs. Bot. F. I. p. 11. *A. dubia*, Leers Herb. leg. t. 9. f. 3.? A. agr.
151. *A. fatua*, paniculata, calycibus trifloris, flosculis omnibus aristatis basi que pilosis. Fl. Sv. 161. Fl. N. 260. Schr. Gr. t. 15. Leers Herb. t. 9. f. 4. A. agr.
152. *A. flavescens*, panicula laxa, calycibus trifloris brevibus, flosculis omnibus aristatis. Fl. Sv. 103. Fl. N. 920. Schr. Gr. t. 9. Leers Herb. t. 10. f. 5. P. pr.
153. *A. pubescens*, subspicata, calycibus subtrifloris basi pilosis, foliis planis pubescentibus. S. N. 122. Leers Herb. 91. t. 9. f. 2. A. pr. ficc.
154. *A. pratensis*, subspicata, calycibus quinquefloris. Fl. Sv. 104. Fl. N. 716. Leers Herb. 92. t. 9. f. 1. P. pr. ster.

ARUNDO.

155. *A. phragmites*, calycibus quinquefloris, panicula laxa. Fl. Sv. 105. Fl. N. 222. Leers Herb. t. 7. f. 1. P. aqu. agr. a.
156. *A. epigejos*, calycibus unifloris, panicula erecta, foliis subtus scabris, culmo simpliciter. Fl. Sv. 106. P. ficc.
157. *A. neglecta*, calycibus unifloris, panicula lanceolata stricta, culmo ramoso. Ehrh. Beytr. 6. p. 137.
158. *A. Calamagrostis*, calycibus unifloris laevibus, corollis lanuginosis, culmo ramoso. Fl. Sv. 107. Fl. N. 486. Fl. D. 280. P. qu.

159. *A.*

159. *A. arenaria*, calycibus unifloris, foliis involutis mucronato-pungentibus. Fl. Sv. 108. Fl. Isl. 4. Fl. D. 917. P. a.

LOLIUM.

160. *L. perenne*, spica mutica, spiculis compressis multifloris. Fl. Sv. 110. Fl. N. 709. Fl. D. 747. Leers Herb. t. 12. f. 1. P. v. agr. viae.

β. *spica ramosa*, Leers Herb. t. 12. f. 1. inf.

γ. *aristatis spiculis*, Vaill. t. 17. f. 3.

161. *L. temulentum*, spica aristata, spiculis compressis multifloris. Fl. Sv. 109. Fl. N. 163. Fl. D. 160. A. agr.

β. *var. mutica*, Hush. Journ. 1778. Oct. p. 33.

ROTTBÛLLIA.

- † 162. *R. incurvata*, spica tereti subulata, gluma calycina subulata adpressa bipartita. S. N. 124. Fl. D. 938. P. apr.

ELYMUS.

163. *E. arenarius*, spica erecta arcta, calycibus tomentosis flosculo longioribus. Fl. Sv. 111. Fl. N. 199. P. a.

164. *E. caninus*, spica nutante arcta, spiculis rectis involucro destitutis: infimis geminis. Fl. Sv. 112. Leers Herb. t. 12. f. 4. 11. P.

HORDEUM.

165. *H. murinum*, flosculis lateralibus masculis aristatis, involucris intermediis ciliatis. Fl. Sv. 113. Fl. D. 629. B. ficc.

166. *H. fecalinum*, flosculis lateralibus masculis aristatis, involucris fetaceis scabris. Schr. Lipf. p. 148. Fl. D. 630. *H. murinum* β. Fl. Sv. 113. it. Scah. 214. B. pr. mar.

TRITI-

TRITICUM.

167. *T. repens*, calycibus quadrifloris subulatis acuminatis, foliis planis. Fl. Sv. 114. Fl. N. 201. Fl. D. 748. Leers Herb. t. 12. f. 3. P. pr. t. r.
 † 168. *T. junceum*, calycibus quinquefloris truncatis, foliis involutis. S. N. 127. Fl. D. 916. P. a. mar.

3. T R I G Y N I A.

MONTIA.

169. *M. fontana*, Fl. Sv. 115. Fl. N. 80. Fl. D. 1317).
 A. font.

HOLOSTEUM.

- † 170. *H. umbellatum*, floribus umbellatis. S. N. 128.
 Weig. Fl. 87. A. apr. arv.

KOENIGIA.

- † 171. *K. islandica*, S. N. 129. Fl. N. 1073. Fl. Isl. 4.
 Fl. D. 418. A. H. X. p. 427. t. 1. f. 1. A. inund. arg.

n) Variat calyce triphylo, staminibus 5, stylis 2.

IV. TETRANDRIA.

I. MONOGYNIA.

GLOBULARIA. *Cal. com. imbricatus, proprius tubulatus inferus. Cor. 2-lab. 5-part. Recept. paleaceum.*

Rad. ramosa. Caul. herbacea, foliosa. Fol. alternata, simplex. Flor. aggregato-capitata, terminalis. Fulcr. o.

DIPSACUS. *Cal. com. polyphyllus, proprius superus. Recept. conicum, paleaceum.*

Rad. fusiformis. Caul. angulata, scabra. Fol. opposita, connata, simplex. Flor. aggregato-capitata. Fulcr. o.

SCABIOSA. *Cal. com. polyphyllus, proprius duplex, superus. Sem. coronata.*

Rad. fibrosa. Caul. herbacea, foliosa. Fol. opposita, simplex et composita. Flor. aggregato-capitata. Fulcr. o.

SHERARDIA. *Cor. 1-pet. infundib. Sem. 2. tridentata.*

Rad. ramosa. Caul. herbacea, ramosa, tetragona, geniculata. Fol. verticillata. Flor. terminalis, fasciculata. Fulcr. involucrium.

ASPERULA. *Cor. 1-pet. infundib. Sem. 2. globosa.*

Rad. ramosa. Caul. herbacea, ramosa, geniculata, tetragona. Fol. verticillata. Flor. fasciculata. Fulcr. o.

GALIUM. *Cor. 1-pet. plana. Sem. 2. globosa.*

Rad. ramosa. Caul. herbacea, ramosa, geniculata, tetragona. Fol. verticillata. Flor. terminalis ramulorum, paniculata. Fulcr. o.

PLANTAGO. *Cal. 4-fid. Cor. 4-fid. limbo reflexo. Stam. longissima. Caps. 2-locul. circumscissa.*

Rad. fibrosa. Caul. herbacea, scaposa. Fol. simplex, radicalis. Flor. spicata. Fulcr. bracteae.

CENTUNCULUS. *Cal.* 4-fid. *Cor.* 4-fid. patens. *Stam.* brevia. *Capsf.* 1-locul. circumscissa.

Rad. fibrosa. *Caul.* herbacea, foliosa. *Fol.* simplex, alterna. *Flor.* solitaria, axillaris. *Fulcr.* o.

SANGUISORBA. *Cal.* 2-phyll. *Cor.* 4-partit. plana. *Capsf.* 4-gona inter calycem et corollam.

Rad. ramosa. *Caul.* herbacea, foliosa. *Fol.* alterna, pinnata. *Flor.* spicata. *Fulcr.* stipulae.

CORNUS. *Invol.* 4-phyll. *Cal.* 4-dent. *Cor.* 4-pet. supera. *Drupa* nucleo biloculari.

Rad. ramosa, lignosa. *Caul.* ramis, oppositis. *Fol.* simplex, opposita. *Flor.* corymbosa. *Fulcr.* o. 1. involucr.

TRAPA. *Cal.* 4-part. *Cor.* 4-pet. *Nux* spinis 4 oppositis.

Rad. fibrosa. *Caul.* submersa. *Fol.* submersa, capillacea, multifida: emersa, simplex, subrotunda, natans. *Flor.* terminalis demum spicata. *Fulcr.* o.

ISNARDIA. *Cal.* 4-fid. *Cor.* o. *Capsf.* 4-locul. calyce cincta.

Rad. fibrosa. *Caul.* natans, geniculata radicans. *Fol.* simplex, opposita. *Flor.* axillaris, solitaria. *Fulcr.* o.

ALCHEMILLA. *Cal.* 8-fid. *Cor.* o. *Sem.* 1. nudum.

Rad. ramosa. *Caul.* herbacea, foliosa. *Fol.* subpalmat. *Flor.* termin. 1. axillaris, glomerata, non colorata. *Fulcr.* stipulae vaginantes.

Convallaria bifolia vid. Hexandr. Monog.

Evonymus europaeus vid. Pentandr. Monog.

Cardamine hirsuta vid. Tetradyn. filiqu.

Verbena officinalis vid. Diandria Monog.

TETRANDRIA.

2. DIGYNIA.

CUSCUTA. *Cal.* 4 fid. *Cor.* 4-fid. *Capf.* 2-loc. circumscissa.

Rad. parasitica caulium. *Caul.* filiformis, volubilis. *Fol.* o. *Flor.* glomerata, sessilis.

Fulcr. bracteae squamiformes.

Gentianae quadrifidae vid. Pentandr. Digyn.

3. TETRAGYNIA.

ILEX. *Cal.* 4-dent. *Cor.* rotata, 4-part. *Styl.* o. *Bacca* 4-sperma.

Rad. lignosa, ramosa. *Caul.* arborea. *Fol.* alterna, simplex. *Flor.* axillaris, glomerata.

Fulcr. o.

POTAMOGETON. *Cal.* o. *Cor.* 4-pet. *Styl.* o. *Sem.* 4.

Rad. articulata, horizontalis. *Caul.* herbacea, articulata, demersa. *Fol.* simplex, varia.

Flor. spicata. *Fulcr.* stipulae.

RUPPIA. *Cal.* o. *Cor.* o. *Sem.* 4. pedicellata.

Rad. fibrosa. *Caul.* herb. articulata, demersa, *Fol.* simplex, alterna, sessilis. *Flor.* pedunculata, solitaria. *Fulcr.* stipulae vagin.

SAGINA. *Cal.* 4-phyll. *Cor.* 4-pet. l. o. *Capf.* 1-locul. 4-valv. polysperma.

Rad. fibrosa. *Caul.* herbacea, articulata. *Fol.* simplex, opposita. *Flor.* solitaria. *Fulcr.* o.

TILLAEA. *Cal.* 4-part. *Cor.* 4-pet. *Capf.* 4. polyspermae.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex, sessilis, sparsa. *Flor.* axillaris, solitaria, sessilis.

Fulcr. o.

Linum Radiola v. Pentandr. Pentagyn.

Thesium alpinum v. Pentandr. Monogyn.

IV. TETRANDRIA.

I. MONOGYNIA.

GLOBULARIA.

172. *G. vulgaris*, caule herbaceo, foliis radicalibus tridentatis: caulinis lanceolatis. Fl. Sv. 116. P. apr.

DIPSACUS.

- † 173. *D. sylvestris*, foliis integris, paleis rectis. S. N. 143. Fl. D. 965. B. rud.
 † 174. *D. pilosus*, petiolis appendiculatis. S. N. 143. Weig. Suppl. 824. P.

SCABIOSA.

175. *S. succisa*, corollulis quadrifidis aequalibus, caule simplici, ramis approximatis, foliis lanceolato-ovatis. Fl. Sv. 119. Fl. N. 187. Fl. D. 279. P. pr.
 176. *S. arvensis*, corollulis quadrifidis radiantibus, foliis pinnatifidis incisfis, caule hispido. Fl. Sv. 117. Fl. N. 186. Fl. D. 447. P. pr.
 177. *S. columbaria*, corollulis quinquefidis radiantibus, foliis radicalibus ovatis crenatis: caulinis pinnatis fetaceis. Fl. Sv. 118. Fl. N. 726. Fl. D. 314. *Scab. polymorpha*, Weig. Obs. p. 23. P. sicc.

SHERARDIA.

178. *S. arvensis*, foliis omnibus verticillatis, floribus terminalibus. Fl. Sv. 120. Fl. N. 1018. Fl. D. 439. A. c.

ASPERULA.

179. *A. odorata*, foliis octonis lanceolatis, florum fasciculis pedunculatis. Fl. Sv. 121. Fl. N. 270. Fl. D. 562. P. sylv.

34 TETRANDRIA. MONOGYNIA. Galium.

180. *A. tinctoria*, foliis linearibus: inferioribus senis: intermediis quaternis. caule flauido, floribus plerisque trifidis. Fl. Sv. 122. P. sicc.

GALIUM.

181. *G. palustre*, foliis quaternis obovatis inaequalibus, caulibus diffusis. Fl. Sv. 126. Fl. N. 931. Fl. D. 423. P. ul.
182. *G. trifidum*, foliis quaternis linearibus, caule procumbente scabro, corollis trifidis. S. N. 150. N. Act. Holm. 1789. p. 40. Fl. N. 788. Fl. D. 48.
183. *G. uliginosum*, foliis senis lanceolatis retrorsum ferrato - aculeatis mucronatis rigidis, corollis fructu maioribus. Fl. Sv. 127. Fl. N. 78. P. hum.
184. *G. saxatile*, foliis senis obovatis obtusis, caule ramosissimo procumbente. Fl. Sv. 1294.
- † 185. *G. pomeranicum*, foliis lineari - lanceolatis scabris: infimis octonis: superioribus senis: supremis quaternis, caulibus tetragono - angulatis o). P.
186. *G. verum*, foliis octonis linearibus fulcatis, ramis floriferis brevibus. Fl. Sv. 123. Fl. N. 105. P. v. agr. pr.

187. *G.*

- o) *Radix perennis*. *Caules* rectiusculi, tetragoni, angulis marginatis, laeves, geniculi inferne incrassatis. *Rami* breves patentissimi. *Folia* lineari - lanceolata, mucronata, subtus pallidiora carina pubescente, margine tenuissime serrulata, plerumque reflexa. *Panicula* ramis oppositis, nonnunquam quaternis, subtrichotomis. *Pedicelli* versus germen incrassati. *Corolla* alba, laciniis acutis in medio linea exaratis. *Stigmata* globosa. *Fructus* ovatus, laevis. I Pomerania legi, locis nī fallor arenosis crescens. An *Galium scabrum* Jacqu, vel *Mollugo Rqth.* fl. Germ. 2. p. 183?

187. *G. Mollugo*, foliis octonis ovato-linearibus subferratis patentissimis mucronatis, caule flauido, ramis patentibus. Fl. Sv. 125. Fl. D. 455. P. r.
 β. *angustifolium*. Roth. Fl. Germ. I. p. 65. 2.
 p. 182. Weig. Fl. 826.
- †188. *G. sylvaticum*, foliis octonis laevibus subtus scabris: floralibus binis, pedunculis capillaribus, caule laevi. S. N. 150. *Gal. glaucum*. Fl. D. 609. P. sylv.
189. *G. boreale*, foliis quaternis lanceolatis trinerviis glabris, caule erecto, feminibus hispidis. Fl. Sv. 124. Fl. N. 104. P. ster.
190. *G. Aparine*, foliis octonis lanceolatis carinatis scabris retrosum aculeatis, geniculis villosis, fructibus hispidis. Fl. Sv. 128. Fl. N. 155. Fl. D. 495. A. f.

PLANTAGO.

191. *P. major*, foliis ovatis glabris, scapo tereti, spica flosculis imbricatis. Fl. Sv. 129. Fl. N. 389. Fl. D. 461. P. v. pr.
192. *P. media*, foliis ovato-lanceolatis pubescentibus, spica cylindrica, scapo tereti. Fl. Sv. 130. Fl. N. 345. Fl. D. 581. P. pr.
193. *P. lanceolata*, foliis lanceolatis, spica subovata nuda, scapo angulato. Fl. Sv. 131. Fl. N. 645. Fl. D. 437. B. v. agr.
 β. *Pl. altissima*, foliis lanceolatis quinquenerviis dentatis glabris, spica oblongo-cylindrica, scapo angulato. S. N. 155 p).
 γ. *Pl. capitata*, spica capitata q).

C 2

194. *P.*

p) Confer Cel. Afzelii Diss. de Plantis Svec. p. 26.

q) Misit et hanc Cel. Afzelius, praeter spicae formam, similitudinem *altissimae*.

36 TETRANDRIA. MONOGYNIA. Centunculus.

194. *P. maritima*, foliis semicylindraceis integerrimis basi lanatis, scapo tereti. Fl. Sv. 133. Fl. N. 198. Fl. D. 243. B. a. m.

β. *Plantago squamata*. Fl. D. 691.

γ. *Pl. serpentina*, foliis linearibus planis glabris basi lanatis, spica erecta cylindrica longiore, scapo tereti hirsuto pilis adpressis. A. Holm. 1769. p. 246. *Pl. dubia*. Fl. Sv. 132. Obs. Bot. Fasc. I. p. 11.

195. *P. ferraria*, foliis lanceolatis quinquenerviis dentato-ferratis, scapo tereti. S. N. 156. r). P.

196. *P. coronopifolia*, foliis linearibus dentatis, scapo tereti. Fl. Sv. 143. Fl. D. 272. A. a. m.

- † 197. *P. Phyllium*, caule ramoso herbaceo, foliis subdentatis recurvatis, capitulis aphyllis. S. N. 156. s). A.

CENTUNCULUS.

198. *C. minimus*, Fl. Sv. 136. Fl. N. 390. Fl. D. 177. Anagallis Centunculus. Afz. Diff. p. 1. A. a. ficc.

SANGUISORBA.

199. *S. officinalis*, spicis ovatis. Fl. Sv. 137. Fl. N. 790. Fl. D. 97. P. pr. ficc.

CORNUS.

200. *C. sanguinea*, arborea, cymis nudis, ramis rectis. Fl. Sv. 138. Fl. D. 481. F. nem.

201. *C. Suecica*, herbacea, ramis binis. Fl. Sv. 139. Fl. L. t. 5. f. 3. Fl. N. 300. Fl. D. 5. P. pr. mont.

ISNAR-

r) Auctoritate Cel. Afzelii in lit.

s) Cel. Kölpin.

• TETRANDRIA. MONOGYNIA. Isnardia. 37

ISNARDIA.

202. *I. palustris*. S. N. 162. A. pal.

TRAPA.

203. *T. natans*, nucibus quadricornibus: spinis patentibus. S. N. 163. Fl. Sv. 141. P. lac.

ALCHEMILLA.

204. *A. vulgaris*, foliis lobatis. Fl. Sv. 141. Fl. N. 192. Fl. D. 693. P. pr.

205. *A. alpina*, foliis digitatis ferratis. Fl. Sv. 142. Fl. N. 192. Fl. D. 49. P. alp.

206. *A. Aphanes*, foliis trilobatis, floribus monandris. Afz. Diff. p. 2. *Aphanes arvensis*. Fl. Sv. 143. Fl. N. 1013. Fl. D. 973. A. arv.

2. D I G Y N I A.

CUSCUTA.

207. *C. europaea*, floribus sessilibus confertis quadridis. Fl. Sv. 144. Fl. N. 639. Fl. D. 199. A.

208. *C. Epithymum*, floribus sessilibus solitariis quinquefidis bracteis obvallatis. S. N. 167. Fl. D. 427. A.

3. T E T R A G Y N I A.

ILEX.

- †209. *I. Aquifolium*, foliis ovatis acutis spinosis. S. N. 168. Fl. N. 518. Fl. D. 508. F.

POTAMOGETON.

- 210. *P. natans*, foliis oblongo-ovatis petiolatis nantibus. Fl. Sv. 145. Fl. N. 399. P. fl. lac.
- 211. *P. perfoliatum*, foliis cordatis amplexicaulibus. Fl. Sv. 146. Fl. D. 196. P. fl.
- 212. *P. lucens*, foliis lanceolatis planis in petiolos definentibus. Fl. Sv. 147. Fl. N. 944. Fl. D. 195. P. fl. lac.
- 213. *P. crispum*, foliis lanceolatis alternis oppositisve undulatis ferratis. Fl. Sv. 148. Fl. D. 927. P. qu.
- 214. *P. compressum*, foliis linearibus obtusis, caule compresso. Fl. Sv. 149. Fl. D. 203. P. qu.
- 215. *P. pectinatum*, foliis setaceis parallelis approximatis distichis. Fl. Sv. 150. P. qu.
- 216. *P. gramineum*, foliis lineari-lanceolatis alternis sessilibus stipula latoribus. Fl. Sv. 151. Fl. D. 222. P. qu.
- 217. *P. marinum*, foliis linearibus alternis distinctis inferne vaginantibus. Fl. Sv. 152. Fl. D. 186. P. mar. qu.
- 218. *P. fusillum*, foliis linearibus oppositis alternisque distinctis basi patentibus, caule tereti. Fl. Sv. 153. P. qu.

RUPPIA.

- 219. *R. maritima*, Fl. Sv. 154. Fl. N. 731. Fl. D. 364. Pl. l. m.

SAGINA.

220. *S. procumbens*, ramis procumbentibus, floribus plerumque apetalis. Fl. Sv. 155. Fl. N. 505. Fl. D. 845. A. ster. ficc.
- † 221. *S. erecta*, caule erecto subunifloro. S. N. 169. Fl. D. 845. Vaill. Par. t. 3. f. 2.

TILLAEA.

222. *T. aquatica*, erecta dichotoma, foliis acutis, floribus quadrifidis. Fl. Sv. 156. Fl. N. 1074. A. H. X. p. 429. A. inund.

V. PENTANDRIA.

I. MONOGYNIA.

MYOSOTIS. *Cor.* hypocraterif. 5-fid. emarginata: fauce clausa fornicibus. *Sem.* 4.

Rad. fibrosa. *Caul.* herbacea, teres. *Fol.* sessilis, alterna, hirta, simplex. *Flor.* terminalis, racemosa. *Fulcr.* o.

LITHOSPERMUM. *Cor.* infundib. fauce perforata, nuda. *Cal.* 5-part. *Sem.* 4.

Rad. ramosa. *Caul.* herbacea, teres. *Fol.* alterna, simplex, sessilis. *Flor.* solitaria. *Fulcr.* o.

ANCHUSA. *Cor.* infundib. fauce clausa fornicibus. *Sem.* 4. basi insculpta.

Rad. ramosa. *Caul.* herbacea, teres, ramosa. *Fol.* alterna, simplex, sessilis. *Flor.* terminalis, spicata. *Fulcr.* o.

CYNOGLOSSUM. *Cor.* infundib. fauce clausa fornicibus. *Sem.* 4. depressa, interiore tantum latere stylo affixa.

Rad. ramosa, crassa. *Caul.* herbacea, ramosa. *Fol.* alterna, simplex, sessilis, mollis. *Flor.* terminalis, ramosa. *Fulcr.* o.

PULMONARIA. *Cor.* infundib. fauce pervia. *Cal.* prismatico-pentagonus. *Sem.* 4.

Rad. ramosa. *Caul.* herbacea, teres, foliosa. *Fol.* alterna, simplex, punctata. *Flor.* paniculata, terminalis. *Fulcr.* o.

SYMPHYTUM. *Cor.* limbus tubulato-ventricosus: fauce clausa radiis lanceolatis. *Sem.* 4.

Rad. fusiformis. *Caul.* herbacea, angulata. *Fol.* simplex, alterna, decurrens. *Flor.* subracemosa. *Fulcr.* o.

ASPERUGO. *Cor.* infundib. *Cal.* fructus compressus: lamellis plano-parallelis sinuatis. *Sem.* 4.

Rad. ramosa. *Caul.* herbacea, ramosa, debilis.

Fol. simplex, sessilis, sparsa. *Flor.* axillaris, congesta, subsessilis, sparsa. *Fulcr.* 0.

LYCOPSIS. *Cor.* infundib. tubo curvo. *Sem.* 4.

Rad. ramosa. *Caul.* herbacea, ramosa, fetosa.

Fol. simplex, alterna, sessilis. *Flor.* solitaria, pedunculata. *Fulcr.* 0.

ECHIUM. *Cor.* campanulata, irregul. fauce nuda. *Sem.* 4.

Rad. ramosa. *Caul.* herbacea, teres, tuberculata. *Fol.* simplex, sessilis, alterna. *Flor.* spicata, secunda. *Fulcr.* 0.

DIAPENSIA. *Cal.* 8-phyll. 3. exterioribus. *Cor.* hypocraterif. *Stam.* tubo corollae imposita. *Capsf.* 3-locul. 3-valv.

Rad. fibrosa. *Caul.* ramosa, foliosa. *Fol.* simplex, sessilis, imbricata. *Flor.* solitaria, terminalis. *Fulcr.* 0.

ANDROSACE. *Umbella* involucrata. *Cor.* tubus ovatus, ore glanduloso. *Capsf.* 1-locul. glob. apice 5-fido dehiscens.

Rad. simplex. *Caul.* scaposa. *Fol.* simplex, radicalis. *Flor.* terminalis, umbellata. *Fulcr.* involucreum.

PRIMULA. *Umb.* invol. *Cor.* tubus cylindricus ore patulo. *Capsf.* teres 1-loc. apice 10-dent. dehiscens, polysperma.

Rad. ramosa. *Caul.* scaposa. *Fol.* simplex, radicalis. *Flor.* terminalis, umbellata. *Fulcr.* involucreum.

MENYANTHES. *Cor.* hirsuta. *Stigma* 2-fid. *Capsf.* 1-loc.

Rad. horizontalis, articulata. *Caul.* herbacea, demersa. *Fol.* ternata l. simplex. *Flor.* scaposa, axillaris. *Fulcr.* bracteae.

HOTTONIA. *Cor.* hypocraterif. *Stam.* tubo corollae imposita. *Capsf.* 1-loc.

Rad. fibrosa. *Caul.* demersa, repens, radicans, foliosa. *Fol.* demersa, verticillata, multifida. *Flor.* spicato-verticillata. *Fulcr.* bracteae.

LYSIMACHIA. *Cor.* rotata. *Capsf.* globosa, mucronata, 10-valv.

Rad. repens. *Caul.* herbacea, ramosa. *Fol.* simplex, opposita l. verticillata. *Flor.* varia. *Fulcr.* o.

ANAGALLIS. *Cor.* rotata. *Capsf.* circumscissa.

Rad. fibrosa. *Caul.* herbacea, debilis, tetragona. *Fol.* simplex, opposita, sessilis. *Flor.* axillaris, solitaria, pedunculata. *Fulcr.* o.

AZALEA. *Cor.* campanulata. *Stam.* receptaculo inserta. *Capsf.* 5-locul.

Rad. lignosa, fibrosa. *Caul.* fruticulosa, ramosa, foliosa. *Fol.* simplex, conferta, semper-virens. *Flor.* terminalis, subterna, pedunculata. *Fulcr.* o.

CONVOLVULUS. *Cor.* campanulata, plicata. *Stigm.* 2. *Capsf.* 2-loc. 4-sperma.

Rad. repens. *Caul.* herbacea, foliosa, volubilis. *Fol.* simplex, alterna. *Flor.* axillaris, pedunculata. *Fulcr.* bracteae.

POLEMONIUM. *Cor.* 5-part. fundo clauso valvis staminiferis. *Stigm.* 3-fid. *Capsf.* 3-loc. supera.

Rad. fibrosa. *Caul.* herbacea, striata, foliosa. *Fol.* alterna, sessilis, pinnata. *Flor.* racemosa. *Fulcr.* o.

CAMPANULA. *Cor.* campanulata fundo clauso valvis staminiferis. *Stigm.* 3-fid. *Capsf.* infera, poris lateralibus dehiscens.

Rad. ramosa, tuberosa, fibrosa. *Caul.* herbacea, foliosa. *Fol.* simplex, varia. *Flor.* varia. *Fulcr.* bracteae.

PHYTEUMA. *Cor.* rotata, 5-part. laciniis linearibus. *Stam.* reflexa. *Stigma* 2-fid. *Capsf.* 2-locul. infera.

Rad. fusiformis. *Caul.* herbacea, foliosa. *Fol.* simplex, alterna. *Flor.* terminalis, capitata. *Fulcr.* bracteae.

SAMOLUS. *Cor.* hypocraterif. *Stam.* munita squamulis corollae. *Capsf.* 1-loc. infera.

Rad. fibrosa. *Caul.* herbacea, foliosa. *Fol.* simplex, alterna, sessilis. *Flor.* terminalis, racemosa. *Fulcr.* o.

LONICERA. *Cor.* 1-pet. irregul. *Stigm.* capitat. *Bacca* 2-loc. infera, polysperma.

Rad. lignosa. *Caul.* fruticosa. *Fol.* opposita, simplex. *Flor.* varia. *Fulcr.* o.

VERBASCUM. *Cor.* rotata, subinaequ. *Stam.* lanuginosa. *Capsf.* 2-loc. 2-valv.

Rad. ramosa. *Caul.* herbacea, stricta, foliosa. *Fol.* simplex. *Flor.* longe spicata. *Fulcr.* bracteae.

DATURA. *Cor.* infundib. plicata. *Cal.* tubulosus, angulatus, deciduus. *Capsf.* 4-loc.

Rad. ramosa. *Caul.* herbacea, fastigiata. *Fol.* simplex, angulata, gemina. *Flor.* solitaria, axillaris. *Fulcr.* o.

HYOSCYAMUS. *Cor.* infundib. obtusa. *Stam.* inclinata. *Capsf.* 2-loc. operculata.

Rad. fusiformis. *Caul.* herbacea, foliosa. *Fol.* simplex, alterna, mollis. *Flor.* solitaria, sessilis, axillaris, secunda. *Fulcr.* o.

ATROPA. *Cor.* campanul. *Stam.* distantia, incurvata. *Bacca* 2-locul.

Rad. ramosa. *Caul.* herbacea, foliosa. *Fol.* simplex, binata. *Flor.* solitaria, pedunculata, axillaris. *Fulcr.* o.

PHYSALIS. *Cor.* rotata. *Stam.* conniventia. *Bacca* 2-locul. intra calycem inflatum.

Rad. repens. *Caul.* herbacea, foliosa. *Fol.* simplex, gemina. *Flor.* axillaris. *Fulcr.* o.

SOLANUM. *Cor.* rotata. *Antherae* subcoalitae apice poro gemino dehiscentes. *Bacca* 2-loc.

Rad. ramosa. *Caul.* varia, foliosa. *Fol.* simplex, alterna. *Flor.* oppositifolia, corymbosa. *Fulcr.* o.

RHAMNUS. *Cal.* tubul. squamis petalodeis stamina muni-
entibus. *Cor.* o. *Bacca.*

Rad. lignosa. *Caul.* arborea. *Fol.* simplex, alterna. *Flor.* axillaris, pedunculata. *Fulcr.* o.

EVONYMUS. *Cal.* 4-fid. *Cor.* 4-5-pet. *Caps.* 4-5-gona-
locul. - valv. *Sem.* arillata.

Rad. lignosa. *Caul.* fruticosa. *Fol.* simplex, opposita. *Flor.* axillaris, subumbellata. *Fulcr.* o.

RIBES. *Petala* et *stamina* calyci cavo inserta. *Styl.* 2-fid. *Bacca* infera, polysperma.

Rad. lignosa. *Caul.* fruticosa. *Fol.* simplex, alterna, lobata. *Flor.* axillaris, racemosa l. solitaria. *Fulcr.* bracteae.

HEDERA. *Petala* 5. oblonga. *Bacca* 5-sperma, calyce cincta.

Rad. lignosa. *Caul.* fruticosa, serpens, radicans. *Fol.* simplex, alterna. *Flor.* terminalis, glomerata. *Fulcr.* radiculae.

ILLECEBRUM. *Cal.* 5-phyll. cartilagineus. *Cor.* σ. *Caps.* 5-valv. 1-sperma.

Rad. simplex. *Caul.* herbacea, ramosa, articulata. *Fol.* simplex, opposita. *Flor.* verticillata, axillaris. *Fulcr.* stipulae.

GLAUX. *Cal.* 1-phyll. *Cor.* o. *Capf.* 1-locul. 5-valv. 5-sperma.

Rad. geniculata. *Caul.* herbacea, foliata. *Fol.* simplex, sessilis. *Flor.* solitaria, axillaris sessilis. *Fulcr.* o.

THESIUM. *Cal.* 1-phyll. cui *stamina* inserta. *Sem.* 1. inferum.

Rad. simplex. *Caul.* simplex, herbacea, foliosa. *Fol.* simplex, sessilis. *Flor.* racemosa. *Fulcr.* bracteae.

Polygonum amphibium v. Octandr. Monog.

2. DIGYNIA.

ASCLEPIAS. *Cor.* contorta. *Nect.* 5. ovata, concava, corniculum exferentia. *Folliculi* 2.

Rad. fibrosa. *Caul.* herbacea, foliosa. *Fol.* simplex, opposita. *Flor.* umbellata, terminalis et alaris. *Fulcr.* o.

HERNIARIA. *Cal.* 5-part. *Cor.* o. *Stam.* 5. sterilia. *Capf.* 1-sperm.

Rad. subramosa. *Caul.* herbacea, articulata, procumbens. *Fol.* simplex, opposita, sessilis. *Flor.* axillaris, sessilis, glomerata. *Fulcr.* stipulae.

CHENOPODIUM. *Cal.* 5-phyll. 5-gon. *Cor.* o. *Sem.* 1. lenticulare.

Rad. ramosa. *Caul.* herbacea, foliosa. *Fol.* simplex, alterna. *Flor.* racemoso-glomerata, varia. *Fulcr.* o.

SALSOLA. *Cal.* 5-phyll. *Cor.* o. *Capf.* 1-sperm. *Sem.* 1. cochleatum.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex, sessilis, succulenta. *Flor.* solitaria, sessilis. axillaris. *Fulcr.* o.

ULMUS.

ULMUS. *Cal.* 5-fid. *Cor.* o. *Bacca* exsucca, compressa, membranacea.

Rad. lignosa. *Caul.* arborea. *Fol.* alterna, simplex. *Flor.* axillaris. *Fulcr.* stipulae deciduae.

SWERTIA. *Cor.* rotata. *Nect.* pori 5 ad basin laciniarum corollae. *Capsf.* 1-loc. 2-valv.

Rad. fibrosa. *Caul.* herbacea, foliosa. *Fol.* simplex, opposita. *Flor.* axillaris et terminalis, pedunculata. *Fulcr.* o.

GENTIANA. *Cor.* 1-pet. tubul. 5-4-fid. *Capsf.* 1-locul. 2-valv.

Rad. subsimplex. *Caul.* herbacea, articulata, subulata. *Fol.* opposita, simplex, sessilis. *Flor.* varia. *Fulcr.* o. l. bracteae.

ERYNGIUM. *Flores* capitati. *Recept.* paleaceum.

Rad. ramosa, subhorizontalis. *Caul.* herbacea. *Fol.* simplex, coriacea, caesia, spinosa. *Flor.* capitata. *Fulcr.* involucria.

HYDROCOTYLE. *Umb.* simplex. *Invol.* 4-phyll. *Pet.* integra. *Sem.* suborbiculato-compressa.

Rad. horizontalis. *Caul.* scaposa. *Fol.* simplex, radicalis, peltata. *Flor.* umbellata. *Fulcr.* involucrium.

SANICULA. *Umb.* confertae, subcapitatae. *Fructus* scaber. *Flor.* disci abortientes.

Rad. fusiformis. *Caul.* herbacea, subnuda. *Fol.* simplex, lobata radicalis; caulina sessilis. *Flor.* terminalis, umbellato-glomerata. *Fulcr.* o.

BUPLEURUM. *Invol.* umbellulae majora 5-phyll. *Pet.* involuta. *Fruct.* subrotund. compress. striat.

Rad. subramosa. *Caul.* herbacea. *Fol.* simplex, varia. *Flor.* umbellata, varia. *Fulcr.* involucria.

TORDYLIUM. *Invol. univ. et part. longa, indivisa. Cor. radiatae, omnes hermaphrod. Fruct. sub-orbic. margine crenat.*

Rad. fusiformis. Caul. herbacea. Fol. alterna, pinnato-incisa. Flor. umbellata. Fulcr. involucra.

DAUCUS. *Invol. univ. et part. pinnat. Cor. subradiatae, omnes hermaphr. Fruct. pilis hispidus.*

Rad. fusiformis. Caul. herbacea. Fol. alterna, multifido-pinnata. Flor. umbellata. Fulcr. involucra.

BUNIUM. *Invol. univ. et part. fetaceum. Umb. conferta. Cor. uniform. Fruct. ovat.*

Rad. bulbosa. Caul. herbacea. Fol. alterna, multifido-pinnata. Flor. terminalis, umbellata. Fulcr. involucra.

CONIUM. *Invol. univ. et part. dimidiata, subtriphylla. Fruct. subglob. 5-striat. utrinque crenatus.*

Rad. fusiformis. Caul. herbacea, maculosa. Fol. alterna, pinnatifido-composita. Flor. umbellata. Fulcr. involucra.

PEUCEDANUM. *Invol. univ. et part. simplex, breviss. Flor. flosc. abort. Fruct. ovat. striat. ala cinctus.*

Rad. ramosa. Caul. herbacea. Fol. alterna, laciniata, sublinearis. Flor. umbellata. Fulcr. involucra.

SELINUM. *Invol. univ. et part. reflexum. Fruct. ovali-oblongus, compresso-planus in medio striatus. Cor. aequal.*

Rad. fusiformis. Caul. herbacea. Fol. alterna, multifido-pinnata. Flor. umbellata. Fulcr. involucra.

ATHAMANTA. *Invol. univ. et part. Flor. omnes fertiles. Pet. inflexa emarginata. Fruſt. ovato-oblong. ſtriatus.*

Rad. fuſiformis. Caul. herbacea. Fol. alterna, multifido-pinnata. Flor. umbellata. Fulcr. involucra.

LASERPITIUM. *Invol. univ. et part. Flor. abortientes. Pet. inflexa, emarginata. Fruſt. ovato-oblong. angulis 8 membranaceis.*

Rad. fuſiformis. Caul. herbacea. Fol. alterna, bipinnata. Flor. umbellata. Fulcr. involucra.

HERACLEUM. *Invol. univ. et part. decidua. Flor. radii abort. Fruſt. ellipticus, emargin. comprefſus, ſtriatus, marginatus.*

Rad. fuſiformis. Caul. herbacea, ſtrigofa. Fol. alterna, pinnata, hiſpida. Flor. umbellata. Fulcr. involucra.

LIGUSTICUM. *Invol. univ. et part. membranac. Flor. aequal. fertil. Pet. involuta. Fruſt. oblongus utrinque 5-fulcatus.*

Rad. ramoſa. Caul. herbacea. Fol. alterna, bipinnata. Flor. umbellata. Fulcr. involucra.

ANGELICA. *Invol. univ. et part. Umbellulae globosae. Cor. aequal. Fruſt. ſubrot. angulatus, ſtylis reflexis.*

Rad. ramoſa. Caul. herbacea. Fol. alterna, bipinnata. Flor. umbellata. Fulcr. involucra.

SIUM. *Invol. univ. et part. polyphylla. Pet. cordata. Fruſt. ſubovat. ſtriat.*

Rad. tuberoſa, fibroſa, repens. Caul. herbacea. Fol. alterna, pinnata, diſſimilis. Flor. umbellata. Fulcr. involucra.

SISON.

SISON. *Invol. univ. et part. sub 4-phylla. Umbellae depauperatae. Fruct. ovat. striat.*

Rad. fibrosa. Fol. immerfa multifido-capillacea, emerfa pinnata lobata. Caul. demerfa, repens. Flor. umbellata, flosculis in umbella sessilibus. Fulcr. involucra.

OENANTHE. *Invol. univ. et part. Flor. disci sessiles steriles. Fruct. calyce et pistillo coronat. sessilis.*

Rad. varia. Caul. herbacea. Fol. pinnata. Flor. umbellata. Fulcr. involucra.

PHELLANDRIUM. *Invol. part. Flor. disci minores fertiles. Fruct. ovatus, laevis, coronatus calyce et pistillo.*

Rad. praemorsa fibris verticillatis. Caul. herbacea, divaricata. Fol. alterna, bipinnata. Flor. umbellata. Fulcr. involucella.

SESELI. *Invol. part. Flores flosculosi fertiles. Umb. glob. rigidulae. Fruct. ovat. striat.*

Rad. fusiformis. Caul. herbacea, stricta. Fol. alterna, bipinnata, linearis. Flor. umbellata. Fulcr. involucella.

CICUTA. *Invol. part. setacea. Pet. planiuscula. Fruct. subovatus, sulcatus.*

Rad. ovata basi fibrosa. Caul. herbacea. Fol. tripinnata, serrata. Flor. umbellata. Fulcr. involucella.

AETHUSA. *Invol. part. dimidiata, 3-phyll. pendula. Fruct. striatus.*

Rad. fusiformis. Caul. herbacea. Fol. alterna, bipinnata, laciniata. Flor. umbellata. Fulcr. involucella.

SCANDIX. *Invol. part. Cor. radiata. Fruct. subulatus. Flosc. disci saepe masculi.*

Rad. fusiformis. Caul. herbacea. Fol. alterna, pinnato-multifida. Flor. umbellata. Fulcr. involucella.

V. PENTANDRIA.

I. MONOGYNIA.

MYOSOTIS. *Cor.* hypocraterif. 5-fid. emarginata: fauce clausa fornicibus. *Sem.* 4.

Rad. fibrosa. *Caul.* herbacea, teres. *Fol.* sessilis, alterna, hirta, simplex. *Flor.* terminalis, racemosa. *Fulcr.* o.

LITHOSPERMUM. *Cor.* infundib. fauce perforata, nuda. *Cal.* 5-part. *Sem.* 4.

Rad. ramosa. *Caul.* herbacea, teres. *Fol.* alterna, simplex, sessilis. *Flor.* solitaria. *Fulcr.* o.

ANCHUSA. *Cor.* infundib. fauce clausa fornicibus. *Sem.* 4. basi insculpta.

Rad. ramosa. *Caul.* herbacea, teres, ramosa. *Fol.* alterna, simplex, sessilis. *Flor.* terminalis, spicata. *Fulcr.* o.

CYNOGLOSSUM. *Cor.* infundib. fauce clausa fornicibus. *Sem.* 4. depressa, interiore tantum latere stylo affixa.

Rad. ramosa, crassa. *Caul.* herbacea, ramosa. *Fol.* alterna, simplex, sessilis, mollis. *Flor.* terminalis, ramosa. *Fulcr.* o.

PULMONARIA. *Cor.* infundib. fauce pervia. *Cal.* prismatico-pentagonus. *Sem.* 4.

Rad. ramosa. *Caul.* herbacea, teres, foliosa. *Fol.* alterna, simplex, punctata. *Flor.* paniculata, terminalis. *Fulcr.* o.

SYMPHYTUM. *Cor.* limbus tubulato-ventricosus: fauce clausa radiis lanceolatis. *Sem.* 4.

Rad. fusiformis. *Caul.* herbacea, angulata. *Fol.* simplex, alterna, decurrens. *Flor.* subracemosa. *Fulcr.* o.

ASPERUGO. *Cor.* infundib. *Cal.* fructus compressus: lamellis plano - parallelis sinuatis. *Sem.* 1.

Rad. ramosa. *Caul.* herbacea, ramosa, debilis.

Fol. simplex, sessilis, sparsa. *Flor.* axillaris, congesta, subsessilis, sparsa. *Fulcr.* 0.

LYCOPSI. *Cor.* infundib. tubo curvo. *Sem.* 4.

Rad. ramosa. *Caul.* herbacea, ramosa, fetosa.

Fol. simplex, alterna, sessilis. *Flor.* solitaria, pedunculata. *Fulcr.* 0.

ECHIUM. *Cor.* campanulata, irregul. fauce nuda. *Sem.* 4.

Rad. ramosa. *Caul.* herbacea, teres, tuberculata. *Fol.* simplex, sessilis, alterna. *Flor.* spicata, secunda. *Fulcr.* 0.

DIAPENSIA. *Cal.* 8-phyll. 3. exterioribus. *Cor.* hypocraterif. *Stam.* tubo corollae imposita. *Capsf.* 3-locul. 3-valv.

Rad. fibrosa. *Caul.* ramosa, foliosa. *Fol.* simplex, sessilis, imbricata. *Flor.* solitaria, terminalis. *Fulcr.* 0.

ANDROSACE. *Umbella* involucrata. *Cor.* tubus ovatus, ore glanduloso. *Capsf.* 1-locul. glob. apice 5-fido dehiscens.

Rad. simplex. *Caul.* scaposa. *Fol.* simplex, radicalis. *Flor.* terminalis, umbellata. *Fulcr.* involucreum.

PRIMULA. *Umb.* invol. *Cor.* tubus cylindricus ore patulo. *Capsf.* teres 1-loc. apice 10-dent. dehiscens, polysperma.

Rad. ramosa. *Caul.* scaposa. *Fol.* simplex, radicalis. *Flor.* terminalis, umbellata. *Fulcr.* involucreum.

MENYANTHES. *Cor.* hirsuta. *Stigma* 2-fid. *Capsf.* 1-loc.

Rad. horizontalis, articulata. *Caul.* herbacea, demersa. *Fol.* ternata l. simplex. *Flor.* scaposa, axillaris. *Fulcr.* bracteae.

HOTTONIA. *Cor.* hypocraterif. *Stam.* tubo corollae imposita. *Capsf.* 1-loc.

Rad. fibrosa. *Caul.* demersa, repens, radicans, foliosa. *Fol.* demersa, verticillata, multifida.

Flor. spicato-verticillata. *Fulcr.* bracteae.

LYSIMACHIA. *Cor.* rotata. *Capsf.* globosa, mucronata, 10-valv.

Rad. repens. *Caul.* herbacea, ramosa. *Fol.* simplex, opposita l. verticillata. *Flor.* varia. *Fulcr.* o.

ANAGALLIS. *Cor.* rotata. *Capsf.* circumscissa.

Rad. fibrosa. *Caul.* herbacea, debilis, tetragona.

Fol. simplex, opposita, sessilis. *Flor.* axillaris, solitaria, pedunculata. *Fulcr.* o.

AZALEA. *Cor.* campanulata. *Stam.* receptaculo inserta. *Capsf.* 5-locul.

Rad. lignosa, fibrosa. *Caul.* fruticulosa, ramosa, foliosa. *Fol.* simplex, conferta, semper-virens. *Flor.* terminalis, subterna, pedunculata. *Fulcr.* o.

CONVOLVULUS. *Cor.* campanulata, plicata. *Stigm.* 2. *Capsf.* 2-loc. 4-sperma.

Rad. repens. *Caul.* herbacea, foliosa, volubilis. *Fol.* simplex, alterna. *Flor.* axillaris, pedunculata. *Fulcr.* bracteae.

POLEMONIUM. *Cor.* 5-part. fundo clauso valvis staminiferis. *Stigm.* 3-fid. *Capsf.* 3-loc. supera.

Rad. fibrosa. *Caul.* herbacea, striata, foliosa. *Fol.* alterna, sessilis, pinnata. *Flor.* racemosa. *Fulcr.* o.

CAMPANULA. *Cor.* campanulata fundo clauso valvis staminiferis. *Stigm.* 3-fid. *Capsf.* infera, poris lateralibus dehiscens.

Rad. ramosa, tuberosa, fibrosa. *Caul.* herbacea, foliosa. *Fol.* simplex, varia. *Flor.* varia. *Fulcr.* bracteae.

PHYTEUMA. *Cor.* rotata, 5-part. laciniis linearibus.
Stam. reflexa. *Stigma* 2-fid. *Capsf.* 2-locul.
infera.

Rad. fusiformis. *Caul.* herbacea, foliosa. *Fol.*
simplex, alterna. *Flor.* terminalis, capitata.
Fulcr. bracteae.

SAMOLUS. *Cor.* hypocraterif. *Stam.* munita squamulis
corollae. *Capsf.* 1-loc. infera.

Rad. fibrosa. *Caul.* herbacea, foliosa. *Fol.* sim-
plex, alterna, sessilis. *Flor.* terminalis, race-
mosa. *Fulcr.* o.

LONICERA. *Cor.* 1-pet. irregul. *Stigm.* capitat. *Bacca*
2-loc. infera, polysperma.

Rad. lignosa. *Caul.* fruticosa. *Fol.* opposita,
simplex. *Flor.* varia. *Fulcr.* o.

VERBASCUM. *Cor.* rotata, subinaequ. *Stam.* lanugi-
nosa. *Capsf.* 2-loc. 2-valv.

Rad. ramosa. *Caul.* herbacea, stricta, foliosa.
Fol. simplex. *Flor.* longe spicata. *Fulcr.*
bracteae.

DATURA. *Cor.* infundib. plicata. *Cal.* tubulosus, an-
gulatus, deciduus. *Capsf.* 4-loc.

Rad. ramosa. *Caul.* herbacea, fastigiata. *Fol.*
simplex, angulata, gemina. *Flor.* solitaria,
axillaris. *Fulcr.* o.

HYOSCYAMUS. *Cor.* infundib. obtusa. *Stam.* inclinata.
Capsf. 2-loc. operculata.

Rad. fusiformis. *Caul.* herbacea, foliosa. *Fol.*
simplex, alterna, mollis. *Flor.* solitaria, ses-
silis, axillaris, secunda. *Fulcr.* o.

ATROPA. *Cor.* campanul. *Stam.* distantia, incurvata.
Bacca 2-locul.

Rad. ramosa. *Caul.* herbacea, foliosa. *Fol.* sim-
plex, binata. *Flor.* solitaria, pedunculata,
axillaris. *Fulcr.* o.

PHYSALIS. *Cor.* rotata. *Stam.* conniventia. *Bacca* 2-locul. intra calycem inflatum.

Rad. repens. *Caul.* herbacea, foliosa. *Fol.* simplex, gemina. *Flor.* axillaris. *Fulcr.* o.

SOLANUM. *Cor.* rotata. *Antherae* subcoalitae apice poro gemino dehiscentes. *Bacca* 2-loc.

Rad. ramosa. *Caul.* varia, foliosa. *Fol.* simplex, alterna. *Flor.* oppositifolia, corymbosa. *Fulcr.* o.

RHAMNUS. *Cal.* tubul. squamis petalodeis stamina muni-
entibus. *Cor.* o. *Bacca.*

Rad. lignosa. *Caul.* arborea. *Fol.* simplex, alterna. *Flor.* axillaris, pedunculata. *Fulcr.* o.

EVONYMUS. *Cal.* 4-fid. *Cor.* 4-5-pet. *Capsf.* 4-5-gona-
locul. - valv. *Sem.* arillata.

Rad. lignosa. *Caul.* fruticosa. *Fol.* simplex, opposita. *Flor.* axillaris, subumbellata. *Fulcr.* o.

RIBES. *Petala* et *stamina* calyci cavo inserta. *Styl.* 2-fid. *Bacca* infera, polysperma.

Rad. lignosa. *Caul.* fruticosa. *Fol.* simplex, alterna, lobata. *Flor.* axillaris, racemosa l. solitaria. *Fulcr.* bracteae.

HEDERA. *Petala* 5. oblonga. *Bacca* 5-sperma, calyce cincta.

Rad. lignosa. *Caul.* fruticosa, serpens, radicans. *Fol.* simplex, alterna. *Flor.* terminalis, glomerata. *Fulcr.* radiculae.

ILLECEBRUM. *Cal.* 5-phyll. cartilagineus. *Cor.* o. *Capsf.* 5-valv. 1-sperma.

Rad. simplex. *Caul.* herbacea, ramosa, articulata. *Fol.* simplex, opposita. *Flor.* verticillata, axillaris. *Fulcr.* stipulae.

GLAUX. *Cal.* 1-phyll. *Cor.* o. *Caps.* 1-locul. 5-valv. 5-sperma.

Rad. geniculata. *Caul.* herbacea, foliata. *Fol.* simplex, sessilis. *Flor.* solitaria, axillaris sessilis. *Fulcr.* o.

THÆSIUM. *Cal.* 1-phyll. cui *stamina* inserta. *Sem.* 1. inferum.

Rad. simplex. *Caul.* simplex, herbacea, foliosa. *Fol.* simplex, sessilis. *Flor.* racemosa. *Fulcr.* bracteae.

Polygonum amphibium v. Octandr. Monog.

2. DIGYNIA.

ASCLEPIAS. *Cor.* contorta. *Nect.* 5. ovata, concava, corniculum exferentia. *Folliculi* 2.

Rad. fibrosa. *Caul.* herbacea, foliosa. *Fol.* simplex, opposita. *Flor.* umbellata, terminalis et alaris. *Fulcr.* o.

HERNIARIA. *Cal.* 5-part. *Cor.* o. *Stam.* 5. sterilia. *Caps.* 1-sperm.

Rad. subramosa. *Caul.* herbacea, articulata, procumbens. *Fol.* simplex, opposita, sessilis. *Flor.* axillaris, sessilis, glomerata. *Fulcr.* stipulae.

CHENOPODIUM. *Cal.* 5-phyll. 5-gon. *Cor.* o. *Sem.* 1. lenticulare.

Rad. ramosa. *Caul.* herbacea, foliosa. *Fol.* simplex, alterna. *Flor.* racemoso-glomerata, varia. *Fulcr.* o.

SALSOLA. *Cal.* 5-phyll. *Cor.* o. *Caps.* 1-sperm. *Sem.* 1. cochleatum.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex, sessilis, succulenta. *Flor.* solitaria, sessilis. axillaris. *Fulcr.* o.

ULMUS.

ULMUS. *Cal.* 5-fid. *Cor.* o. *Bacca* exsucca, compressa, membranacea.

Rad. lignosa. *Caul.* arborea. *Fol.* alterna, simplex. *Flor.* axillaris. *Fulcr.* stipulae deciduae.

SWERTIA. *Cor.* rotata. *Nest.* pori 5 ad basin laciniarum corollae. *Capsf.* 1 loc. 2-valv.

Rad. fibrosa. *Caul.* herbacea, foliosa. *Fol.* simplex, opposita. *Flor.* axillaris et terminalis, pedunculata. *Fulcr.* o.

GENTIANA. *Cor.* 1-pet. tubul. 5-4-fid. *Capsf.* 1-locul. 2-valv.

Rad. subsimplex. *Caul.* herbacea, articulata, subalata. *Fol.* opposita, simplex, sessilis. *Flor.* varia. *Fulcr.* o. l. bracteae.

ERYNGIUM. *Flores* capitati. *Recept.* paleaceum.

Rad. ramosa, subhorizontalis. *Caul.* herbacea. *Fol.* simplex, coriacea, caesia, spinosa. *Flor.* capitata. *Fulcr.* involucria.

HYDROCOTYLE. *Umb.* simplex. *Invol.* 4-phyll. *Pet.* integra. *Sem.* suborbiculato-compressa.

Rad. horizontalis. *Caul.* scaposa. *Fol.* simplex, radicalis, peltata. *Flor.* umbellata. *Fulcr.* involucrium.

SANICULA. *Umb.* confertae, subcapitatae. *Fruſtus* scaber. *Flor.* disci abortientes.

Rad. fusiformis. *Caul.* herbacea, subnuda. *Fol.* simplex, lobata radicalis; caulina sessilis. *Flor.* terminalis, umbellato-glomerata. *Fulcr.* o.

BUPLEURUM. *Invol.* umbellulae majora 5-phyll. *Pet.* involuta. *Fruſt* subrotund. compress. striat.

Rad. subramosa. *Caul.* herbacea. *Fol.* simplex, varia. *Flor.* umbellata, varia. *Fulcr.* involucria.

TORDYLIUM. *Invol. univ. et part. longa, indivisa. Cor. radiatae, omnes hermaphrod. Fruct. sub-orbic. margine crenat.*

Rad. fusiformis. Caul. herbacea. Fol. alterna, pinnato-incisa. Flor. umbellata. Fulcr. involucra.

DAUCUS. *Invol. univ. et part. pinnat. Cor. subradiatae, omnes hermaphr. Fruct. pilis hispidus.*

Rad. fusiformis. Caul. herbacea. Fol. alterna, multifido-pinnata. Flor. umbellata. Fulcr. involucra.

BUNIAM. *Invol. univ. et part. setaceum. Umb. conferta. Cor. uniform. Fruct. ovat.*

Rad. bulbosa. Caul. herbacea. Fol. alterna, multifido-pinnata. Flor. terminalis, umbellata. Fulcr. involucra.

CONIUM. *Invol. univ. et part. dimidiata, subtriphylla. Fruct. subglob. 5-striat. utrinque crenatus.*

Rad. fusiformis. Caul. herbacea, maculosa. Fol. alterna, pinnatifido-composita. Flor. umbellata. Fulcr. involucra.

PEUCEDANUM. *Invol. univ. et part. simplex, breviss. Flor. flosc. abort. Fruct. ovat. striat. ala cinctus.*

Rad. ramosa. Caul. herbacea. Fol. alterna, laciniata, sublinearis. Flor. umbellata. Fulcr. involucra.

SELINUM. *Invol. univ. et part. reflexum. Fruct. ovali-oblongus; compresso-planus in medio striatus. Cor. aequal.*

Rad. fusiformis. Caul. herbacea. Fol. alterna, multifido-pinnata. Flor. umbellata. Fulcr. involucra.

ATHAMANTA. *Invol. univ. et part. Flor. omnes fertiles. Pet. inflexa emarginata. Fruct. ovato-oblong. striatus.*

Rad. fusiformis. Caul. herbacea. Fol. alterna, multifido-pinnata. Flor. umbellata. Fulcr. involucri.

LASERPITIUM. *Invol. univ. et part. Flor. abortientes. Pet. inflexa, emarginata. Fruct. ovato-oblong. angulis 8 membranaceis.*

Rad. fusiformis. Caul. herbacea. Fol. alterna, bipinnata. Flor. umbellata. Fulcr. involucri.

HERACLEUM. *Invol. univ. et part. decidua. Flor. radii abort. Fruct. ellipticus, emargin. compressus, striatus, marginatus.*

Rad. fusiformis. Caul. herbacea, strigosa. Fol. alterna, pinnata, hispida. Flor. umbellata. Fulcr. involucri.

LIGUSTICUM. *Invol. univ. et part. membranac. Flor. aequal. fertil. Pet. involuta. Fruct. oblongus utrinque 5-sulcatus.*

Rad. ramosa. Caul. herbacea. Fol. alterna, bipinnata. Flor. umbellata. Fulcr. involucri.

ANGELICA. *Invol. univ. et part. Umbellulae globosae. Cor. aequal. Fruct. subrot. angulatus, stylis reflexis.*

Rad. ramosa. Caul. herbacea. Fol. alterna, bipinnata. Flor. umbellata. Fulcr. involucri.

SIUM. *Invol. univ. et part. polyphylla. Pet. cordata. Fruct. subovat. striat.*

Rad. tuberosa, fibrosa, repens. Caul. herbacea. Fol. alterna, pinnata, dissimilis. Flor. umbellata. Fulcr. involucri.

SISON. *Invol. univ. et part. sub 4-phylla. Umbellae depauperatae. Fruct. ovat. striat.*
Rad. fibrosa. Fol. immerfa multifido-capillacea, emerfa pinnata lobata. Caul. demerfa, repens. Flor. umbellata, flosculis in umbella sessilibus. Fulcr. involucri.

OENANTHE. *Invol. univ. et part. Flor. disci sessiles steriles. Fruct. calyce et pistillo coronat. sessilis.*
Rad. varia. Caul. herbacea. Fol. pinnata. Flor. umbellata. Fulcr. involucri.

PHELLANDRIUM. *Invol. part. Flor. disci minores fertiles. Fruct. ovatus, laevis, coronatus calyce et pistillo.*

Rad. praemorsa fibris verticillatis. Caul. herbacea, divaricata. Fol. alterna, bipinnata. Flor. umbellata. Fulcr. involucri.

SESSELY. *Invol. part. Flores flosculosi fertiles. Umb. glob. rigidulae. Fruct. ovat. striat.*

Rad. fusiformis. Caul. herbacea, stricta. Fol. alterna, bipinnata, linearis. Flor. umbellata. Fulcr. involucri.

CICUTA. *Invol. part. setacea. Pet. planiuscula. Fruct. subovatus, sulcatus.*

Rad. ovata basi fibrosa. Caul. herbacea. Fol. tripinnata, serrata. Flor. umbellata. Fulcr. involucri.

AETHUSA. *Invol. part. dimidiata, 3-phyll. pendula. Fruct. striatus.*

Rad. fusiformis. Caul. herbacea. Fol. alterna, bipinnata, laciniata, Flor. umbellata. Fulcr. involucri.

SCANDIX. *Invol. part. Cor. radiata. Fruct. subulatus. Flosc. disci saepe masculi.*

Rad. fusiformis. Caul. herbacea. Fol. alterna, pinnato-multifida. Flor. umbellata. Fulcr. involucri.

CHAEROPHYLLUM. *Invol. part.* reflexum, concavum, sub5-phyll. *Fruſt.* oblongus, laevis.

Rad. fusiformis. *Caul.* herbacea, geniculato-tumida. *Fol.* alterna, multifido-pinnata. *Flor.* umbellata. *Fulcr.* involucella.

IMPERATORIA. *Invol. part.* parciſſimum. *Umb.* expanso-plana. *Fruſt.* subrot. compress. medio gibbus, marginatus.

Rad. tuberosa, annulata. *Caul.* herbacea, simplex. *Fol.* triternata, serrata. *Flor.* umbellata. *Fulcr.* plerumque nulla.

PASTINACA. *Invol.* o. *Pet.* involuta, integra. *Fruſtus* ellipticus, compresso-planus.

Rad. fusiformis. *Caul.* herbacea. *Fol.* alterna, pinnata, serrata. *Flor.* umbellata. *Fulcr.* o.

CARUM. *Invol.* o. l. 1-phyll. *Pet.* carinata, inflexo-emarginata. *Fruſt.* ovato-oblongus, striatus.

Rad. fusiformis. *Caul.* herbacea. *Fol.* alterna, bipinnato-incisa. *Flor.* umbellata. *Fulcr.* subnulla.

PIMPINELLA. *Invol.* o. *Umb.* ante florescentiam nutantes. *Stigm.* subglobosa. *Fruſtus* ovato oblongus.

Rad. fusiformis, longa. *Caul.* herbacea. *Fol.* composita, difformis. *Flor.* umbellata. *Fulcr.* o.

APIUM. *Invol.* o. l. 1-phyll. *Pet.* aequal. *Fruſtus* minutus, ovatus, striatus.

Rad. praemorsa. *Caul.* herbacea, sulcata. *Fol.* pinnata, difformis. *Flor.* umbellata. *Fulcr.* subnulla.

AEGOPodium. *Invol.* o. *Pet.* cordata. *Sem.* gibba striata.

Rad. repens. *Caul.* herbacea. *Fol.* pinnato-ternata. *Flor.* umbellata. *Fulcr.* o.

Cuscuta Epithymum v. Tetrandr. digyn.

PENTANDRIA.

51

3. TRIGYNIA.

VIBURNUM. *Cal.* 5-part. *super.* *Cor.* 5-fid. *Bacca* 1-sperma.

Rad. lignosa. *Caul.* fruticosa. *Fol.* simplex, opposita. *Flor.* corymbosa, radiata. *Fulcr.* o.

SAMBUCUS. *Cal.* 5-part. *Cor.* 5-fid. *Bacca* trisperma.

Rad. lignosa. *Caul.* fruticosa. *Fol.* pinnata, opposita. *Flor.* cymosa. *Fulcr.* stipulae.

TAMARIX. *Cal.* 5-part. *Cor.* 5-pet. *Capsf.* 1-loc. 3-valv. *Sem.* papposa.

Rad. lignosa. *Caul.* fruticosa, virgata. *Fol.* simplex, minuta, imbricata. *Flor.* racemosa. *Fulcr.* stipulae.

CORRIGIOLA. *Cal.* 5-phyll. *Cor.* 5-pet. *Sem.* 1. triquetrum.

Rad. fibrosa. *Caul.* herbacea, prostrata. *Fol.* simplex, alterna, sessilis. *Flor.* glomerata, terminalis. *Fulcr.* o.

ALSINE. *Cal.* 5-phyll. *Cor.* 5-pet. *Stam.* 3-10. *Capsf.* 1-loc. 3-valv.

Rad. fibrosa. *Caul.* herbacea, geniculata. *Fol.* simplex, opposita. *Flor.* axillaris, pedunculata. *Fulcr.* o.

4. TETRAGYNIA.

PARNASSIA. *Cal.* 5-part. *Cor.* 5-pet. *Nect.* 5. cordata, ciliata, ciliis capitatis. *Styl.* o. *Capsf.* 4-valv.

Rad. fibrosa. *Caul.* simplex, monophylla. *Fol.* simplex, cordata. *Flor.* solitaria, terminalis. *Fulcr.* o.

5. PENTAGYNIA.

STATICE. *Cal.* 1-phyll. plicatus, scariosus. *Cor.* 5-pet. *Sem.* 1. superum.

Rad. fibrosa. *Caul.* scaposa. *Fol.* simplex, radicalis. *Flor.* capitata l. paniculato-racemosa. *Fulcr.* bracteae.

LINUM. *Cal.* 5-phyll. *Cor.* 5-pet. *Capf.* 5-valv. 10-loc.
Sem. solitaria.

Rad. simplex. *Caul.* herbacea, foliosa. *Fol.* simplex, sessilis. *Flor.* paniculata. *Fulcr.* o.

DROSERA. *Cal.* 5-fid. *Cor.* 5-pet. *Capf.* 1-loc. apice
5-valvi, polysperma.

Rad. fibrosa. *Caul.* scaposa. *Fol.* simplex, radicalis, glandulis stipitatis. *Flor.* racemosa, secunda. *Fulcr.* o.

CRASSULA. *Cal.* 5-phyll. *Cor.* 5-pet. *Squamae* 5. nectariferae ad basin germinis. *Capf.* 5.

Rad. fibrosa. *Caul.* herbacea, succulenta. *Fol.* simplex, opposita, succulenta. *Flor.* cymosa. *Fulcr.* o.

SIBBALDIA. *Cal.* 10-fid. *Cor.* 5-pet. calyci inserta.
Styli e latere germinis. *Sem.* 5.

Rad. horizontalis. *Caul.* herbacea. *Fol.* ternata.
Flor. subumbellata. *Fulcr.* bracteae.

6. POLYGYNIA.

MYOSURUS. *Cal.* 5-phyll. adnatus. *Nect.* 5. petaliformia. *Recept.* elongatum.

Rad. fibrosa. *Caul.* scaposa. *Fol.* simplex, radicalis, linearis. *Flor.* solitaria, terminalis. *Fulcr.* o.

V. PENTANDRIA.

I. MONOGYNIA.

MYOSOTIS.

223. *M. scorpioides*, feminibus laevibus, foliorum apicibus callofis. Fl. Sv. 157. Fl. N. 285. P.
α. arvensis, hirsuta. Fl. D. 583. pr. pasc.
β. palustris, glabra. riv. l. lac.
224. *M. Lappula*, feminum aculeis glochidibus, foliis lanceolatis pilosis. Fl. Sv. 158. Fl. D. 692. A. ficc. ster.

LITHOSPERMUM.

225. *L. officinale*, feminibus laevibus, corollis calycem vix superantibus, foliis lanceolatis. Fl. Sv. 159. Fl. N. 1034. P. ficc.
226. *L. arvense*, feminibus rugosis, corollis calycem vix superantibus, foliis lineari-lanceolatis. Fl. Sv. 160. Fl. N. 613. Fl. D. 456. A. agr.

ANCHUSA.

227. *A. officinalis*, foliis lanceolatis, spicis imbricatis secundis. Fl. Sv. 161. Fl. N. 1026. Fl. D. 572. B. r. an.

CYNOGLOSSUM.

228. *C. officinale*, staminibus corolla brevioribus, foliis lato-lanceolatis tomentosis sessilibus. Fl. Sv. 162. Fl. N. 1014. B. r. arv.

PULMONARIA.

229. *P. officinalis*, foliis radicalibus ovato-cordatis scabris. Fl. Sv. 163. Fl. N. 612. Fl. D. 482. 483. P. nem.
α. foliis maculosis.
β. foliis non maculosis.

54 PENTANDRIA. MONOGYNIA. Symphytum.

230. *P. angustifolia*, foliis radicalibus lanceolatis. Fl. Sv. 164. Fl. N. 1115. P. pr. apr.
231. *P. maritima*, calycibus abbreviatis, foliis ovatis, caule ramoso procumbente. Fn. Sv. 557. Fl. N. 17. Fl. D. 25. P. l. mar.

SYMPHYTUM.

232. *S. officinale*, foliis ovato-lanceolatis alternis decurrentibus. Fl. Sv. 165. Fl. N. 382. Fl. D. 664. P. pisc. foss.

ASBERUGO.

233. *A. procumbens*, calycibus fructus compressis. Fl. Sv. 166. Fl. N. 508. Fl. D. 552. A. fr. r.

LYCOPSIS.

234. *L. arvensis*, foliis lanceolatis hispidis, calycibus florescentibus erectis. Fl. Sv. 167. Fl. N. 498. Fl. D. 435. A. c.

ECHIUM.

235. *E. vulgare*, caule tuberculato hispido, foliis caulinis lanceolatis hispidis, floribus spicatis lateralibus. Fl. Sv. 168. Fl. N. 1033. Fl. D. 445. B. arv. r.

DIAPENSIA.

236. *D. lapponica*, floribus pedunculatis. Fl. Sv. 169. Fl. L. 88. t. 1. f. 1. Fl. N. 118. Fl. D. 47. A. H. X. t. 6. f. 20. P. alp.

ANDROSACE.

237. *A. septentrionalis*, foliis lanceolatis, dentatis glabris, perianthiis angulatis corolla brevioribus. Fl. Sv. 170. Fl. N. 618. Fl. D. 7. B. apr. a.
β. *acaulis*, umbella radicali acauli. mr.

245. *L. thyrsiflora*, racemis lateralibus pedunculatis.
Fl. Sv. 176. Fl. N. 82. Fl. D. 517. P. pr. ul.
- † 246. *L. punctata*, foliis subquaternis subsessilibus nigro-punctatis, pedunculis verticillatis unifloris.
S. N. 196. Clus. Hist. 2. p. 52 v). P.
247. *L. nemorum*, foliis ovatis acutis floribus solitariis, caule procumbente. S. N. 196. Fl. D. 174. Weig. Fl. 132. P. nem. ul.
248. *L. Nummularia*, foliis subcordatis, caule repente. Fl. Sv. 177. Fl. N. 1031. Fl. D. 493. P. pr. hum. sylv.

ANAGALLIS.

249. *A. arvensis*, foliis indivisis, caule procumbente. Fl. Sv. 178. Fl. D. 88. A. c.

AZALEA.

250. *A. Lapponica*, foliis adpersis punctis excavatis. Fl. Sv. 180. Fl. L. t. 6. f. 1. A. H. X. p. 430. Fl. D. 966. S. alp.
251. *A. procumbens*, ramis diffusis procumbentibus. Fl. Sv. 179. Fl. L. t. 6. f. 2. Fl. D. 9. Fl. N. 235. S. alp.

CONVOLVULUS.

252. *C. arvensis*, foliis sagittatis utrinque acutis, pedunculis subunifloris. Fl. Sv. 181. Fl. N. 721. Fl. D. 459. P. c.
253. *C. sepium*, foliis sagittatis postice truncatis, pedunculis tetragonis unifloris. Fl. Sv. 182. Fl. N. 960. Fl. D. 458. P. sep. nem.

POLE-

- v) *E Selandia* mihi misit *Cel. Florman*. Morisoni icon II. S. 5. t. 10. f. 15. sat bona, at descriptio pag. 491. ex Clusio partim mutuata misere, cum descriptione *Lyfimachiae thyrsiflorae* Clusiana confundit.

POLEMONIUM.

254. *P. coeruleum*, foliis pinnatis, floribus erectis, calycibus tubo corollae longioribus. Fl. Sv. 183. Fl. N. 83. Fl. D. 255. P. pr. m.

CAMPANULA.

255. *C. uniflora*, caule unifloro, calyce corollam aequante. Fl. Sv. 185. Fl. L. t. 9. f. 5. 6. A. H. X. p. 432, t. 6, f. 19. Fl. N. 1080. P. alp.
256. *C. rotundifolia*, foliis radicalibus reniformibus: caulinis linearibus. Fl. Sv. 184. Fl. N. 362. P. pr. v. agr.
- β. *parviflora*. Fl. D. 189. alp.
257. *C. patula*, foliis strictis: radicalibus lanceolato-ovalibus, panicula patula. Fl. Sv. 186. Fl. D. 373. B. f.
- † 258. *C. Rapunculus*, foliis undulatis: radicalibus lanceolato ovalibus, panicula coarctata. S. N. 206. Fl. D. 855. ? B. pr. sylv.
259. *C. persicifolia*, foliis radicalibus obovatis, caulinis lanceolato-linearibus subserratis sessilibus remotis. Fl. Sv. 187. P. nem.
260. *C. latifolia*, foliis ovato-lanceolatis, caule simplicissimo tereti, floribus solitariis pedunculatis, fructibus cernuis. Fl. Sv. 188. Fl. N. 55. Fl. D. 782. P. nem.
- ♀. flore albo etc. Fl. D. 85.
261. *C. rapunculoides*, foliis cordato lanceolatis, caule ramoso, floribus secundis sparsis, calycibus reflexis. Fn. Sv. 557. Weig. Fl. 140. P. pr.
262. *C. Trachelium*, caule angulato, foliis petiolatis ciliatis, pedunculis trifidis. Fl. Sv. 189. P. sylv.
263. *C. glomerata*, caule angulato simplici, floribus sessilibus, capitulo terminali, foliis ovato-lanceolatis subtomentosis. Fl. Sv. 190. it. Scan. 306. P. pr.

58. PENTANDRIA. MONOGYNIA. Phyteuma.

264. *C. Cervicaria*, hispida, floribus sessilibus, capitulo terminali, foliis lanceolato-linearibus undulatis. Fl. Sv. 191. Fl. D. 787. P. sylv.

† 265. *C. hederacea*, foliis cordatis quinquelobis petiolatis glabris, caule laxo. S. N. 210. Fl. D. 330. P. pr. hum.

PHYTEUMA.

† 266. *P. spicata*, spica oblonga, capsulis bilocularibus, foliis radicalibus cordatis. S. N. 218. Fl. D. 362. Weig. Fl. 143. P. sylv. w).

SAMOLUS.

267. *S. Valerandi*, Fl. Sv. 192. Fl. D. 198. A. l. m.

LONICERA.

268. *L. periclymenum*, capitulis ovatis imbricatis terminalibus, foliis omnibus distinctis. Fl. Sv. 193. Fl. N. 307. Fl. D. 908. F. sylv.

269. *L. Xylosteum*, pedunculis bifloris, baccis distinctis, foliis integerrimis pubescentibus. Fl. Sv. 194. Fl. N. 1029. Fl. D. 808. F. nem.

270. *L. coerulea*, pedunculis bifloris, baccis coadunatis globosis, stylis indivisis. S. N. 216 x). F.

VERBASCUM.

271. *V. Thapsus*, foliis decurrentibus utrinque tomentosis, caule simplici. Fl. Sv. 195. Fl. N. 349. Fl. D. 631. B. ster. a.

272. *V. Lychnitis*, foliis cuneiformi-oblongis. Fl. Sv. 196. Fl. D. 586. A. r.

273. *V. nigrum*, foliis cordato-oblongis petiolatis. Fl. Sv. 197. Fl. N. 633. P. r.

DATU-

w) etiam in Norvegia. Ström.

x) in Nericia.

DATURA.

274. *D. Stramonium*, pericarpis spinosis erectis ovatis glabris. Fl. Sv. 198. Fl. D. 436. A. r.

HYOSCYAMUS.

275. *H. niger*, foliis amplexicaulibus sinuatis, floribus sessilibus. Fl. Sv. 199. Fl. N. 350. A. B. r.

ATROPA.

- † 276. *A. Belladonna*, caule herbaceo, foliis ovatis integris. S. N. 221. Fl. D. 758. P. sylv.

PHYSALIS.

- † 277. *P. Alkekengi*, foliis geminis integris acutis, caule herbaceo inferne subramoso. S. N. 222. Fl. Fr. 191. P. nem.

SOLANUM.

278. *S. nigrum*, caule inermi herbaceo, foliis ovatis dentato-angulatis, racemis distichis nutantibus. Fl. Sv. 200. Fl. D. 460. A. r. c.
279. *S. Dulcamara*, caule inermi frutescente flexuoso, foliis superioribus hastatis, racemis cymosis. Fl. Sv. 201. Fl. N. 23. Fl. D. 607. S. f. hum.

RHAMNUS.

280. *R. catharticus*, spinis terminalibus, floribus quadrifidis dioicis polygamicisque, foliis ovatis. Fl. Sv. 202. Fl. N. 239. Fl. D. 850. Arb. sylv. nem.
281. *R. Frangula*, inermis, floribus monogynis hermaphroditis, foliis integerrimis. Fl. Sv. 203. Fl. N. 18. Fl. D. 278. Arb. hum.

EVONYMUS.

282. *E. europaeus*, floribus plerisque quadrifidis, foliis sessilibus. Fl. Sv. 204. F. nem.

RIBES.

RIBES.

283. *R. rubrum*, inerme, racemis glabris pendulis, floribus planiusculis. Fl. Sv. 205. Fl. N. 94. Fl. D. 967. F. fylv.
284. *R. alpinum*, inerme, racemis erectis, bracteis flore longioribus. Fl. Sv. 206. Jacqu. Fl. Austr. I. t. 47. Obs. Bot. Fasc. II. 17. a. F. fylv.
285. *R. petraeum*, inerme, racemis florentibus erectis fructescentibus pendulis. S. N. 243. *R. alpinum* β. Obs. Bot. Fasc. II. p. 11. Fl. N. 395. t. 2. f. 1. 2. Fl. D. 968. F. fylv.
286. *R. nigrum*, inerme, racemis pilosis, floribus oblongis. Fl. Sv. 207. Fl. N. 351. Fl. D. 556. F. fylv.
287. *R. Grossularia*, ramis aculeatis petiolorum ciliis pilosis, baccis hirsutis. S. N. 243. F. fylv.
288. *R. Uva crispa*, ramis aculeatis, baccis glabris, pedicellis bractea monophylla. Fl. Sv. 208. Fl. D. 546 y). F. fylv.

HEDERA.

289. *H. Helix*, foliis ovatis lobatisque. Fl. Sv. 209. F. fylv.

ILLECEBRUM.

- †290. *I. verticillatum*, floribus verticillatis nudis, caulibus procumbentibus. S. N. 248. Fl. D. 335. A. a. inund.

GLAUX.

291. *G. maritima*, Fl. Sv. 210. Fl. N. 62. Fl. D. 548. P. l. m. a.

THE-

y) vix et species ab antecedente diversa considerari debet.

THESIUM.

292. *T. alpinum*, racemo foliato, foliis linearibus, floribus quadrifidis tetrandris. S. N. 249. Jacqu. Vind. p. 213. Fl. Austr. V. t. 416. P. apr. mont.
293. *T. linophyllum*, panicula foliacea, foliis linearibus. Fl. Sv. 211 z). B. pr. apr.

2. DIGYNIA.

ASCLEPIAS.

294. *A. Vincetoxicum*, foliis ovatis basi barbatis, caule erecto, umbellis proliferis. Fl. Sv. 212. Fl. D. 849. P. mont.

HERNIARIA.

- 294 *. *H. glabra*, herbacea glabra. Fl. Sv. 213. Fl. N. 926. Fl. D. 529. A. ficc.

CHENOPODIUM.

295. *C. Bonus Henricus*, foliis triangulari-sagittatis integerrimis, spicis compositis aphyllis axillaribus. Fl. Sv. 214. Fl. N. 611. Fl. D. 579. P. r.
296. *C. urbicum*, foliis triangularibus subdentatis, racemis confertis strictissimis, cauli approximatis longissimis. Fl. Sv. 215. Fl. N. 924. A. r.
297. *C. rubrum*, foliis cordato-triangularibus obtusiusculis dentatis, racemis erectis compositis subfoliosis caule brevioribus. Fl. Sv. 217. Fl. N. 540. A. r.
298. *C. murale*, foliis ovatis nitidis dentatis acutis, racemis ramosis nudis. Fl. Sv. 216. A. r.

299. *C.*

z) obscura adhuc satis harum differentia specifica, quare utriusque descriptio graphica desideratur.

299. *C. viride*, foliis rhomboideis dentato-sinuatis, racemis ramosis subfoliatis. Fl. Sv. 219. Fl. N. 298. A. r. c.

β. *C. album*, foliis rhomboideo-triangularibus erosius postice integris: summis oblongis, racemis erectis. Fl. Sv. 218. Fl. N. 297. Vaill. Par. t. 7. f. 1. a).

300. *C. hybridum*, foliis cordatis angulato-acuminatis, racemis ramosis nudis. Fl. Sv. 220. Vaill. Par. t. 7. f. 2. A. c.

301. *C. ambrosioides*, foliis lanceolatis dentatis, racemis foliatis simplicibus. S. N. 262. Obs. Bot. Fasc. I. p. 13. A. r.

302. *C. glaucum*, foliis ovato-oblongis repandis, racemis nudis simplicibus glomeratis. Fl. Sv. 221. A. r.

303. *C. Vulvaria*, foliis integerrimis rhomboideo-ovatis, floribus conglomeratis axillaribus. Fl. Sv. 222. Fl. N. 925. A. r.

304. *C. polyspermum*, foliis integerrimis ovatis, caule decumbente, cymis dichotomis aphyllis axillaribus. Fl. Sv. 223. A. c. h.

305. *C. maritimum*, foliis subulatis semicylindricis. Fl. Sv. 244. Fl. N. 296. Fl. D. 489. A. l. m.

SALSOLA.

306. *S. Kali*, herbacea, foliis subulatis spinosis, calycibus marginatis axillaribus. Fl. Sv. 225. A. l. m.

α. *decumbens* piloso-scabra, digitalis, semipedalis. Fl. D. 818.

β. *erecta* hispida, bipedalis et ultra.

γ. *erecta* glaberrima, bipedalis et ultra β).

† 307. *S.*

a) conf. Cl. Afzelii Diff. cit. p. 28.

b) Si *Salsola Tragus* Sp. Pl. 322. distincta sit species, β et γ ad illam referendae; meras autem varietates adhuc credo, caulem licet habent striatum, in α. quasi farinaceo-scabrum.

- † 307. *S. hirsuta*, herbacea diffusa, foliis teretibus obtusis tomentosis. S. N. 263. Fl. D. 187. A. l. m.

ULMUS.

308. *U. campestris*, foliis duplicato-ferratis basi inaequalibus. Fl. Sv. 226. Fl. N. 295. Fl. D. 632. Arb.

SWERTIA.

- † 309. *S. fulcata*, corollis quinquepartitis campanulatis mucronatis eglandulosis, germinibus compressis utrinque sulcatis. A. H. X. p. 438. t. 1. f. 4. *S. rotata* Fl. N. 1077. *Gentiana islandica* Fl. D. 343. Fl. N. 380. c)
- † 310. *S. perennis*, corollis quinquefidis, foliis radicalibus ovalibus. S. N. 266. Jacqu. Flor. Austr. III. t. 243. P. d).

GENTIANA.

- † 311. *G. purpurea*, corollis subquinquefidis campanulatis verticillatis, calycibus truncatis. S. N. 267. Fl. N. 97. Fl. D. 50. P. alp.
312. *G. Pneumonanthe*, corollis quinquefidis campanulatis, foliis linearibus. Fl. Sv. 228. Fl. N. 1108. Fl. D. 269. P. pr. hum.
313. *G. nivalis*, corollis quinquefidis infundibuliformibus, ramis unifloris alternis. Fl. Sv. 231. Fl. N. 183. Fl. D. 17. A. H. X. t. 2. f. 5. e). A. alp.

314. *G.*c) Diversissima a *S. rotata* Linn.

d) In Pomerania legit Cl. Helwig.

e) Figurae *Rottböllii*, *Oederi* et *Halleri* Stirp. Helv. t. 17. diversae speciei videntur plantas referre, sed eandem omnino plantam sistunt.

314. *G. pulchella*, corolla quinquefida infundibiliformi, tubo elongato, stylo simplici, caule simplicissimo. N. A. Holm. 4. p. 85. t. 3. f. 8. 9. f). A.
315. *G. Centaurium*, corollis quinquefidis infundibiliformibus, caule dichotomo, pistillo duplici. Fl. Sv. 232. Fl. N. 723. Fl. D. 617. A. pr. hum.
316. *G. involocrata*, corollis quinquefidis infundibiliformibus involocratis, caule acute tetragono, foliis oblongo-ovatis. A. H. X. p. 434. t. 1. f. 2. A. B. *G. aurea*, Spec. Pl. 331. Fl. N. 548. *G. quinquefolia*, Fl. N. 838. Fl. Isl. 5. Fl. D. 344. g).
317. *G. Amarella*, corollis quinquefidis hypocrateriformibus fauce barbatis. Fl. Sv. 229. Fl. N. 95. Fl. D. 328. h). A. pr.
318. *G. campestris*, corollis quadrifidis fauce barbatis, caule alato. Fl. Sv. 230. Fl. N. 96. Fl. D. 367. A. pr.
- † 319. *G. tenella*, corollis quadrifidis fauce barbatis, caule filiformi tetragono, foliis ovatis basi solutis. A. H. X. p. 436. t. 2. f. 6. *G. Königii* Fl. N. 822. G. etc. Fl. D. 318. St. Helv. 652. A. alp.
- † 320. *G. ciliata*, corollis quadrifidis margine ciliatis. S. N. 269.
 β. *G. detonsa*, A. H. X. p. 435. t. 1. f. 3. Fl. N. 733. t. 2. f. 3. 4. 5.
 γ. *G. ferrata*, quinquefida. Fl. N. 819. *G. bavarica*, Fl. Isl. 5. Fl. D. 317.
- † 321. *G. filiformis*, corollis quadrifidis imberbibus, caule dichotomo filiformi. S. N. 270. Fl. D. 324. A. pasc. ster.

ERYN-

- f) varietatem sequentis credunt alii, negat inventor *Cel. Swartz.*
- g) Vix dubium est perill. Linné hanc intellexisse et ex specimine suo *auream* dixisse.
- h) Flores inferiores saepe quadrifidi.

ERYNGIUM.

322. *E. maritimum*, foliis radicalibus subrotundis plicatis spinosis, capitulis pedunculatis, paleis tricuspидatis. Fl. Sv. 233. Fl. D. 875. P. a. mar.
 † 323. *E. campestre*, foliis radicalibus amplexicaulibus pinnato-lanceolatis. S. N. 271. Fl. D. 554. P. a.

HYDROCOTYLE.

324. *H. vulgaris*, foliis peltatis, umbellis quinquefloris. Fl. Sv. 234. Fl. Sv. 90. P. hum.

SANICULA.

325. *S. europaea*, foliis radicalibus simplicibus, fasciculis omnibus sessilibus. Fl. Sv. 235. Fl. N. 530. Fl. D. 283. P. sylv.

BUPLEURUM.

- † 326. *B. falcatum*, involucrellis pentaphyllis acutis: universali subpentaphyllo, foliis lanceolatis, caule flexuoso. S. N. 273. Weig. Suppl. 858. Jacqu. Fl. Austr. II. t. 128. P. sep.
 327. *B. tenuissimum*, umbellis simplicibus alternis pentaphyllis subtrifloris. S. N. 274. Obs. Bot. F. I. 14. A. r.

TORDYLIUM.

328. *T. Anthriscus*, umbellis confertis, foliolis ovato-lanceolatis pinnatifidis. Fl. Sv. 236. Fl. D. 919. B. r. v. agr.

DAUCUS.

329. *D. Carota*, seminibus hispidis, petiolis subtus nervosis. Fl. Sv. 237. Fl. N. 647. Fl. D. 723. B. pr. v. agr.

BUNIUM.

- † 330. *B. Bulbocastanum*, Fn. Sv. 557. Fl. N. 361. Fl. D. 220. P. a.

CONIUM.

331. *C. maculatum*, feminibus striatis. Fl. Sv. 238.
B. f. c. arg.

SELINUM.

332. *S. palustre*, sublaetescens, radice unica. Fl. Sv. 239.
Fl. N. 593. Fl. D. 257. P. pal.
333. *S. carvisolia*, caule fulcato acutangulo involuero
universali evanido, stylis erectis, petalis conniven-
tibus. S. N. 278. Weig. Suppl. 859. Fl. D. 667.
P. pr.

ATHAMANTA.

334. *A. Libanotis*, foliis bipinnatis planis, umbella he-
misphaerica, feminibus hirsutis. Fl. Sv. 240. Fl.
D. 754. P. pr.
335. *A. Oreoselinum*, foliolis divaricatis. Fl. Sv. 241.
P. pr. a.

LASERPITIUM.

336. *L. latifolium*, foliolis cordatis inciso-ferratis. Fl.
Sv. 242. Jacqu. Fl. Austr. II. t. 146. P. pr.
† 337. *L. prutenicum*, foliolis lanceolatis integerrimis:
extimis coalitis. S. N. 281. Jacqu. Fl. Austr. II.
t. 153. i). P.

HERACLEUM.

338. *H. Sphondylium*, foliolis pinnatifidis, floribus dif-
formibus. Fl. Sv. 243. Fl. N. 218. P. f. v. agr.
339. *H. angustifolium*, foliis cruciato-pinnatis, folio-
lis linearibus, corollis flosculosis. S. N. 282.
Mant. Pl. 57, k). P. pr. fylv.

LIGUSTICUM.

340. *L. Levisticum*, fons multiplicibus: foliolis superne
incisis. S. N. 283. P. f. r.
341. *L. Scoticum*, foliis biternatis. Fl. Sv. 244. Fl.
N. 254. Fl. D. 207. P. l. m. a.

ANGE-

- i) Diversum omnino ab illo Florae Austriacae.
k) *Cel. Kölpin*.

ANGELICA.

342. *A. Archangelica*, foliorum impari lobato. Fl. Sv. 245. Fl. N. 98. Fl. D. 206. P. hum. alp.

343. *A. sylvestris*, foliolis aequalibus ovato-lanceolatis ferratis. Fl. Sv. 246. Fl. N. 217. P. pr. hum. sylv.

SIUM.

345. *S. latifolium*, foliis pinnatis, umbella terminali. Fl. Sv. 247. Fl. N. 651. Fl. D. 246. P. aqu.

346. *S. angustifolium*, foliis pinnatis inaequaliter argutaeque ferratis auriculatis, umbellis oppositifoliis pedunculatis. Obs. Bot. F. I. p. 14. l). S. N. 284. *S. nodiflorum*, Ros. Obs. p. 14. Fl. Sv. 248. Weig. Obs. p. 27. St. Helv. 778. Fl. D. 247. P. aqu.

† 347. *S. nodiflorum*, foliis pinnatis, umbellis axillaribus sessilibus. S. N. 284. Morif. 3. p. 383. S. IX. t. 5. f. 3. P. aqu.

† 348. *S. Falcaria*, foliolis linearibus decurrentibus connatis. S. N. 284. Weig. Suppl. 861. P. pr.

SISON.

349. *S. inundatum*, repens, umbellis bifidis. Fl. Sv. 249. Fl. D. 89. A. l. lac.

OENANTHE.

350. *Oe. fistulosa*, stolonifera, foliis caulinis pinnatis filiformibus fistulosis. Fl. Sv. 250. Fl. D. 846. P. riv.

351. *Oe. crocata*, foliis omnibus multifidis obtusis subaequalibus. Fl. Sv. 251. m). P. riv.

† 352. *Oe. pimpinelloides*, foliolis radicalibus cuneatis fissis: caulinis integris linearibus longissimis simplicibus. S. N. 286. Jacqu. Fl. Austr. IV. t. 39. m). P.

E 2

PHEL-

l) Quo jure me hanc cum sequente confusione dici poterat?

m) Juxta Lundam olim observata ubi a b. Leche fuit, jam desideratur.

n) Cel. Hellwig.

68 PENTANDRIA. DIGYNIA. Phellandrium.

PHELLANDRIUM.

353. *P. aquaticum*, foliorum ramificationibus divaricatis. Fl. Sv. 252. B. l. lac.

AETHUSA.

354. *Ae. Cynapium*, foliis conformibus. Fl. Sv. 254. Fl. N. 915. A. c.

CICUTA.

355. *C. virosa*, umbellis oppositifoliis, petiolis marginatis obtusis. Fl. Sv. 253. Fl. N. 42. t. 2. Fl. D. 208. P. aqu.

SCANDIX.

356. *S. odorata*, feminibus fulcatis angulatis. S. N. 287. Fn. Sv. 257. Fl. N. 370. P. a. alp.
357. *S. Petten*, feminum rostro longissimo. S. N. 287. Fh. Sv. 557. Fl. D. 844. A. c.
358. *S. Cerefolium*, feminibus nitidis ovato-fubulatis, umbellis sessilibus lateralibus. Fl. Sv. 255. A. c. r.
359. *S. Anthriscus*, feminibus ovatis hispidis, corollis uniformibus, caule laevi. Fl. Sv. 256. Fl. D. 863. A. r. f.
360. *S. infesta*, femine exteriori hispido, umbellulis confertissimis hemisphaericis. S. N. 287. Jacqu. Fl. Austr. I. t. 46. o).

CHAEROPHYLLUM.

361. *C. sylvestre*, caule laevi striato, geniculis tumidiusculis. Fl. Sv. 257. Fl. N. 294. A. r. v. agr. f.
362. *C. temulum*, caule scabro, geniculis tumidis. Fl. Sv. 258. Fl. D. 918. B. r.
† 363. *C. bulbosum*, caule laevi geniculis tumido, basi hirta. S. N. 288. Fl. N. 460. Jacqu. Fl. Austr. I. t. 63. B.

IMPERATORIA.

364. *I. Ostruthium*, S. N. 289. Fl. Sv. p. 464. Fl. Isl. 5. P. alp.

SESELI.

- o) Cel. Afzelius.

SESELI.

365. *S. montanum*, petiolis ramiferis, membranaceis oblongis integris, foliis caulinis angustissimis. S. N. 289. 6*). P. pr. mont.

PASTINACA.

366. *P. sativa*, foliis pinnatis subbipinnatisque auriculatis. Obs. Bot. Fasc. V. p. 3. Fl. Sv. 259. Fl. N. 290. B. ad urb.

CARUM.

367. *C. Carvi*, Fl. Sv. 260. Fl. N. 352. B. P. pr. f.

PIMPINELLA.

368. *P. Saxifraga*, foliis inferioribus simpliciter pinnatis, foliolo terminali trilobo. Obs. Bot. Fasc. III. p. 29. Fl. N. 353. P. pr. pasc.

β. *P. magna*. Pr. Flor. Scand. 326. P. fol. infer. latis subcordatis auriculatis superioribus subbipinnatis.

γ. P. fol. inf. ovato-lanceolatis auriculatis, supremis subbipinnatis, pinnis lineari-incisis.

δ. P. fol. rad. subrotundis ferratis, caulinis linearibus. Fl. Sv. 261. Fl. D. 669. p).

369. *P. dissecta*, foliis omnibus pinnatis. pinnis multipartitis, segmentis subfalcatis acutis. Obs. Bot. Fasc. III. p. 30. t. 2. P. pr. viae.

APIUM.

370. *A. graveolens*, foliis caulinis cuneiformibus umbellis sessilibus. Fl. Sv. 262. Fl. N. 496. Fl. D. 790. B. pasc. m.

E 3

AEGO-

6*) Cel. Afzelius.

p) Ab hac nullo modo differt Pimpinella nigra, nisi quod succo nigrescente gaudeat radix. Crescit ad Malmogiam.

AEGOPODIUM.

371. *Ae. Podagraria*, foliis caulinis summis ternatis.
Fl. Sv. 263. Fl. N. 304. Fl. D. 670. P. c. pr. r. fylv.

3. T R I G Y N I A.

VIBURNUM.

372. *V. Opulus*, foliis lobatis, petiolis glandulosis.
Fl. Sv. 264. Fl. N. 7. Fl. D. 661. F. pr. ul.

SAMBUCUS.

373. *S. nigra*, cymis quinquepartitis, caule arboreo.
Fl. Sv. 265. Fl. N. 407. Fl. D. 545. F. ad dom. f.
374. *S. Ebulus*, cymis tripartitis, stipulis foliaceis, caule
herbaceo. Fl. Sv. 266. P.

TAMARIX.

375. *T. germanica*, floribus decandris. S. N. 296. Fl.
N. 152. Fl. D. 234. q). F. inund.

CORRIGIOLA.

- † 376. *C. littoralis*. S. N. 297. Fl. D. 334. A. l. a.

ALSINE.

377. *A. media*, petalis bipartitis, foliis ovato-cordatis.
Fl. Sv. 267. Fl. N. 191. r). A. c. r.
α. *pentandra*. Fl. D. 525.
β. *decandra*. Fl. D. 438.

- † 378. *A. viscosa*, caule paniculato foliisque linearibus
pubescentibus, petalis capsulisque calyce pube-
scente brevioribus. Schr. Lips. p. 30. F. D. 389. A.

4. TE-

q) Monogyna est secundum *Gunnerum*.

r) *Holsteo* commodè jungitur hæc species, observante *Cal.*
Swarze Obs. Bot. 118.

3. T E T R A G Y N I A.

PARNASSIA.

379. *P. palustris*, Fl. S. 268. Fl. N. 231. Fl. D. 584.
P. pr. ul.

5. P E N T A G Y N I A.

STATICE.

380. *S. Armeria*, scapo simplici capitato, foliis linearibus. Fl. Sv. 269. Fl. N. 113. P. pr. apr. a.
381. *S. Limonium*, scapo paniculato tereti, foliis laevibus nerviis subtus mucronatis. Fl. Sv. 270. Fl. D. 315. P. l. m.

LINUM.

382. *L. catharticum*, foliis oppositis ovato-lanceolatis, caule dichotomo, corollis acutis. Fl. Sv. 271. Fl. N. 86. Fl. D. 851. A. a.
383. *L. Radiola*, foliis oppositis ovatis, caule dichotomo, floribus tetrandris tetragynis. Fl. Sv. 272. Fl. N. 722. Fl. D. 178. A. a.

DROSERA.

384. *D. rotundifolia*, scapis radicatis, foliis orbiculatis. Fl. Sv. 273. Fl. N. 234. A. pr. ul.
385. *D. longifolia*, scapis radicatis, foliis oblongis. Fl. Sv. 274. Fl. N. 279. A. pr. ul.

CRASSULA.

- †386. *C. rubens*, foliis fusiformibus subdepressis, cyma quadrifida foliosa, floribus sessilibus, staminibus reflexis. S.N. 306. Fl. N. 1106. Fl. D. 82. A. l. m.

72 PENTANDRIA. PENTAGYNIA. Sibbaldia.

SIBBALDIA.

387. *S. procumbens*, foliolis tridentatis. Fl. Sv. 275.
Fl. N. 107. Fl. D. 32. P. alp;

6. P O L Y G Y N I A.

MYOSURUS.

388. *M. minimus*, Fl. Sv. 276. Fl. N. 759. Fl. D. 406, s).
A. r. ficc. f.

s) Quin *Ranunculis* affinis sit, nullum est dubium, at ideo
affociari eis non debet, quum praeter numerum staminum
aliae adsint differentiae.

VI. HEXANDRIA.

I. MONOGYNIA.

ALLIUM, *Spatha* multiflora. *Umb.* congesta. *Cor.* 6-part. patens. *Capsf.* supera.

Rad. bulbosa, tunicata. *Caul.* scaposa. *Fol.* simplex, succulenta. *Flor.* capitato-umbellata. *Fulcr.* o.

LILIUM, *Flor.* nudi. *Cor.* 6-pet. campanulata *rima* longitudinali nectarifera.

Rad. bulbosa, squamata. *Caul.* foliosa. *Fol.* simplex, crassa. *Flor.* terminalis, subumbellata. *Fulcr.* bracteae.

FRITILLARIA. *Flor.* nudi. *Cor.* 6-pet. campanulata supra ungues fovea nectarifera.

Rad. bulbosa, solida. *Caul.* foliosa. *Flor.* terminalis, cornua. *Fulcr.* bracteae.

TULIPA, *Flor.* nudi. *Cor.* 6-pet. campanul. *Nest.* o. *Stylus* o.

Rad. bulbosa, solida. *Caul.* scaposa. *Fol.* simplex, radicalis. *Flor.* terminalis, solitaria. *Fulcr.* o.

ORNITHOGALUM, *Flor.* nudi. *Cor.* 6-pet. erecta, persistens, supra medium patens. *Filamenta* alterna basi dilatata.

Rad. bulbosa, solida. *Caul.* scaposa. *Fol.* simplex, radicalis. *Flor.* terminalis, corymbosa vel racemosa.

SCILLA, *Flor.* nudi. *Cor.* 6-pet. patens, decidua. *Filam.* filiformia.

Rad. bulbosa, solida. *Caul.* scaposa. *Fol.* simplex, radicalis. *Flor.* terminalis, corymbosa. *Fulcr.* bracteae.

HYACINTHUS, *Flor.* nudi. *Cor.* 6-fida, infera. *Pori* 3 melliferi ad apicem germinis.

Rad. bulbosa, tunicata. *Caul.* scaposa. *Fol.* simplex, radicalis. *Flor.* racemosa. *Fulcr.* bracteae.

ANTHERICUM. *Cor.* 6-pet. plana.

Rad. stolonifera vel fibrosa. *Caul.* scaposa. *Fol.* simplex, radicalis. *Flor.* terminalis, racemosa. *Fulcr.* bracteae.

ASPARAGUS. *Cor.* 6-partit. laciniis alternis reflexis.

Bacca 3-loc. 2-sperma.

Rad. filiformis. *Caul.* ramosa, foliosa. *Fol.* simplex, setacea, fasciculata. *Flor.* axillaris, pendunculata. *Fulcr.* stipulae.

CONVALLARIA. *Cor.* 6-fida. *Bacca* 3-loc. polysperma, maculosa.

Rad. horizontalis, nodosa, l. repens. *Caul.* herbacea, varia. *Fol.* simplex, diversa. *Flor.* subracemosa. *Fulcr.* o. l. bracteae.

ACORUS. *Spadix* cylindricus tectus floribus. *Cor.* 6-pet. nuda. *Styli* o. *Capsf.* 3-locul.

Rad. horizontalis, articulata. *Caul.* scapus foliaceus. *Fol.* simplex, radicalis, ensiformis. *Flor.* spicata e sinu folii. *Fulcr.* o.

JUNCUS. *Cal.* 6-phyll. *Cor.* o. *Capsf.* 1-loc.

Rad. fibrosa. *Caul.* culmosa, sarcta. *Fol.* graminea, aut longe subulata. *Flor.* varia. *Fulcr.* bracteae.

BERBERIS. *Cal.* 6-phyll. *Cor.* 6-pet. glandulis 2 ad ungues. *Styl.* o. *Bacca* 2-sperma.

Rad. lignosa. *Caul.* fruticosa, aculeis furcatis. *Fol.* simplex, alterna, ciliata. *Flor.* racemosa, axillaris. *Fulcr.* bracteae.

PEPLIS. *Cal.* campanul. 12-dentat. *Pet.* 6 calyci inserta l. o. *Capsf.* 2-loc.

Rad. repens. *Caul.* procumbens, geniculata. *Fol.* simplex, opposita, conjugata. *Flor.* axillaris, sessilis, solitaria. *Fulcr.* o.

2. DIGYNIA.

Rumex digynus v. Hexandr. trigyn.

Polygonum Hydropiper et Persicaria v. Octandr. trigyn.

3. TRI-

3. TRIGYNIA.

RUMEX. *Cal.* 3-phyll. *Cor.* 3-pet. *Styli* capillares.
Sem. 1. triquetrum.

Rad. ramosa. *Caul.* herbacea, foliosa. *Fol.* alternata, simplex. *Flor.* racemosa. *Fulcr.* o.

SCHEUCHZERIA. *Cal.* 6-part. *Cor.* o. *Styli* o. *Capsf.* 3. inflatae, 1-spermae.

Rad. repens, geniculata. *Caul.* culmosa, foliosa. *Fol.* simplex, graminea. *Flor.* racemosa, *Fulcr.* o.

TRIGLOCHIN. *Cal.* 3-phyll. *Cor.* 3-pet. calyciformis. *Styli* o. *Capsf.* basi dehiscens.

Rad. fibrosa. *Caul.* scaposa. *Fol.* simplex, filiformis, radicalis. *Flor.* spicata. *Fulcr.* o.

Anthericum calyculatum v. Hexandr. monogyn.

4. POLYGYNIA.

ALISMA. *Cal.* 3-phyll. *Cor.* 3-pet. *Sem* plurima.

Rad. fibrosa. *Caul.* scaposa. *Fol.* simplex, radicalis, nervosa. *Flor.* varia. *Fulcr.* bracteae.

VI. HEXANDRIA.

I. MONOGYNIA.

ALLIUM.

389. *A. Scorodoprasum*, caule planifolio bulbifero, foliis crenulatis, vaginis ancipitibus, staminibus tricuspидatis. Fl. Sv. 278. P. pr. sylv.
390. *A. arenarium*, caule planifolio bulbifero, vaginis teretibus, spatha mutica, staminibus tricuspидatis. Fl. Sv. 279. Fl. D. 290. P. agr. a.
391. *A. carinatum*, caule planifolio bulbifero, staminibus subulatis. Fl. Sv. 281. Ros. Obs. 20. P. pr. a.
392. *A. oleraceum*, caule teretifolio bulbifero, foliis scabris semiteretibus subtus fulcatis, staminibus simplicibus. Fl. Sv. 280. Fl. N. 430. B.
393. *A. urfinum*, scapo nudo subtriquetro, foliis lanceolatis petiolatis, umbella fastigiata. Fl. Sv. 277. Fl. N. 432. Fl. D. 757. P. sylv.
394. *A. schoenoprasum*, scapo nudo adaequante, folia teretia subulato-filiformia. Fl. Sv. 282. Fl. N. 429. Fl. D. 971. P. l. mar.

LILIUM.

395. *L. bulbiferum*, foliis sparsis, corollis campanulatis erectis intus scabris. S. N. 324. Fn. Sv. 557. P. sylv.
396. *L. Martagon*, foliis verticillatis, floribus reflexis, corollis revolutis. S. N. 324. t). P.

FRITILLARIA.

397. *F. Meleagris*, foliis omnibus alternis, caule unifloro. Fl. Sv. 283. Fl. D. 972. P. pr.

TULIPA.

- 2) *Cel. Afzelius*, utraque ex hortis olim aufugo, ut etiam *Ornithogalum nuzans*, *umbellatum* etc.

TULIPA.

398. *T. sylvestris*, flore subnutante, filamentis basi lanatis, foliis lanceolatis. Fl. Sv. 284. Fl. D. 375. P. pr.

ORNITHOGALUM.

399. *O. luteum*, scapo anguloso diphylo, pedunculis umbellatis simplicibus. Fl. Sv. 285. Fl. N. 79. Fl. D. 378. P. n. c.
400. *O. minimum*, scapo angulato diphylo, pedunculis umbellatis ramosis. Fl. Sv. 286. Fl. N. 1114. Fl. D. 612. P. n. c.
401. *O. umbellatum*, floribus corymbosis, pedunculis scapo altioribus, filamentis basi dilatatis. S. N. 328. A. Holm. 1769. p. 246. P. pr.
402. *O. nutans*, floribus secundis pendulis, nectario stamineo campaniformi. S. N. 328. A. Holm. 1769. p. 246. Fl. D. 912. P. pr.

SCILLA.

403. *S. italica*, racemo conico oblongo. S. N. 328. Obs. Bot. Fasc. I. p. 15. P. pr. hort.
- † 404. *S. bifolia*, floribus erectiusculis unilateralibus. Obs. Bot. Fasc. VI. p. 27. S. N. 329. Fl. N. 1068. Fl. D. 568. P.

ANTHERICUM.

405. *A. ramosum*, foliis planis, scapo ramoso, corollis planis, pistillo recto. Fl. Sv. 289. Jacq. Fl. Austr. II. t. 161. P. a.
406. *A. Liliago*, foliis planis, scapo simplicissimo, corollis planis, pistillo declinato. Fl. Sv. 290. Fl. D. 616. P. agr. a.

78 HEXANDRIA. MONOGYNIA. Asparagus.

407. *A. ossifragum*, foliis ensiformibus, filamentis lanatis. Fl. Sv. 287. Fl. N. 136. Fl. D. 42. u). P. pr. hum.
 408. *A. calyculatum*, foliis ensiformibus, perianthiis trilobis, filamentis glabris, floribus trigynis. Fl. Sv. 288. Fl. L. t. 10. f. 3. Fl. N. 190. Fl. D. 36. P. pr. ul.

ASPARAGUS.

409. *A. officinalis*, caule herbaceo tereti erecto, foliis fetaceis, stipulis paribus. Fl. Sv. 291. Fl. D. 805. P. pr. sylv.

CONVALLARIA.

410. *C. majalis*, scapo nudo. Fl. Sv. 292. Fl. N. 179. Fl. D. 854. P. pr. sylv.
 411. *C. verticillata*, foliis verticillatis. Fl. Sv. 293. Fl. N. 181. Fl. D. 86. P. pr. m.
 412. *C. Polygonatum*, foliis alternis amplexicaulibus, caule ancipiti, pedunculis axillaribus subunifloris. Fl. Sv. 294. Fl. N. 248. Fl. D. 377. P. pr. n. sylv. mont.
 413. *C. multiflora*, foliis alternis amplexicaulibus, caule tereti, pedunculis axillaribus multifloris. Fl. Sv. 295. Fl. N. 249. Fl. D. 152. P. pr. n. sylv. mont.
 414. *C. bifolia*, foliis cordatis, floribus tetrandris. Fl. Sv. 296. Obs. Bot. Fasc. I. p. 18. Fl. N. 180. Fl. D. 291. P. sylv.

HYA-

- *) De hujus et sequentis cum Anthericis affinitate confer Spec. Plant. ed. 2. p. 447. *Anthericum calyculatum* Svecicum et *Pseudoasphodelus Jacquinii* omnino diversae videntur ex specimenibus siccis collatis. Ceterum omnino ab Antherici genere divelli debent, ac descriptionem Anthericorum in Flora Capensi ill. Thunberg, occurrentium expectandam puto.

HYACINTHUS.

415. *H. botryoides*, corollis globosis uniformibus, foliis canaliculato-cylindricis strictis. S. N. 336. v). P.

ACORUS.

416. *A. Calamus*. Fl. Sv. 297. P. qu.

JUNCUS.

* *culmis nudis*.

- † 417. *J. austus*, culmo subnudo tereti submucronato, panicula terminali, involucro diphylo spinoso. S. N. 340. Moris. 3. f. 8. t. 10. f. 15. w). P. pal. m.

418. *J. conglomeratus*, culmo nudo stricto, capitulo laterali. Fl. Sv. 298. Fl. N. 473. Leers Herb. t. 13. f. 1. x). P. ul.

419. *J. effusus*, culmo nudo stricto, panicula laterali. Fl. Sv. 299. Fl. N. 516. Leers Herb. t. 13. f. 2. y). P.

α. floribus pedunculatis. Fl. Lapp. 117. qu.

β. floribus sessilibus. Fl. Lapp. 116. z). alp.

γ. culmo non facto. Obs. Bot. Fasc. I. p. 16. riv.

420. *J. inflexus*, culmo nudo, apice membranaceo incurvo, panicula laterali. S. N. 340. Leers Herb. t. 13. f. 3. P. foss. a.

421. *J. filiformis*, culmo nudo filiformi nutante, panicula laterali. Fl. Sv. 300. Fl. N. 935. Leers Herb. t. 13. f. 4. P. pal. sylv.

422. *J.*

v) Cl. Afzelius.

w) In Pomerania Cl. Helwig.

x) flores triandri. Leers, Ehrhart.

y) flores triandri Ehrhart, hexandri Leers, Roth.

z) Diversus hic videtur, at ex specimine sicco determinare speciem non potui: Icon Scheuchz. itin. alp. erronea. Accuratam determinationem dabit Cel. Vahl.

80 HEXANDRIA. MONOGYNIA. *Juncus*.

422. *J. trifidus*, culmo nudo, foliis floribusque tribus terminalibus. Fl. Sv. 301. Fl. N. 484. Fl. D. 107. P. alp.

423. *J. squarrosus*, culmo nudo, foliis setaceis, capitulis glomeratis aphyllis. Fl. Sv. 302. Fl. N. 936. Fl. D. 430.

**** culmis foliosis.**

424. *J. capitatus*, culmis foliisque filiformibus sulcatis, capituli terminalis involucre pentaphyllo inaequali. *J. capitatus* Weig. Obs. p. 28. t. 2. f. 5. Ehrh. Beytr. 2. p. 78. A. a. inund.

α. capitulo unico terminali.

β. capitulo uno alterove pedunculato ex basi capituli terminalis.

425. *J. setifolius*, culmis foliisque filiformibus, capitulis sessilibus pedunculatisque, floribus acuminatis. Ehrh. Calam. 86. St. Helv. 1320. Moris. 3. f. 8. t. 9. f. 3. a). A. l.

426. *J. articulatus*, foliis nodoso-articulatis compressis, petalis acutis. Fl. Sv. 303. Fl. N. 643. P. pal. l. lac.

α. *aquaticus* foliis compressis, glumis mucronatis. St. Helv. 1321.

β. Gramen junceum folio articulato cum utriculis. Bauh. prodr. p. 12. b).

427. *J. bulbosus*, foliis linearibus canaliculatis, capsulis obtusis. Fl. Sv. 304. Fl. N. 567. Fl. D. 431. c). P. pr. hum.

428. *J.*

a) Similis variet. β. Junci capitati, at flores minus acuminati et pedunculi magis divaricati, modo haec sufficient. Vide-tur *Cl. Roth* hunc cum suo *uliginoso* confudisse.

b) Cum post foenifecium caulis palmaris exsurgat, cui raro capitula corymbacea insident, sed utriculi unciales paleacei in capillamenta fissi, modo virides, modo purpurascens etc. *C. Bauh.* Prodr. p. 12. c. icone.

c) Quae a *Müllero* tab. 817 pingitur vix hujus est varietas,

428. *J. bufonius*, culmo dichotomo, foliis angulatis, floribus solitariis sessilibus, petalis lineari-acutis, capsulis oblongis. Fl. Sv. 305. Fl. N. 502. Leers Herb. t. 13. f. 8. A. inund. v.
β. viviparus. St. Helv. 1319.
γ. pumilus, digitalis. Obs. Bot. Fasc. I. p. 16.
429. *J. Tenageja*, culmo ramoso paniculato, floribus solitariis sessilibus, petalis ovato-oblongis, capsulis globosis. Suppl. Pl. 208. Vaill. Par. t. 20. f. 1. A. inund. a.
430. *J. supinus*, culmo subdichotomo, foliis canaliculatis filiformibus, capitulo subtrifloro. Mönch Hass. 296. t. 5. A. pr. pal.
431. *J. stygius*, culmo simplici, foliis setaceis depressiusculis, pedunculis geminis terminalibus, glumis solitariis subbifloris. Spec. Plant. 467. P. pal.
432. *J. biglumis*, floribus terminalibus subgeminis, bractea altera floribus longiore acuminata. Skr. Nat. Selfk. 2. p. 38. Fl. Sv. 307. Am. Acad. 2. t. 3. f. 3. Fl. N. 480. Fl. D. 120. A. H. X. t. 2. f. 7. P. alp.
433. *J. triglumis*, floribus terminalibus subternis, bracteis duabus ovatis flores subaequantibus. Skr. Nat. Selfk. 2. p. 38. Fl. Sv. 306. Fl. L. t. 10. f. 5. Fl. N. 482. Fl. D. 132. A. H. X. t. 2. f. 8. P. alp.
- 433*. *J. vernalis*, foliis planis pilosis, paniculae simplicis floribus solitariis sessilibus. Ehrh. Beytr. 6. p. 137.
J. pilosus L. Sp. Pl. 468. α. Morif. 3. f. 8. t. 9. f. 1. Leers Herb. t. 13. f. 10. Fl. N. 497. P. sylv.
- 434*. *J. parviflorus*, foliis planis nudis, paniculae ramosae floribus solitariis pedicellatis. Ehrh. Beytr. 6. p. 139. *J. pilosus* L. Sp. Pl. 468. γ. Fl. Sv. 497. β. Fl. L. 424. Scheuchz. Agr. t. 6. P. sylv.

434. *J. maximus*, foliis planis pilosis, paniculae ramosae floribus subternis subsessilibus. Ehrh. Beytr. 6. p. 139. *J. pilosus* L. Sp. Pl. 468. *δ*. Morif. 3. f. 8. t. 9. f. 2. Fl. D. 441. P. sylv.
- 435*. *J. multiflorus*, foliis planis nudis, culmo basi folioso, corymbo subramoso, capitulis multifloris terminalibus axillaribusque. *Juncus* Hall. St. Helv. 1329. ? *d*) P. sylv.
435. *J. campestris*, foliis planis subpilosis, spicis sessilipedunculatisque. Fl. Sv. 309. Fl. N. 323. P. pr. sylv.
- β*. *Lapponicus*. Fl. Sv. 10. f. 2.
- γ*. *tenuis*, capitulo unico vel duplici pedunculato. Obs. Bot. Fasc. I. p. 16. *J. pediformis* Villars Fl. Delph.
- δ*. *strictus*. Gr. hirsutum capitulo globofo. Mor. 3. f. 8. t. 9.
436. *J. spicatus*, foliis planis, spica racemosa nutante. Fl. Sv. 310. Fl. L. t. 10. f. 4. Fl. N. 807. Fl. D. 270. P. alp.
- † 437. *J. subverticillatus*, culmo subdichotomo, foliis subulato-setaceis, floribus fasciculato-subverticillatis. Jacqu. Coll. 3. p. 51. Fl. D. 817. *e*). fōss.

BERBERIS.

438. *B. vulgaris*, pedunculis racemosis. Fl. Sv. 311. Fl. N. 424. Fl. D. 904. F. sylv.

PEPLIS.

439. *P. Portula*, floribus subapetalis. Fl. Sv. 312. Fl. N. 650. Fl. D. 64. A. inund.

3. TRI-

d) Habitus *J. pilosi*, at capitula *J. articulati*. Folium in culmo nullum. Flores acuti spadicei.

e) Varieratem *J. bulbos.* credit Müller, diversam speciem Wulffen.

2. T R I G Y N I A.

RUMEX.

440. *R. crispus*, floribus hermaphroditis: valvulis integris graniferis, foliis lanceolatis: inferioribus oblongo-cordatis undulatis acutis. Fl. Sv. 314. Fl. N. 35. P. v. agr. r.
441. *R. Nemolapathum*, floribus hermaphroditis: valvulis linearibus obtusis integerrimis graniferis, verticillis remotis, ramis patentibus, foliis inferioribus cordato-lanceolatis, superioribus lanceolatis. Suppl. Pf. p. 212. *R. dubius* Fl. N. 412. Prodr. Fl. Scand. 387. *R. palustris* Weig. Suppl. 866. P. pr. sylv. pal.
442. *R. maritimus*, floribus hermaphroditis, valvulis dentatis graniferis, foliis linearibus. Fl. Sv. 313. Fl. N. 318.
443. *R. acutus*, floribus hermaphroditis, valvulis dentatis graniferis, foliis cordato-oblongis acuminatis. Fl. Sv. 316. Fl. N. 531. P. pr. r.
444. *R. obtusifolius*, floribus hermaphroditis, valvulis dentatis graniferis, foliis cordato-oblongis obtusiusculis crenulatis. S. N. 347. Fn. Sv. 557. P. pr. r.
445. *R. aquaticus*, floribus hermaphroditis, valvulis integerrimis nudis, foliis cordatis glabris acutis. Fl. Sv. 315. P. aqu.
446. *R. digynus*, floribus hermaphroditis digynis. Fl. Sv. 317. Fl. N. 36. Fl. D. 14. P. alp.
447. *R. Acetosa*, floribus dioicis, foliis oblongis sagittatis. Fl. Sv. 318. Fl. N. 167. P. pr.
 a. vulgaris.
 β. alpina major.
448. *R. Acetosella*, floribus dioicis, foliis lanceolato-hastatis. Fl. Sv. 319. Fl. N. 654. B. arv. c. tect.

SCHEUCHZERIA.

449. *S. palustris*. Fl. Sv. 320. Fl. L. t. 10. f. 1. Fl. N. 483.
Fl. D. 76. P. pal.

TRIGLOCHIN.

450. *T. palustre*, capsulis trilocularibus sublinearibus.
Fl. Sv. 321. Fl. N. 162. Fl. D. 490. P. pr. hum.
451. *T. maritimum*, capsulis sexlocularibus ovatis. Fl.
Sv. 322. Fl. N. 153. Fl. D. 306. P. pr. hum. m.

4. P O L Y G Y N I A.

ALISMA.

452. *A. Plantago aquatica*, foliis ovatis acutis, fructibus obtuse trigonis. Fl. Sv. 323. Fl. N. 672.
Fl. D. 561. P. aqu.
β. foliis angustioribus fere lanceolatis.
453. *A. natans*, foliis ovatis obtusis, pedunculis solitariis. Fl. Sv. 324. P. aqu.
454. *A. Ranunculoides*, foliis lineari-lanceolatis, fructibus globofo-squarrosis. Fl. Sv. 325. Obs. Bot. Fl. p. 16. Fl. D. 122. P. ? pr. a. inund.

VII. HEPTANDRIA.

I. MONOGYNIA.

TRIENTALIS. *Cal.* 7 phyll. *Cor.* 7-part. plana. *Bacca*
ficca 1-locul.

Rad. fibrosa. *Caul.* herbacea, simplex, biflora.

Fol. simplex, subverticillata ad apicem caulis.

Flor. terminalis, solitaria, longe peduncu-
lata. *Fulcr.* squama caulis.

VII. HEPTANDRIA.

I. M O N O G Y N I A.

TRIENTALIS.

455. *T. europaea*, foliis lanceolatis integerrimis. Fl.
Sv. 326. Fl. N. 230. Fl. D. 84. P. sylv.

VIII. OCTANDRIA.

I. MONOGYNIA.

OENOTHERA. *Cal.* 4-fid. *Cor.* 4-pet. *Capsf.* cylindr. infera. *Sem.* nuda.

Rad. ramosa. *Caul.* herbacea, foliosa, obliqua. *Fol.* simplex, altera, decurrens. *Flor.* sessilis, axillaris. *Fulcr.* o.

EPILOBIUM. *Cal.* 4-fid. *Cor.* 4-pet. *Capsf.* 4-gona, infera. *Sem.* papposa.

Rad. ramosa. *Caul.* herbacea, foliosa. *Fol.* simplex, varia. *Flor.* axillaris, subterminalis. *Fulcr.* o.

VACCINIUM. *Cal.* 4-dent. superus. *Cor.* 1-pet. *Bacca* 4-locul. polysperma.

Rad. lignosa. *Caul.* fruticulosa. *Fol.* simplex. *Flor.* varia. *Fulcr.* o.

ERICA. *Cal.* 4-phyll. *Cor.* 4-fid. *Antherae* bifidae. *Capsf.* 4-locul.

Rad. lignosa. *Caul.* fruticulosa. *Fol.* simplex, opposita l. verticillata. *Flor.* racemosa. *Fulcr.* bracteae.

DAPHNE. *Cal.* o. *Cor.* 4-fid. marcescens, stamina includens. *Bacca* 1-sperma.

Rad. lignosa. *Caul.* fruticosa. *Fol.* simplex, sparsa. *Flor.* subspicata, terna. *Fulcr.* o.

Monotropa v. Decandria monogyn.

2. TRIGYNIA.

POLYGONUM. *Cal.* o. *Cor.* 5-part. *Sem.* 1. angulatum.

3. TETRAGYNIA.

PARIS. *Cal.* 4-phyll. *Cor.* 4-pet. linearis. *Bacca* 4-loc.
Rad. repens. *Caul.* herbacea, simplicissima,
 tetraphylla. *Fol.* simplex, quaterna. *Flor.*
 terminalis, solitaria. *Fulcr.* o.

ADOXA. *Cal.* 2-fid. inferus. *Cor.* 4-5-fida. *Bacca* 4-5-loc.
 calyce coalita.
Rad. dentata. *Caul.* herbacea, simplex, diphylla.
Fol. opposita, composita. *Flor.* terminalis,
 capitata. *Fulcr.* o.

ELATINE. *Cal.* 4-phyll. *Cor.* 4-pet. *Caps.* 4-locul.
 4-valv. depressa.
Rad. fibrosa. *Caul.* inferne demersa, radicans.
Fol. opposita, verticillata. *Flor.* solitaria,
 axillaris, breviter pedicellata. *Fulcr.* o.

VIII. OCTANDRIA.

I. M O N O G Y N I A.

OENOTHERA.

456. *Oe. biennis*, foliis ovato-lanceolatis planis, caule muricato-villoso. S. N. 358. Fl. D. 446. B. pr.

EPILOBIUM.

457. *E. angustifolium*, foliis sparsis lineari-lanceolatis, floribus inaequalibus. Fl. Sv. 327. Fl. N. 57. Fl. D. 289. P. sylv. nem. f.

- †458. *E. Halleri*, foliis sparsis linearibus integerrimis, floribus inaequalibus. All. Fl. Pedem. 1016, Stirp. Helv. 1001. f). P. alp.

- †459. *E. frigidum*, foliis oppositis alternisque lanceolatis subdentatis glabris, floribus axillaribus solitariis. *E. corymbosum* A. H. X. p. 440. t. 8. f. 23. *E. latifolium* Fl. N. 1066. Fl. D. 565. g). P.

460. *E. hirsutum*, foliis oppositis lanceolatis ferratis decurrenti-amplexicaulibus. Sp. Pl. 494. α. Fl. Sv. 328. β. Fl. D. 326. P. sylv. hum.

461. *E. parviflorum*, foliis oppositis alternisque lanceolatis ferratis sessilibus cauleque villosis. Schr. Lips. p. 146. *E. hirsutum* Fl. Sv. 328. α. Fl. D. 347. *E. villosum* H. Kew. 2. p. 5.

- †462. *E. obscurum*, foliis oppositis alternisque lanceolatis glabris ferratis basi decurrentibus. Schr. Lips. p. 147. h). P.

F 5

463. *E.*

f) Specimen possideo ex Islandia allatum. Nomen *Cel. Allioni* non retinui, quia *pseudo-Lysimachium purpureum minus* Dod. certe alia est planta.

g) Huic omnino similem ex alpinis Rhaeticis possideo siccam, at flores in hoc minores et acuti, ac folia nonnihil angustiora, quae forte est Dodonaei planta.

h) In Pomerania Cl. *Helwig*.

90 OCTANDRIA! MONOGYNIA. Vaccinium.

463. *E. montanum*, foliis oppositis ovatis denticatis. Fl. Sv. 329. Fl. N. 56. Fl. D. 922. P. f. lap.
464. *E. roseum*, foliis oppositis alternisque ovato-lanceolatis petiolatis ferratis glabris. Schr. Lipf. p. 147. Obs. Bot. Fasc. I. p. 16. P. riv.
- † 465. *E. tetragonum*, foliis lanceolatis denticulatis: imis oppositis, caule tetragono. S. N. 359. Weig. Suppl. 868. Fl. Isl. 6. P.
466. *E. palustre*, foliis oppositis lanceolatis integerrimis, petalis emarginatis, caule erecto. Fl. Sv. 330. Fl. N. 71. P. hum.
 β. *alpinum*, ramoso florens. Fl. L. 148. alp.
467. *E. alpinum*, foliis oppositis ovato-lanceolatis integerrimis i), filiquis sessilibus, caule repente. Fl. Sv. 331. Fl. N. 701. Fl. D. 322. P. alp.

VACCINIUM.

468. *V. Myrtillus*, pedunculis unifloris foliis ferratis ovatis deciduis, caule angulato. Fl. Sv. 333. Fl. N. 292. Fl. D. 974. S. sylv. hum.
469. *V. uliginosum*, pedunculis unifloris, foliis integerrimis obovatis obtusis laevibus. Fl. Sv. 332. Fl. N. 291. Fl. D. 231. S. pr. ul.
470. *V. Vitis idaea*, racemis terminalibus nutantibus, foliis obovatis revolutis integerrimis subtus punctatis. Fl. Sv. 334. Fl. N. 109. Fl. D. 40. S. pr. ster.
471. *V. Oxycoccus*, foliis integerrimis revolutis ovatis, caulibus repentibus filiformibus nudis. Fl. Sv. 335. Fl. N. 72. Fl. D. 80. k). S. pal.

ERICA.

- i) Folia certe in meis exemplaribus leviter denticulata sunt.
 k) Corolla quadripartita revoluta, non tetrapetala.

ERICA.

472. *E. vulgaris*, antheris aristatis, corollis campanulatis subaequalibus, calycibus duplicatis, foliis oppositis sagittatis. Fl. Sv. 336. Fl. N. 15. Fl. D. 677. S. pasc. pr. apr.
473. *E. Tetralix*, antheris aristatis, corollis ovatis, stylo incluso, foliis quaternis ciliatis, floribus capitatis. Fl. Sv. 337. Fl. N. 369. Fl. D. 81. S. pasc. fabul. ster. pal.
- † 474. *E. cinerea*, antheris cristatis, corollis ovatis, stylo subexserto, stigma capitato, foliis ternis. S. N. 366. Fl. N. 367. Fl. D. 38. S. ster.

DAPHNE.

475. *D. Mezereum*, floribus sessilibus ternis caulinis, foliis lanceolatis deciduis. Fl. Sv. 338. Fl. N. 22. Fl. D. 268. F. fylv.

2. T R I G Y N I A.

POLYGONUM.

476. *P. Bisflorta*, caule simplicissimo monostachyo, foliis ovatis in petiolum desinentibus. S. N. 376. Fl. D. 421. P. pr. ul.
477. *P. viviparum*, caule simplicissimo monostachyo, foliis lanceolatis. Fl. Sv. 340. Fl. N. 9. Fl. D. 13. P. pr. alp. mont.
478. *P. amphibium*, floribus pentandris semidigynis, spica ovata. Fl. Sv. 341. Fl. N. 724. P.
 α . *terrestre* erectum, staminibus corolla longioribus. v. agr. arg.
 β . *aquaticum* fluitans, staminibus corolla brevioribus. Fl. D. 282. aqu.

479. *P.*

479. *P. Hydropiper*, floribus hexandris semidigynis, foliis lanceolatis, stipulis submuticis. Fl. Sv. 343. Fl. N. 406. P. inund. hum.
480. *P. Perficaria*, floribus hexandris digynis, spicis ovato-oblongis, foliis lanceolatis, stipulis ciliatis. Fl. Sv. 342. Fl. N. 601. A. r. c. agr.
 α. *latifolia*,
 β. *angustifolia*,
 γ. *maculata* Fl. D. 702.
 δ. *pufilla*,
 ε. *tomentosa* subtus.
481. *P. aviculare*, floribus octandris trigynis axillaribus, foliis lanceolatis, caule procumbente herbaceo. Fl. Sv. 339. Fl. N. 309. Fl. D. 803. A. c. r. v.
 α. *latifolium*,
 β. *angustifolium*,
 γ. *brevifolium*,
 δ. *purpurascens*,
 ε. *aphyllum*.
482. *P. Fagopyrum*, foliis cordato sagittatis, caule erectiusculo inermi, seminum angulis aequalibus. Fl. Sv. 345. A. agr.
483. *P. Convolvulus*, foliis cordatis, caule volubili angulato, floribus obtusis. Fl. Sv. 344. Fl. N. 34. Fl. D. 744. A. c. n.
- † 484. *P. dumetorum*, foliis cordatis, caule volubili laevi, floribus carinato-alatis. S. N. 379. Fl. D. 756. Weig. Suppl. 869. A. fylv.

3. T E T R A G Y N I A.

PARIS.

485. *P. quadrifolia*, Fl. Sv. 346. Fl. N. 129. Fl. D. 139. P. fylv.

ADOXA.

ADOXA.

486. *A. Moschatellina*, Fl. Sv. 347. Fl. N. 616. Fl. D. 94. P. n.

ELATINE.

487. *E. Hydropiper*, foliis oppositis. Fl. Sv. 348. Fl. N. 1037. Fl. D. 156. A. l. lac.
488. *E. Alfinastrum*, foliis verticillatis. Fl. Sv. 349. qu.

X. DECANDRIA.

I. MONOGYNIA.

MONOTROPA. *Cal.* 0. *Cor.* 8-10 pet. petalis exterioribus basi excavata mellifera. *Stam.* 8-10. *Caps.* 5-locul.

Rad. parasitica. *Caul.* simplex, squamosa. *Fol.* 0. *Flor.* spicata, secunda. *Fulcr.* bracteae.

LEDUM. *Cal.* 5-dent. *Cor.* 5-pet. plana. *Caps.* 5-locul. basi dehiscens.

Rad. lignosa. *Caul.* fruticulosa. *Fol.* simplex, linearis, revoluta, biennis. *Flor.* corymbosa, terminalis. *Fulcr.* 0.

ANDROMEDA. *Cal.* 5-part. *Cor.* ovata, ore 5-fido. *Caps.* 5-locul.

Rad. fibrosa. *Caul.* suffruticulosa. *Fol.* simplex. *Flor.* pedunculata, nutans. *Fulcr.* 0.

ARBUTUS. *Cal.* 5-dent. *Cor.* ovata, ore 5-fido révoluto, basi pellucida. *Bacca* 5-locul.

Rad. fibrosa. *Caul.* fruticulosa, decumbens. *Fol.* simplex, sparsa. *Flor.* racemosa. *Fulcr.* bracteae.

PYROLA. *Cal.* 5-part. *Cor.* 5-pet. *Caps.* 5-loc. angulis dehiscens.

Rad. filiformis, fibrosa. *Caul.* herbacea, angulata. *Fol.* simplex, duriuscula. *Flor.* varia. *Fulcr.* bracteae et squamae.

2. DIGYNIA.

CHRYSOSPLENIUM. *Cal.* 4-5-fidus, coloratus. *Cor.* 0. *Stam.* 8-10. *Caps.* 2-rostris, 1-locul.

Rad. fibrosa. *Caul.* herbacea, angulata, foliosa. *Fol.* simplex, reniformis. *Flor.* subumbellata. *Fulcr.* bracteae.

SAXIFRAGA. *Cal.* 5-part. *Cor.* 5-pet. *Capsf.* 2-rostris, 1-locul.

Rad. fibrosa l. granulosa. *Caul.* herbacea, varia.

Fol. simplex, crassiuscula, varia. *Flor.* subpaniculata. *Fulcr.* o.

SCLERANTHUS. *Cal.* 5-fid. *Cor.* o. *Sem.* 2. calyce inclusa.

Rad. fibrosa. *Caul.* herbacea, articulata, rigida. *Fol.* simplex, opposita, subulata. *Flor.* terminalis, fastigiata. *Fulcr.* o.

GYPSOPHILA. *Cal.* 1-phyll. campanulatus, angulatus.

Pet. 5. ovata, sessilia. *Capsf.* globosa, 1-locul.

Rad. fibrosa. *Caul.* herbacea, articulata. *Fol.* opposita, sessilis, simplex. *Flor.* terminalis, fastigiata. *Fulcr.* o.

SAPONARIA. *Cal.* 1 phyll. nudus. *Pet.* 5. unguiculata.

Capsf. oblonga 1-locul.

Rad. filiformis, repens. *Caul.* herbacea, articulata. *Fol.* simplex, nervosa, opposita. *Flor.* fastigiata. *Fulcr.* o.

DIANTHUS. *Cal.* cylindricus, 1-phyll. basi squamis 4.

Pet. 5. unguiculata. *Capsf.* cylindrica, 1-loc.

Rad. fibrosa. *Caul.* herbacea, articulata. *Fol.* simplex, linearis, opposita. *Flor.* varia. *Fulcr.* o.

3. TRIGYNIA.

CUCUBALUS. *Cal.* inflatus. *Pet.* 5. unguicul. absque corona faucis. *Capsf.* 3-locul.

Rad. fibrosa. *Caul.* herbacea, articulata. *Fol.* simplex, opposita. *Flor.* varia. *Fulcr.* o.

SILENE. *Cal.* ventricosus. *Pet.* 5. unguicul. coronata, ad faucem. *Capsf.* 3-locul.

Rad. fibrosa. *Caul.* herbacea. *Fol.* simplex, opposita. *Flor.* paniculata. *Fulcr.* o.

STELLARIA. *Cal.* 5-phyll. patens. *Pet.* 5-bipartita.
Capsf. 1-locul. polysperma.

Rad. filiformis. *Caul.* herbacea, debilis. *Fol.*
simplex, opposita. *Flor.* subpaniculata.
Fulcr. o.

ARENARIA. *Cal.* 5-phyll. patens. *Pet.* 5. integra.
Capsf. 1-loc. polysperma.

Rad. fibrosa. *Caul.* herbacea, articulata. *Fol.*
simplex, opposita. *Flor.* varia. *Fulcr.* stipulae.

Tamarix germanica vid. Pentandr. trigyn.

Alfne vid. Pentandr. trigyn.

4. PENTAGYNIA.

SEDUM. *Cal.* 5-fid. *Cor.* 5-pet. *Squamae* nectariferae
5 ad basin germinis (Elastra Hall.) *Capsf.* 5.

Rad. fibrosa. *Caul.* herbacea. *Fol.* simplex, ses-
silis, gibba, succulenta. *Flor.* corymbosa vel
cymosa. *Fulcr.* o.

OXALIS. *Cal.* 3-phyll. *Pet.* 5. unguibus connexa.
Capsf. angulis dehiscens. *Sem.* elastice prosi-
lientia.

Rad. dentata. *Caul.* scaposa. *Fol.* ternata, ra-
dicalis. *Flor.* terminalis, solitaria. *Fulcr.*
bractaeae.

AGROSTEMMA. *Cal.* 1-phyll. coriaceus. *Pet.* 5. un-
guicul. limbo indiviso obtuso. *Capsf.* 1-loc.

Rad. fusiformis. *Caul.* herbacea. *Fol.* simplex,
opposita. *Flor.* terminalis, solitaria. *Fulcr.* o.

LYCHNIS. *Cal.* 1-phyll. oblongus, laevis. *Pet.* 5. un-
guicul. limbo subbifido. *Capsf.* 5-locul.

Rad. subramosa. *Caul.* herbacea, articulata.
Fol. simplex, opposita. *Flor.* subpaniculata.
Fulcr. o.

CERASTIUM. *Cal.* 5-phyll. *Pet.* 2-fid. *Capsf.* 1-locul.
apice dehiscens.

Rad. filiformis, vel fibrosa. *Caul.* herbacea, articu-
lata. *Fol.* simplex, opposita, sessilis.
Flor. terminalis. *Fulcr.* o.

SPERGULA. *Cal.* 5-phyll. *Pet.* integra. *Capsf.* ovata
1-locul. 5-valv.

Rad. fibrosa. *Caul.* herbacea, articulata. *Fol.*
simplex, linearis. *Flor.* varia. *Fulcr.* o.

X. DECANDRIA.

I. MONOGYNIA.

MONOTROPA.

491. *M. Hypopithys*, floribus lateralibus octandris, terminali decandro. Fl. Sv. 251. Fl. D. 232. P. sylv.

LEDUM.

492. *L. palustre*, foliis linearibus, staminibus corolla longioribus. Obs. Bot. Fasc. IV. p. 26. Fl. Sv. 352. Fl. N. 333. S. pal. sylv.
 †493. *L. groenlandicum*, foliis ellipticis, staminibus corolla brevioribus. Pr. Fl. Scand. 436. Fl. N. 1067. Fl. D. 567. m).

ANDROMEDA.

494. *A. tetragona*, pedunculis solitariis lateralibus, corollis campanulatis, foliis oppositis imbricatis obtusis revolutis. Fl. Sv. 356. Fl. L. t. 1. f. 4. Fl. N. 1077. *A. tetraquetra* A. H. X. p. 443. t. 5. f. 18. S. alp.
 495. *A. hypnoides*, pedunculis solitariis terminalibus, corollis campanulatis, foliis confertis subulatis. Fl. Sv. 355. Fl. L. t. 1. f. 3. Fl. N. 667. Fl. D. 10. S. alp.
 496. *A. coerulesca*, pedunculis aggregatis, corollis ovatis, foliis sparsis linearibus obtusis planis. Fl. Sv. 354. Fl. L. t. 1. f. 5. Fl. N. 28. Fl. D. 57. A. H. X. p. 442. t. 5. f. 17. S. alp.
 497. *A. polifolia*, pedunculis aggregatis, corollis ovatis, foliis alternis lanceolatis revolutis. Fl. Sv. 353. Fl. L. t. 1. f. 2. Fl. N. 293. Fl. D. 54. S. pal. sylv.
 498. *A.*

m) Flores in hoc subpentandri. H. Kew.

498. *A. calyculata*, racemis fecundis foliaceis, corollis subcylindricis, foliis alternis lanceolatis obtusis punctatis. Fl. Sv. 357. S. sylv. hum.

ARBUTUS.

499. *A. alpina*, caulibus procumbentibus, foliis rugosis ferratis. Fl. Sv. 359. Fl. L. 161. Fl. N. 178. Fl. D. 73. S. alp.
500. *A. Uva Ursi*, caulibus procumbentibus, foliis integerrimis. Fl. Sv. 358. Fl. L. t. 6. f. 3. Fl. N. 165. Fl. D. 33. S. sylv.

PYROLA.

501. *P. rotundifolia*, staminibus adscendentibus, pistillo declinato. Fl. Sv. 360. Obs. Bot. Fasc. I. p. 17. Fl. N. 119. Fl. D. 110. P. sylv.
502. *P. minor*, floribus racemosis dispersis, staminibus pistilloque rectis. Fl. Sv. 361. Obs. Bot. Fasc. I. p. 17. Fl. N. 121. Fl. D. 55. P. sylv.
503. *P. secunda*, racemo unilaterali. Fl. Sv. 362. Obs. Bot. Fasc. I. p. 17. Fl. N. 120. Fl. D. 402. P. sylv.
504. *P. umbellata*, pedunculis subumbellatis. Fl. Sv. 363. Obs. Bot. Fasc. I. p. 17. P. sylv. acer.
505. *P. uniflora*, scapo unifloro. Fl. Sv. 364. Obs. Bot. Fasc. I. p. 17. Fl. N. 122. Fl. D. 8. P. sylv.

2. D I G Y N I A.

CHRYSO SPLENIUM.

506. *C. alternifolium*, foliis alternis. Fl. Sv. 365. Fl. N. 54. Fl. D. 366. P. hum. umbr.
† 507. *C. oppositifolium*, foliis oppositis. S. N. 411. floribus octandris Ehrh. Beytr. 3. p. 60. Fl. D. 365. P. hum.

SAXIFRAGA.

508. *S. Cotyledon*, foliis radicatis aggregatis lingulatis cartilagineo-ferratis, caule paniculato. Fl. Sv. 366. Fl. L. t. 2. f. 2. Fl. N. 13. Fl. D. 241. A. H. X. t. 2. f. 9. P. alp.

509. *S. stellaris*, foliis ferratis, caule nudo ramoso, petalis acuminatis. Fl. Sv. 367. Fl. N. 265. Fl. D. 23. Jacqu. Coll. I. p. 202. t. 13. B. alp.
β. *comosa*.

510. *S. nivalis*, foliis obovatis crenatis subsessilibus, caule nudo, floribus congestis. Fl. Sv. 368. Fl. L. t. 2. f. 5. 6. Fl. N. 545. Fl. D. 28. Skr. Nat. H. Selfk. II. p. 52. P. alp.

† 511. *S. cuneifolia*, foliis cuneiformibus obtusissimis repandis, caule nudo paniculato. S. N. 413. *S. punctata* Fl. N. 1076. A. H. X. p. 445. t. 3. f. 10. n). P.

512. *S. oppositifolia*, foliis caulinis ovatis oppositis imbricatis: summis ciliatis. Fl. Sv. 369. Fl. N. 53. Fl. D. 34. P. alp.

β. *indurata*, foliorum apicibus cartilagineis. Fl. L. p. 142. t. 2. f. 1. o).

513. *S. Hirculus*, foliis caulinis lanceolatis alternis nudis inermibus, caule erecto, pedunculo ferrugineo-tomentoso. Fl. Sv. 370. Obs. Bot. Fasc. I. p. 17. Fl. N. 335. Fl. D. 200. P. pal.

514. *S. Aizoides*, foliis caulinis lanceolatis sparsis ciliato-aculeatis, caulibus procumbentibus. Obs. Bot. F. I. p. 18. *S. Aizoides* Fl. Sv. 371. Fl. N. 541. *S. autumnalis* Fl. D. 72. Fl. N. 24. P. alp.

515. *S.*

n) Monente *Cel. Dryander* in litt. Planta Islandica haec pro *umbrosa* a me in priori editione habita est. *S. cuneata* Linn. et Scop. T. I. t. 13.

o) Plures hujus varietates extricavit *Cel. Vahl* in Nat. H. Selfk. Skr. 2. p. 51.

515. *S. granulata*, foliis caulinis reniformibus lobatis, caule ramoso, radice granulata. Fl. Sv. 372. Fl. N. 615. Fl. D. 514. P. pr. pasc. apr.
- † 516. *S. bulbifera*, foliis palmato-lobatis: caulinis sessilibus, caule ramoso bulbifero. S. N. 413. Fl. N. 1006. Fl. D. 390. P. alp.
517. *S. cernua*, foliis caulinis palmatis petiolatis, caule simplicissimo unifloro bulbifero. Fl. Sv. 373. Fl. L. t. 2. f. 4. Fl. N. 528. Fl. D. 22. P. alp.
 β. *biflora*. Fl. N. t. 8.
518. *S. rivularis*, foliis caulinis palmatis, summo florali ovato, caule simplici subbifloro. Fl. Sv. 374. Fl. L. t. 2. f. 7. Fl. N. 479. Fl. D. 118. A. H. X. t. 3. f. 13. A. riv. alp.
519. *S. tridactylites*, foliis caulinis cuneiformibus trifidis alternis, caule erecto ramoso. Fl. Sv. 375. A. a.
 β. *integrifolia*, Obs. Bot. Fasc. I. p. 18.
520. *S. petraea*, foliis cuneiformibus: radicalibus integris tridentatisque: caulinis quinquedentatis, superioribus trifidis, pedunculis subtrifloris. Vahl. Skr. Nat. H. Selfk. II. p. 10. S. N. 414. Fl. N. 427. Sax. foliis radicalibus congestis subquinquedentatis. Fl. D. 68. *S. tridactylites* Fl. N. 544. t. 9. f. 1-3. *Saxifr. adscendens* Jacqu. Coll. I. p. 197. t. 11 et 12. f. 1. 2. P. alp.
521. *S. caespitosa*, foliis radicatis aggregatis linearibus integris palmatis trifidisque, caule erecto subnudo subbifloro. Fl. Sv. 376. Fl. N. 1047. t. 7. f. 3. 4. Fl. D. 71. P. alp.
 β. *S. groenlandica*, foliis caulinis palmato-multifidis: laciniis acutis, caule erecto. S. N. 414. Fl. N. 689. t. 7. f. 1.

104 DECANDRIA. DIGYNIA. Scleranthus.

- † 522. *S. tricuspidata*, foliis lanceolatis: radicalibus congestis tricuspidatis, caulibus adscendentibus multifloris. A. H. X. p. 446. t. 6. f. 21. Fl. N. 1046. Fl. D. 976. P. pal.
523. *S. hypnoides*, foliis caulinis sublinearibus integris trifidisque, stolonibus procumbentibus multifloris. S. N. 414. Fl. N. 834. Fl. D. 348. P. alp.

SCLERANTHUS.

524. *S. annuus*, calycibus fructus patulis. Fl. Sv. 377. Fl. N. 729. Fl. D. 504. A. c.
525. *S. perennis*, calycibus fructus clausis. Fl. Sv. 378. Fl. D. 563. P. a.

GYPSOPHILA.

526. *G. fastigiata*, foliis lanceolato-linearibus obsolete triquetris laevibus obtusis secundis. Fl. Sv. 379. Gmel. Sib. 4. t. 61. f. 1. P. ficc.
527. *G. muralis*, foliis linearibus planis, calycibus aphyllis, caule dichotomo, petalis crenatis. Fl. Sv. 380. B. ficc.

SAPONARIA.

528. *S. officinalis*, calycibus cylindricis, foliis ovato-lanceolatis. S. N. 416. Ros. Obs. 24. Fl. D. 543. P. coem.

DIANTHUS.

- † 529. *D. Carthusianorum*, floribus subaggregatis, squamis calycinis ovatis aristatis tubum subaequantibus, foliis trinerviis. S. N. 417. Weig. Suppl. 872. B.
530. *D. Armeria*, floribus aggregatis, squamis calycinis lanceolatis villosis tubum aequantibus. Fl. Sv. 381. Fl. D. 230. B. pr.
531. *D. prolifer*, floribus aggregatis capitatis, squamis calycinis ovatis muticis tubum superantibus. S. N. 417. Fn. Sv. p. 557. Fl. D. 221. A. ficc.

532. *D.*

532. *D. deltoides*, floribus solitariis, squamis calycinis lanceolatis binis, corollis crenatis. Fl. Sv. 382. Fl. N. 1070. Fl. D. 577. P. pr.
533. *D. superbus*, floribus paniculatis, squamis calycinis brevibus acuminatis, corollis multifido-capillaribus, caule erecto. Fl. Sv. 383. Fl. D. 578. *D. plumarius*, Fl. N. 532. B. pr. hum.
534. *D. arenarius*, caulibus subunifloris, squamis calycinis ovatis obtusis, petalis multifidis, foliis linearibus. Fl. Sv. 384. Fl. N. 728. P. a.

3. T R I G Y N I A.

CUCUBALUS.

535. *C. Behen*, calycibus subglobosis glabris reticulato-venosis, capsulis trilocularibus: corollis subnudis. Fl. Sv. 385. Fl. N. 8. Fl. D. 914. P. a. ficc.
β. repens, maritima. Fl. D. 857. l. m.
536. *C. viscosus*, floribus-lateralibus undique decumbentibus, caule indiviso, foliis basi reflexis. Fl. Sv. 386. B.
- †537. *C. Otites*, floribus polygamis, petalis linearibus indivisis. S. N. 419. Fl. D. 518. B.

SILENE.

538. *S. nutans*, petalis bifidis, floribus lateralibus secundis, panicula nutante. Fl. Sv. 388. Fl. N. 1109. Fl. D. 242. P. pr.
539. *S. noctiflora*, calycibus decemangularibus: dentibus tubum aequantibus, caule dichotomo, petalis bifidis. Fl. Sv. 389. Weig. Suppl. 873. B. r. c.
- †540. *S. Armeria*, floribus fasciculatis fastigiatis, foliis superioribus cordatis glabris, petalis integris. S. N. 422. Fl. N. 182. Fl. D. 559. A. c.

541. *S. rupestris*, floribus erectis, petalis emarginatis, calycibus teretibus, foliis lanceolatis. Fl. Sv. 390. Fl. N. 67q. Fl. D. 4. A. pr. mont.
542. *S. acaulis*, acaulis depressa, petalis emarginatis. S. N. 422. *Cucubalus acaulis* Fl. Sv. 387. Fl. N. 117. Fl. D. 21. P. alp.
 β . *S. exscapa*, All. Fl. Ped. 1584. t. 79. f. 2. p).

STELLARIA.

543. *St. nemorum*, foliis cordatis petiolatis, paniculae pedunculis ramosis. Fl. Sv. 391. Fl. N. 232. Fl. D. 271. P. nem.
- † 544. *St. dichotoma*, foliis ovatis sessilibus, caule dichotomo, floribus solitariis, pedunculis fructiferis reflexis. S. N. 422. q). A.
545. *St. Holostea*, foliis lanceolatis ferrulatis, petalis semibifidis. Fl. Sv. 392. Fl. N. 490. Fl. D. 698. P. pr. umbr.
546. *St. graminea*, foliis linearibus integerrimis, floribus paniculatis, petalis bifidis. Fl. Sv. 493. Fl. N. 237. P.
 α . *major*. pr. ficc.
 β . *minor*. Fl. D. 414. pr. hum.
547. *St. uliginosa*, foliis lanceolatis basi subciliatis, paniculis axillaribus, petalis bifidis calycem vix aequantibus. *St. uliginosa* Roth. Fl. Germ. 2. p. 489. Murr. Pr. Gott. p. 55. *St. graminea* γ . Fl. Sv. 493. Fl. D. 415. P. ul.
548. *St. palustris*, foliis lineari-lanceolatis integerrimis, petalis bipartitis calyce majoribus. *St. graminea* β . Fl. Sv. 493. P. pr. pal.

549. *St.*

p) Skr. Nat. H. Selfk. p. 50.

q) In Pomerania Cel. Kölpin.

549. *St. cerastoides*, foliis oblongis, pedunculis subbifloris, petalis emarginatis. Fl. Sv. 394. Fl. N. 951. t. 6. f. 2. Fl. D. 92. A. H. X. t. 3. f. 12. P. alp.
550. *St. humifusa*, foliis ovatis floribus solitariis breviter pedunculatis, petalis bifidis, caulibus diffusis procumbentibus. *St. humifusa* A. H. X. p. 447. t. 4. f. 14. Fl. D. 978. N. A. Holm. 1789. p. 125. P. alp.
551. *St. biflora*, foliis subulatis, seapis subbifloris, petalis emarginatis, germinibus oblongis, calycibus striatis. Fl. Sv. 395. Fl. N. 91. Fl. D. 12. A. H. X. t. 3. f. 11. N. A. Holm. 1789. p. 42. t. 1. f. 1. P. alp.
- †552. *St. groenlandica*, caulibus decumbentibus subbifloris, foliis linearibus basi subciliatis, petalis emarginatis, fructibus globosis r).

ARENARIA,

553. *A. peploides*, foliis ovatis acutis carnosiss. Fl. Sv. 396. Fl. N. 322. Fl. D. 624. P. a. m.
554. *A. trinervia*, foliis ovatis acutis petiolatis nervosis. Fl. Sv. 397. Fl. N. 1016. Fl. D. 429. A. umbr.
- †555. *A. ciliata*, foliis ovatis nervosis ciliatis acutis. S. N. 423. Fl. N. 1078. Fl. D. 346. A. H. X. p. 448. t. 4. f. 15. Jacqu. Coll. I. p. 245. t. 15. f. 2. P. alp.
- β. *A. norvegica*. Fl. N. 1100. t. 9. f. 7.
- †556. *A. multicaulis*, foliis ovatis enerviis sessilibus acutis, corollis calyce majoribus. S. N. 423. Fl. N. 666. St. Helv. 876, t. 17. P. alp.

557. *A.*

r) Planta pollicaris. Folia subcarnosa, nonnulla e caulinis margine pilis longis ciliata. Flores, ratione plantae, magni. Calyces nudi, foliolis ovatis obtusis. Petala minime bifida vixque emarginata in meo specimine, quod dedit *Amici*. *Vahl*, an ideo potius *Arenariae* species?

108 DECANDRIA. PENTAGYNIA. *Sedum*.

557. *A. serpyllifolia*, foliis ovatis acutis sessilibus, corollis calyce brevioribus. Fl. Sv. 398. Fl. N. 564. Fl. D. 977. A. ficc.
558. *A. rubra*, foliis filiformibus, stipulis membranaceis vaginantibus. Fl. Sv. 399. Fl. N. 919. A.
α. camp. stris. ficc.
β. maritima. Fl. D. 740. l. m. 5).

4. P E N T A G Y N I A.

SEDUM.

559. *S. Telephium*, foliis planiusculis ferratis, corymbo folioso, caule erecto. Fl. Sv. 400. Fl. N. 391. P. tect. lap.
β. purpureum. Fl. D. 686.
560. *S. reflexum*, foliis subulatis sparsis basi solutis, inferioribus recurvatis. Fl. Sv. 1296. et 401. *β.* Fl. N. 475. Fl. D. 113. P. mont.
561. *S. rupestre*, foliis subulatis quinquefariis confertis basi solutis, floribus cymosis. Fl. Sv. 401. P. mont.
- † 562. *S. Oederi*, foliis semiteretibus sparsis basi solutis caule ramoso adscendente, floribus cymosis. St. Helv. 964. *S. rupestre* Fl. N. 636. Fl. D. 59. *S. saxatile* All. Fl. Ped. 1749. t. 65. f. 6. 1). P. alp.
563. *S. album*, foliis oblongis obtusis teretiusculis sessilibus patentibus, cyma ramosa. Fl. Sv. 402. Obs. Bot. Fasc. VI. p. 28. Fl. N. 705. Fl. D. 66. P. ficc.
564. *S. acre*, foliis subovatis adnato-sessilibus gibbis erectiusculis alternis, cyma trifida. Fl. Sv. 403. Fl. N. 149. P. ficc.

565. *S.*

5) Pro diversa specie habet *Rev. Wulffen*. Jacqu. Coll. III.

565. *S. sexangulare*, foliis subovatis adnato-fessilibus gibbis erectiusculis sexfariam imbricatis. Fl. Sv. 404. Obs. Bot. Fasc. I. p. 18. Fl. N. 485. B. ficc.
566. *S. annuum*, caule erecto solitario annuo, foliis ovatis fessilibus gibbis alternis, cyma recurva. Fl. Sv. 405. Obs. Bot. Fasc. I. p. 19. A. ficc.
- †567. *S. anglicum*, foliis subovatis adnato-fessilibus gibbis alternis, cyma ramosa bifida. H. Kew. II. p. 111. Fl. N. 331. u). P. ficc.
- †568. *S. villosum*, caule erecto, foliis planiusculis pedunculisque subpilosis. S. N. 431. Fl. N. 487. Fl. D. 24. A. pr. pal.

OXALIS.

569. *O. Acetosella*, scapo unifloro, foliis ternatis obcordatis, radice dentata. Fl. Sv. 406. Fl. N. 289. Fl. D. 980. P. umbr.
- †570. *O. corniculata*, caule decumbente herbaceo, pedunculis umbelliferis. S. N. 434. Fl. D. 873. A.

AGROSTEMMA.

571. *A. Githago*, hirsuta, calycibus corollam aequantibus, petalis integris nudis. Fl. Sv. 407. Fl. N. 624. Fl. D. 570. A. agr.

LYCHNIS.

572. *L. Flors Cuculi*, petalis quadrifidis, fructu subrotundo. Fl. Sv. 408. Fl. N. 124. Fl. D. 590. P. pr.
573. *L. Viscaria*, petalis subintegris. Fl. Sv. 409. Fl. N. 652. P. pr.
574. *L. alpina*, petalis bifidis, floribus tetragynis. Fl. Sv. 410. Fl. N. 679. Fl. D. 65. P. alp.

575. *L.*

*) Flores albi.

110 DECANDRIA. PENTAGYNIA. *Cerastium*.

575. *L. dioica*, floribus dioicis. Fl. Sv. 411. Fl. N. 123.
P. umbr. hum.
 α . *purpurea*,
 β . *alba* Fl. D. 792.
 γ . *hermaphrodita*.
576. *L. apetala*, calyce inflato, corolla calyce brevior, caule subunifloro hermaphrodito. Fl. Sv. 412. Fl. L. t. 12. f. 1. Fl. N. 680. Fl. D. 806. P. alp.

CERASTIUM.

577. *C. vulgatum*, foliis ovatis, petalis calyci aequalibus, caulibus diffusis. Fl. Sv. 415. Fl. N. 392. Vaill. Par. t. 30. f. 3. A. c. r.
578. *C. viscosum*, erectum villoso-viscosum. Fl. Sv. 414. Fl. N. 536. Vaill. Par. t. 30. f. 1. A. c.
579. *C. semidecandrum*, florum staminibus quinque sterilibus, petalis emarginatis. Fl. Sv. 416. Fl. N. 671. Vaill. Par. t. 30. f. 2. A. ficc.
580. *C. alpinum*, foliis ovato-lanceolatis caule diviso, capsulis oblongis. Fl. Sv. 418. Fl. N. 438. Obs. Bot. Fasc. III. p. 32. P. alp.
 α . *hirsutum*. Fl. D. 6. Fl. Scot. t. 10.
 β . *glabrum*. Fl. D. 979.
581. *C. aquaticum*, foliis cordatis sessilibus, floribus solitariis, fructibus pendulis. Fl. Sv. 413. Fl. N. 1036. P. hum.
- † 582. *C. latifolium*, foliis ovatis subtomentosis, ramis unifloris, capsulis globosis. S. N. 436. Fl. Isl. 7. P. alp.

SPERGULA.

583. *S. arvensis*, foliis verticillatis, floribus decandris, Fl. Sv. 419. Fl. N. 20. A. c. a.
584. *S. nodosa*, foliis oppositis subulatis laevibus, caulibus simplicibus. Fl. Sv. 420. Fl. D. 96. A. hum. inund.

585. *S.*

585. *S. saginoides*, foliis oppositis linearibus laevibus, pedunculis solitariis longissimis, floribus pentandris, caule repente. N. Act. Holm. 1789. p. 44. t. 1. f. 2. S. N. 437. P. alp.
586. *S. subulata*, foliis oppositis subsecundis lineari-subulatis carinatis ciliatis, floribus decandris. N. Act. Holm. 1789. p. 45. t. 3. f. 3. *Spergula saginoides* Pr. Fl. Scand. 523. Obs. Bot. Fasc. I. p. 19, II. p. 18, III. p. 32. *Spergula laricina* Fl. D. 858. A. pasc. ficc.

XI. DODECANDRIA.

1. MONOGYNIA.

ASARUM. *Cal.* 3-fid. *Cor.* 0. *Capsf.* 6-locul.

Rad. repens, geniculata. *Caul.* scaposa. *Fol.* simplex, radicalis, binata. *Flor.* terminalis, solitaria. *Fulcr.* 0.

LYTHRUM. *Cal.* 12-fid., *Cor.* 6-pet. calyci inserta. *Capsf.* 2-loc. polysperma.

Rad. fibrosa. *Caul.* herbacea, angulata. *Fol.* simplex, sessilis. *Flor.* terminalis, spicata. *Fulcr.* bracteae.

2. DIGYNIA.

AGRIMONIA. *Cal.* 5-dent. altero obvallatus. *Pet.* 5. *Sem.* 2.

3. TRIGYNIA.

RESEDA. *Cal.* 1 phyll. partit. *Pet.* laciniata. *Nectar.* glandula plana. *Capsf.* hians, 1-locul.

Rad. fusiformis. *Caul.* herbacea. *Fol.* simplex, undulata, sessilis. *Flor.* spicata. *Fulcr.* bracteae.

EUPHORBIA. *Cor.* 4-5-pet. calyci 1-phyll. ventricoso infidens. *Pet.* peltata. *Capsf.* 3-cocca. *Stam.* successive prorumpentia.

Rad. ramosa. *Caul.* herbacea, lactescens. *Fol.* simplex, lactescens. *Flor.* umbellata. *Fulcr.* involucria.

4. DODECAGYNIA.

SEMPERVIVUM. *Cal.* 12-part. *Cor.* 12-pet. *Capsf.* 12. Numerum quandoque auget.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex, crassa, succulenta, imbricata. *Flor.* cymosa. *Fulcr.* 0.

XI. DODECANDRIA.

I. MONOGYNIA.

ASARUM.

587. *A. europaeum*, foliis reniformibus obtusis binis.
Fl. Sv. 421. Fl. D. 633. P. nem.

LYTHRUM.

588. *L. Salicaria*, foliis oppositis cordato-lanceolatis,
floribus spicatis. Fl. Sv. 422. Fl. N. 539. Fl. D. 671.
P. l. lac, pr. hum.
 β . *trifolia*, foliis ternatis.
 γ . *lapponica*, floribus in alis solitariis.

2. DIGYNIA.

AGRIMONIA.

589. *A. Eupatoria*, foliis caulinis pinnatis, impari-pe-
tiolato, fructibus hispidis. Fl. Sv. 423. Fl. N. 229.
Fl. D. 588. P. pr. v. agr.

3. TRIGYNIA.

RESEDA.

590. *R. Luteola*, foliis lanceolatis integris basi utrinque
unidentatis, calycibus quadrifidis. Fl. Sv. 424.
Weig. Fl. 304. Fl. D. 864. B r. f. a.

EUPHORBIA.

591. *E. Peplus*, umbella trifida: dichotoma, involu-
cellis ovatis foliis integerrimis obovatis petiolatis.
Fl. Sv. 426. Fl. N. 738. A. r. c.
592. *E. exigua*, umbella trifida: dichotoma, involu-
cellis lanceolatis, foliis lineari-lanceolatis acutis w).
E. exigua acuta Sp. Pl. 654. α . Fl. D. 592. A. pr. agr.
593. *E.*

w) Corolla in hac tetrapetala, in *retusa* pentapetala, Cava-
nilles, qui species distinguit. *Usteri* Ann. 2. p. 64.

114 DODECANDRIA. DODECAGYNIA. Sempervivum.

593. *E. Helioscopia*, umbella quinquefida: trifida dichotoma, involucellis obovatis, foliis cuneiformibus ferratis. Fl. Sv. 425. Fl. N. 189. Fl. D. 725. A. c. r.
594. *E. Esula*, umbella multifida: bifida, involucellis subcordatis, petalis subbicornibus, ramis sterilibus, foliis uniformibus. S. N. 453. Ehrh. Beytr. 2. p. 147. x). P.
595. *E. palustris*, umbella multifida: subtrifida: bifida, involucellis ovatis, foliis lanceolatis, ramis sterilibus. Fl. Sv. 427. Fl. D. 866. P. pr. pal.
- † 596. *E. amygdaloides*, umbella multifida: dichotoma, involucellis perfoliatis orbiculatis, foliis obtusis. S. N. 454. y). P.

4 D O D E C A G Y N I A.

SEMPERVIVUM.

597. *S. testorum*, foliis ciliatis, propaginibus patentibus. Fl. Sv. 428. Fl. N. 1116. Fl. D. 601. A. test.

x) *Cel. Afzelius.*

y) *Cel. Kölpin.*

XII. ICOSANDRIA.

I. MONOGYNIA.

PRUNUS. *Cal.* 5-fid. inferus. *Cor.* 5-pet. *Drupae* nux
futuris prominulis.
Rad. lignosa. *Caul.* arborea. *Fol.* simplex, al-
terna. *Flor.* varia. *Fulcr.* o.

2. DIGYNIA.

CRATAEGUS. *Cal.* 5-fid. *Cor.* 5-pet. *Bacca* infera, di-
sperma.
Rad. lignosa. *Caul.* arborea. *Fol.* simplex, al-
terna, sublobata. *Flor.* corymbosa. *Fulcr.*
stipulae deciduae.

3. TRIGYNIA.

SORBUS. *Cal.* 5-fid. *Cor.* 5-pet. *Bacca* infera, tri-
sperma.
Rad. lignosa. *Caul.* arborea. *Fol.* subpinnata,
alterna. *Flor.* corymbosa. *Fulcr.* o.

4. PENTAGYNIA.

MESPILUS. *Cal.* 5-fid. *Cor.* 5-pet. *Bacca* infera,
5-sperma.
Rad. lignosa. *Caul.* fruticosa. *Fol.* simplex, al-
terna. *Flor.* terminalis ramulorum. *Fulcr.*
stipulae deciduae.

PYRUS. *Cal.* 5-fid. *Cor.* 5-pet. *Pomum* inferum, 5-loc.
polyperm.
Rad. lignosa. *Caul.* arborea. *Fol.* simplex, sparsa.
Flor. subumbellata. *Fulcr.* o.

SPIRAEA. *Cal.* 5-fid. *Cor.* 5-pet. *Capf.* plures polyspermae.

Rad. varia. *Caul.* herbacea vel fruticosa. *Fol.* alterna, simplex vel pinnata. *Flor.* corymbosa. *Fulcr.* stipulae.

5. POLYGYNIA.

ROSA. *Cal.* urceolatus, 5-fid. carnosus, collo coarctatus. *Cor.* 5-pet. *Sem.* plurima, hispida, calycis interiori lateri affixa.

Rad. lignosa. *Caul.* fruticosa, aculeata. *Fol.* pinnata, alterna. *Flor.* subumbellata. *Fulcr.* stipulae.

RUBUS. *Cal.* 5-fid. *Cor.* 5-pet. *Bacca* composita acinis monospermis.

Rad. repens. *Caul.* herbacea inermis vel fruticosa aculeata. *Fol.* alterna, simplex, ternata, quinata. *Flor.* varia. *Fulcr.* o. l. stipulae.

FRAGARIA. *Cal.* 10-fid. *Cor.* 5-pet. *Recept.* seminum ovatum, baccatum, deciduum.

Rad. fibrosa. *Caul.* repens, radicans. *Fol.* ternata. *Flor.* corymbosa. *Fulcr.* bracteae.

POTENTILLA. *Cal.* 10-fid. *Cor.* 5-pet. *Sem.* subrotunda, nuda, receptaculo parvo exsucco affixa.

Rad. varia. *Caul.* varia. *Fol.* ternata, quinata, pinnata. *Flor.* varia. *Fulcr.* stipulae.

TORMENTILLA. *Cal.* 8-fid. *Cor.* 4-pet. *Sem.* et *Recept.* Potentillae.

Rad. crassa, nodosa. *Caul.* herbacea. *Fol.* ternata. *Flor.* solitaria, axillaris et terminalis. *Fulcr.* stipulae.

GEUM. *Cal.* 10-fid. *Cor.* 5-pet. *Sem.* arista geniculata. *Rad.* fibrosa. *Caul.* herbacea, subangulata. *Fol.* lyrata. *Flor.* terminalis. *Fulcr.* stipulae.

DRYAS.

DRYAS. *Cal.* 8 fidus. *Cor.* 8-pet. *Sem.* caudata, pilosa.
Rad. subfusiformis. *Caul.* scaposa. *Fol.* simplex,
 radicalis. *Flor.* solitaria, terminalis. *Fulcr.* o.

COMARUM. *Cal.* 10 fid. *Pet.* 5. calyce minora. *Recept.*
 seminum ovatum, spongiosum, persistens.
Rad. fibrosa. *Caul.* adscendens, compressa.
Fol. pinnata, ternata. *Flor.* subpaniculata.
Fulcr. stipulae.

XII. ICOSANDRIA.

1. MONOGYNIA.

PRUNUS.

598. *P. Padus*, floribus racemosis, racemis pendulis, foliis deciduis subrugosis basi biglandulosis. H. Kew. II. p. 162. Fl. Sv. 431. Fl. N. 332. Fl. D. 205. Arb. nem.
599. *P. Cerasus*, umbellis subpedunculatis, foliis ovato-lanceolatis glabris conduplicatis. S. N. 463. Sp. Pl. 679. *η. austera*. Arb. sylv.
600. *P. avium*, umbellis sessilibus, foliis ovato-lanceolatis subtus pubescentibus conduplicatis. Fl. Sv. 429. Arb. sylv.
601. *P. domestica*, pedunculis subsolitariis, foliis lanceolato-ovatis convolutis, ramis muticis. Fl. Sv. 430. Arb.
602. *P. insititia*, pedunculis geminis, foliis ovatis subvillosis convolutis, ramis spinescentibus. S. N. 463. Arb. sylv.
603. *P. spinosa*, pedunculis solitariis, foliis lanceolatis glabris, ramis spinosis. Fl. Sv. 432. Fl. N. 417. Fl. D. 926. F. pasc. pr.

2. DIGYNIA.

CRATAEGUS.

604. *C. Aria*, foliis ovatis incisiss serratis subtus tomentosis. Fl. Sv. 433. Fl. N. 211. Fl. D. 302. Arb.
α. ulmifolia, foliis non sinuatis. *Pyrus Aria* Ehrh. Beytr. 4. p. 20. 2). A. Westrog. p. 161.
β. sinuata, Fl. L. 199. *Pyrus intermedia* Ehrh. Beytr. 4. p. 20.

605. *C.*

*) Haec certe omnia non pertinent Synonyma ibidem allata.

605. *C. torminalis*, foliis cordatis septangulis, lobis infimis divaricatis. S. N. 464. Weig. Fl. 312. Fl. D. 798. Arb.
606. *C. Oxyacantha*, foliis subtrifidis, sinibus acutis. Fl. Sv. 434. Fl. N. 620. Fl. D. 634. F. sylv.
607. *C. monogyna*, foliis subtrifidis, sinibus divaricatis. Jacqu. Fl. Austr. III. t. 292. a). F. sylv.

3. T R I G Y N I A.

SORBUS.

608. *S. aucuparia*, foliis pinnatis utrinque glabris. Fl. Sv. 435. Fl. N. 99. Arb.
609. *S. hybrida*, foliis semipinnatis subtus tomentosis. S. N. 465. Linn. Pl. rar. Fasc. I. p. 11. t. 6. Fl. N. 428. *Crataegus Aria* Fl. Sv. 433. γ. Arb.

4. P E N T A G Y N I A.

MESPILUS.

610. *M. Cotoneaster*, inermis, foliis ovatis integerrimis subtus tomentosis. Fl. Sv. 438. Fl. N. 470. Fl. D. 112. F. sylv.

PYRUS.

611. *P. communis*, foliis ferratis laevibus, floribus corymbosis. Fl. Sv. 436. Arb. sylv.
612. *P. malus*, foliis ferratis, umbellis sessilibus. Fl. Sv. 437. Fl. N. 447. Arb. sylv.

SPIRAEA.

613. *S. salicifolia*, foliis lanceolatis obtusis ferratis nudis, floribus duplicato-racemosis. Fl. N. 471. A. Holm. 1769. p. 247. F.

H 4

614. *S.*

- a) Folia in hac aequae ferrata ut in *Oxyacantha*, et in hac non magis obtusa quam in *monogyna*. Unitas pistilli, facies, magnitudo inferior et foliorum sinus melius distinguunt.

614. *S. Filipendula*, foliis pinnatis: foliolis uniformibus ferratis, caule herbaceo, floribus corymbosis. Fl. Sv. 439. Fl. N. 405. Fl. D. 635. P. pr. ficc.
615. *S. Ulmaria*, foliis pinnatis: impari majore lobato, floribus cymosis. Fl. Sv. 440. Fl. N. 215. Fl. D. 547. P. pr. hum.

5. P O L Y G Y N I A.

ROSA.

* *germinibus subglobosis.*

616. *R. Eglanteria*, germinibus globosis pedunculisque glabris, caule aculeis sparsis rectis, petiolis scabris, foliolis acutis. Fl. Sv. 443. b). F.
617. *R. arvensis*, germinibus globosis pedunculisque glabris, caule petiolisque aculeatis, floribus cymosis. M. Pl. 2. p. 245. Fl. N. 678. Fl. D. 398. F. f.
618. *R. cinnamomea*, germinibus globosis pedunculis petiolisque inermibus, caule aculeis geminis axillaribus, foliolis obovatis basi integris, calycibus integris spathulatis. *R. majalis* Obs. Bot. Fasc. III. p. 33. *R. cinnamomea* S. N. 473. F.
- † 619. *R. fluvialis*, germinibus globosis, pedunculis petiolisque inermibus, caule aculeis sparsis, foliolis ovatis acutis, calycibus linearibus incis. *R. fluvialis* Fl. D. 868 c). F. lit. lac.
620. *R. spinosissima*, germinibus globosis glabris, pedunculis hispidis, caule petiolisque aculeatissimis. Fl. Sv. 442. F. pr. v.
621. *R. villosa*, germinibus globosis pedunculisque hispidis, caule aculeis sparsis, petiolis aculeatis, foliis tomentosis. S. N. 474. Fl. Sv. 1295. F. pr. f.

** *ger-*b) Vix Sueciae indigena et forte loco *R. rubiginosae* in Flora Suecica introducta.

*) Ex icone charactèrem concinnavi.

** *germinibus oblongis.*

622. *R. rubiginosa*, germinibus ovatis petiolisque aculeatis, caule aculeis recurvis, foliis subtus rubiginosis. S. N. 473. *R. juvifolia* Fl. D. 870. d). F. sylv.
623. *R. canina*, germinibus ovatis pedunculisque glabris, caule petiolisque aculeatis. Fl. Sv. 441. Fl. N. 319. Fl. D. 555. F. f. v. pasc.

RUBUS.

624. *R. idaeus*, foliis quinato-pinnatis ternatisque, caule aculeato, petiolis canaliculatis. Fl. Sv. 446. Fl. N. 374. Fl. D. 788. F. f. lap.
625. *R. caesius*, foliis ternatis subnudis: lateralibus bilobis, caule aculeato tereti. Fl. Sv. 445. Fl. N. 375. F. a. f.
626. *R. fruticosus*, foliis quinato-digitatis ternatisque, caule petiolisque aculeatis. Fl. Sv. 443. Fl. N. 376. F. sylv. f.
627. *R. saxatilis*, foliis ternatis nudis, flagellis reptantibus herbaceis. Fl. Sv. 447. Fl. N. 299. Fl. D. 134. P. pr. nem.
628. *R. arcticus*, foliis ternatis, caule inermi unifloro. Fl. Sv. 448. Fl. L. t. 5. f. 2. Fl. N. 379. Fl. D. 488. P. pasc. musc. boreal.
629. *R. Chamaemorus*, foliis simplicibus lobatis, caule inermi unifloro declini. Fl. Sv. 449. Fl. L. t. 5. f. 1. Fl. N. 6. Fl. D. 1. P. pal.

FRAGARIA.

630. *F. vesca*, flagellis reptantibus. Fl. Sv. 450. Fl. N. 148. P. pr. pasc. sylv.

H 5

PO-

d) Haec vera est *Neglahryr* Westrogothorum et Scanorum.

POTENTILLA.

631. *P. fruticosa*, foliis pinnatis, caule fruticoso. Fl. Sv. 451. Morif. 2. f. 2. t. 23. f. 5. S. ul.
632. *P. Anserina*, foliis pinnatis serratis, caule repente, pedunculis unifloris. Fl. Sv. 452. Fl. N. 38. Fl. D. 544. P. v. pasc. pr. arg.
633. *P. rupestris*, foliis pinnatis alternis: foliolis quinis ovatis crenatis, caule erecto. Fl. Sv. 453. Morif. 2. f. 2. t. 20. f. 1. P. pr. mont.
- † 634. *P. supina*, foliis pinnatis, caule dichotomo decumbente. S. N. 477. e). A.
635. *P. argentea*, foliis quinatis cuneiformibus incisissimis subtus tomentosis, caule erecto. Fl. Sv. 454. Fl. N. 144. Fl. D. 865. P. ficc.
- † 636. *P. opaca*, foliis radicalibus quinatis cuneiformibus serratis, caulibus suboppositis, ramis filiformibus decumbentibus. S. N. 478. f). P.
637. *P. verna*, foliis radicalibus quinatis acute serratis retusis, caulibus ternatis, caule declinato. Fl. Sv. 455. Fl. N. 51. P. ficc.
638. *P. aurea*, foliis radicalibus quinatis serratis acuminatis, caulibus ternatis, caule declinato. S. N. 478. Fl. N. 472. Fl. D. 114. g). P. alp.
639. *P. reptans*, foliis quinatis, caule repente, pedunculis unifloris. Fl. Sv. 456. Fl. N. 402. P. ficc.
640. *P. norvegica*, foliis ternatis, caule dichotomo, pedunculis axillaribus. Fl. Sv. 457. Obs. Bot. F. I. p. 20. Fl. N. 145. Fl. D. 171. B. c. alp.

641. *P.*

e) Cl. Hellwig.

f) Cel. Kölpin.

g) Si a *Verna* vere diversa aegre distinguitur.

641. *P. nivea*, foliis ternatis incisfis subtus tomentosis, caule ascendente. Fl. Sv. 458. Fl. N. 512. t. 3. f. 1. A. H. X. p. 451. t. 7. f. 22. Skr. Nat. H. Selfk. 2. p. 23. P. alp.

β. caule erecto, foliis utrinque hirsutis subtus subsericeis. *Vahl* l. cit. p. 24. h).

γ. floribus subseptenis glomeratis. A. H. X. p. 453.

- † 642. *P. retusa*, foliis ternatis hirtis apice retuso tridentato. Fl. D. 799. i). P.

TORMENTILLA.

643. *T. erecta*, caule erectiusculo, foliis sessilibus. Fl. Sv. 459. Fl. N. 66. Fl. D. 589. P. pr. hum. ul.

GEUM.

644. *G. urbanum*, floribus erectis, fructibus globosis villosis, aristis uncinatis nudis, foliis lyratis. Fl. Sv. 460. Fl. N. 63. Fl. D. 672. P. pr. nem.

645. *G. rivale*, floribus nutantibus, fructu oblongo, aristis plumosis tortis. Fl. Sv. 461. Fl. N. 64. Fl. D. 722. P. pr.

β. flore multiplicato.

DRYAS.

646. *D. octopetala*, foliis simplicibus. Fl. Sv. 462. Fl. N. 106. Fl. D. 31. P. alp.

COMARUM.

647. *C. palustre*. Fl. Sv. 463. Fl. N. 65. Fl. D. 636. P. pal. ul.

h) Etiam in Lapponia Svecica.

i) Proxima *Clusianae*, at flores flavi, et caules erecti. *P. tridentata* H. Kew. II. p. 216. t. 9. eadem videtur, cultura procerior facta.

XIII. POLYANDRIA.

I. MONOGYNIA.

ACTAEA. *Cal.* 4-phyll. *Cor.* 4-pet. *Bacca* 1-locul. polysperma.

Rad. nodosa. *Caul.* herbacea. *Fol.* alterna, composita, tripinnata. *Flor.* racemosa. *Fulcr.* stipulae.

CHELIDONIUM. *Cal.* 2-phyll. caducus. *Cor.* 4-pet. *Siliqua* linearis.

Rad. fusiformi-ramosa. *Caul.* herbacea. *Fol.* pinnatifido-sinuata. *Flor.* varia, pedunculata. *Fulcr.* bracteae.

PAPAVER. *Cal.* 2-phyll. caducus. *Cor.* 4-pet. *Caps.* 1-locul. dissepimentis dimidiatis, sub stigmate persistente poris dehiscens.

Rad. fusiformis. *Caul.* herbacea, stricta. *Fol.* pinnatifido-sinuata. *Flor.* terminalis, solitaria. *Fulcr.* o.

NYMPHAEA. *Cal.* 4-5-phyll. *Cor.* polypetala. *Bacca* multilocularis, truncata.

Rad. horizontalis, crassa. *Caul.* demersa, scaposa. *Fol.* simplex, natans. *Flor.* solitaria, natans. *Fulcr.* o.

TILIA. *Cal.* 5 part. *Cor.* 5-pet. *Bacca* sicca, globosa. 5-locul. 5-valv. basi dehiscens.

Rad. lignosa. *Caul.* arborea. *Fol.* simplex, alterna. *Flor.* pedunculata e medio bracteae. *Fulcr.* bracteae.

CISTUS. *Cal.* 5-phyll. foliolis 2 minoribus. *Cor.* 5-pet. *Fruct.* capsula.

Rad. fibrosa. *Caul.* suffruticulosa. *Fol.* simplex, varia. *Flor.* varia. *Fulcr.* bracteae, stipulae nonnullis.

2. TRIGYNIA.

DELPHINIUM. *Cal.* o. *Pet.* 5. *Nectar.* 2-fid. postice cornutum. *Siliquae* 3-1.

Rad. ramosa. *Caul.* herbacea. *Fol.* composita, multifida. *Flor.* paniculata. *Fulcr.* bracteae.

ACONITUM. *Cal.* o. *Pet.* 5. supremo fornicato. *Nect.* 2. pedunculata, recurva. *Siliquae* 3-5.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex, palmato-multifida. *Flor.* racemosa. *Fulcr.* o.

3. TETRAGYNIA.

Myriophyllum verticillatum v. *Monoec. Polyandr.*

4. PENTAGYNIA.

AQUILEGIA. *Cal.* o. *Cor.* 5 pet. *Nect.* 5. cornuta inter petala. *Caps.* 5.

Rad. ramosa. *Caul.* herbacea. *Fol.* triternata. *Flor.* subcorymbosa. *Fulcr.* bracteae.

5. POLYGYNIA.

ANEMONE. *Cal.* o (prima specie excepta). *Pet.* 6-9. *Sem.* plurima.

Rad. fibrosa. *Caul.* herbacea, varia. *Fol.* lobata l. multifida. *Flor.* solitaria. *Fulcr.* o. l. involucrum.

THALICTRUM. *Cal.* o. *Pet.* 4-5. *Sem.* ecaudata.

Rad. fusiformis l. repens. *Caul.* herbacea. *Fol.* pinnata, trilobata. *Flor.* racemoso-paniculata. *Fulcr.* stipulae.

ADONIS. *Cal.* 5-phyll. *Pet.* quinque plura absque nectario. *Sem.* nuda.

Rad. fibrosa. *Caul.* herbacea, foliosa, simplex. *Fol.* multifido-linearis. *Flor.* terminalis, solitaria. *Fulcr.* o.

RANUNCULUS. *Cal.* 5-phyll. *Pet.* 5. intra ungues poro vel squama nectarifera. *Sem.* nuda.

Rad. fibrosa l. tuberosa. l. repens. *Caul.* herbacea. *Flor.* varia, pedunculata. *Fulcr.* o.

TROLLIUS. *Cal.* o. *Cor.* polypetala. *Nectar.* linearia, plana. *Capsf.* plures polyspermae.

Rad. fibrosa. *Caul.* herbacea. *Fol.* 5-partita: radicalis petiolata, caulina sessilis. *Flor.* solitaria. *Fulcr.* o.

HELLEBORUS. *Cal.* o. *Cor.* 5-6-pet. *Nectar.* tubulosa, 2-labiata. *Capsf.* polyspermae.

Rad. fibrosa. *Caul.* scaposa. *Fol.* composita, *Flor.* terminalis. *Fulcr.* bractea remota.

CALTHA. *Cal.* o. *Peti.* 4. *Nectar.* o. *Capsf.* plures, polyspermae.

Rad. subhorizontalis, fibrosa. *Caul.* herbacea, obliqua. *Fol.* simplex, semiorbicularis. *Flor.* solitaria, pedunculata. *Fulcr.* o.

XIII. POLYANDRIA.

I. MONOGYNIA.

ACTAEA.

648. *A. spicata*, racemo ovato, fructibus baccatis. Fl. Sv. 464. Fl. N. 27. Fl. D. 496. P. sylv.

CHELIDONIUM.

649. *C. majus*, pedunculis umbellatis. Fl. Sv. 465. Fl. N. 706. Fl. D. 542. P. f. r.

β. *tenuifolium*, l. foliis quernis. C. Bauh. Fl. D. 676.

- † 650. *C. Glaucium*, pedunculis unifloris, foliis amplexicaulibus sinuatis, caule glabro. S. N. 489. Fl. D. 585. A. B.

PAPAVER.

651. *P. Argemone*, capsulis clavatis hispidis, caule folioso multifloro. Fl. Sv. 466. Fl. D. 867. A. agr.

- † 652. *P. nudicaule*, capsulis hispidis, scapo unifloro nudo hispido, foliis simplicibus pinnato-sinuatis. S. N. 490. Fl. N. 578. Fl. D. 41. *P. radicum* A. H. X. p. 455. t. 8. f. 24. P. alp.

653. *P. Rhoëas*, capsulis glabris globosis, caule piloso multifloro, foliis pinnatifidis incis. Fl. Sv. 468. Fl. N. 455. A. c.

654. *P. dubium*, capsulis glabris oblongis, caule multifloro setis adpressis, foliis pinnatifidis incis. Fl. Sv. 467. Fl. N. 1105. Fl. D. 902. A. c.

NYMPHAEA.

655. *N. lutea*, foliis cordatis integerrimis, calyce pentaphyllo petalis majore. Fl. Sv. 469. Fl. N. 440. Fl. D. 603. P. aqu.

656. *N. alba*, foliis cordatis integerrimis, calyce quadrifido. Fl. Sv. 470. Fl. N. 441. Fl. D. 602. P. aqu.

TILIA.

TILIA.

657. *T. europaea*, floribus nectario destitutis. Fl. Sv. 471.
 Fl. N. 442. Fl. D. 553. Arb. nem.
α. grandifolia.
β. parvifolia.
γ. corallina, viminibus rubris.

CISTUS.

658. *C. Oelandicus*, suffruticosus procumbens exstipulatus, foliis oppositis oblongis utrinque glabris, petiolis ciliatis, petalis emarginatis. Fl. Sv. 473.
 Jacqu. Fl. Austr. IV. t. 399. S. ficc.
 659. *C. Fumana*, suffruticosus procumbens exstipulatus, foliis alternis linearibus margine scabris, pedunculis unifloris. Fl. Sv. 474. Jacqu. Fl. Austr. III. t. 252. S. ficc.
 660. *C. Helianthemum*, suffruticosus procumbens, stipulis lanceolatis, foliis oblongis revolutis subpilis. Fl. D. 101. S. ficc.

2. TRIGYNIA.

DELPHINIUM.

661. *D. Consolida*, nectariis monophyllis, caule subdiviso. Fl. Sv. 475. Fl. D. 683. A. c.

ACONITUM.

662. *A. Lycostomum*, foliis palmatis multifidis villosis. Fl. Sv. 476. Fl. N. 14.
α. coeruleum Fl. D. 123. Aët. Phys. Lund. T. I. P. I. p. 44.
 663. *A. Napellus*, foliorum laciniis linearibus superne latioribus linea exaratis, pedunculis nudis. Fl. Sv. 477. P. nem.

3. P E N T A G Y N I A.

AQUILEGIA.

664. *A. vulgaris*, nectariis incurvis. Fl. Sv. 478. Fl. D. 695. P. mont.

4. P O L Y G Y N I A.

ANEMONE.

665. *A. Hepatica*, foliis trilobis integerrimis. Fl. Sv. 480. Fl. N. 168. Fl. D. 610. P. nem.
 †666. *A. patens*, pedunculo involucrato, foliis digitatis multifidis. S. N. 509. k).
 667. *A. vernalis*, pedunculo involucrato, foliis pinnatis, flore erecto. Fl. Sv. 483. Fl. N. 734. Fl. D. 29. P. sylv. acer.
 668. *A. Pulsatilla*, pedunculo involucrato, foliis bipinnatis, petalis rectis. Fl. Sv. 481. Fl. N. 617. Fl. D. 153. P. ficc.
 669. *A. pratensis*, pedunculo involucrato, foliis bipinnatis, petalis apice reflexis. Fl. Sv. 482. Fl. D. 611. P. a.
 670. *A. sylvestris*, pedunculo nudo, seminibus subrotundis hirsutis muticis. Fl. Sv. 484. Clus. Hist. I. p. 244. P. nem.
 671. *A. nemorosa*, seminibus acutis, foliolis incisfis, caule unifloro. Fl. Sv. 485. Fl. N. 166. Fl. D. 549. P. pr. nem.
 672. *A. ranunculoides*, seminibus acutis, foliolis incisfis, petalis subrotundis, caule subbifloro. Fl. Sv. 486. Fl. N. 1117. Fl. D. 140. P. nem.

THALICTRUM.

673. *T. alpinum*, caule simplicissimo subnudo, racemo simplici terminali. Fl. Sv. 491. Fl. N. 41. Fl. D. 11. P. alp.

674. *T.*

k) *Cel. Kölpin.*

674. *T. minus*, floribus paniculatis cernuis. Fl. Sv. 489. Fl. N. 785. Fl. D. 732. l). P. pr.
675. *T. flavum*, caule folioso sulcato, panicula multiplici erecta. Fl. Sv. 488. Fl. N. 40. Fl. D. 939. P. pr.
676. *T. simplex*, caule folioso simplicissimo angulato. Fl. Sv. 490. Fl. N. 784. Fl. D. 244. P. pr. sylv. ul.
677. *T. aquilegifolium*, fructibus pendulis triangularibus rectis, caule tereti. Fl. Sv. 487. Obs. Bot. Fasc. VI. p. 30. P. pr. nem.

ADONIS.

- † 678. *A. aestivalis*, floribus pentapetalis, fructibus ovatis. S. N. 514. m). A. c.
679. *A. vernalis*, flore dodecapetalo, fructu ovato. Fl. Sv. 492. P. apr.

RANUNCULUS.

680. *R. Flammula*, foliis ovato-lanceolatis petiolatis, caule declinato. Fl. Sv. 494. Fl. N. 467. Fl. D. 575. P. ul.
681. *R. reptans*, foliis linearibus caule repente. Fl. Sv. 495. Fl. L. t. 3. f. 5. Fl. N. 434. Fl. D. 108. n). P. lit. lac.
682. *R. Lingua*, foliis lancolatis, caule erecto. Fl. Sv. 493. Fl. D. 755. A. aqu.
683. *R. Ficaria*, foliis cordatis angulatis petiolatis, caule unifloro polypetalo. Fl. Sv. 496. Fl. N. 433. Fl. D. 499. P. c. f. nem.

684. *R.*

l) Folia sexpartita nondum vidi, sed triloba sunt.

m) Cel. Kölpin.

n) Male omnino a nonnullis pro varietate praecedentis habetur.

684. *R. auricomus*, foliis radicalibus reniformibus crenatis incis: caulinis digitatis linearibus, caule multifloro. Fl. Sv. 498. Fl. N. 493. Fl. D. 665, P. pr. pasc. hum.
685. *R. sceleratus*, foliis inferioribus palmatis: summis digitatis, fructibus oblongis. Fl. Sv. 499. Fl. N. 84. Fl. D. 571. A. aqu.
- †686. *R. platanifolius*, foliis palmatis laevibus incis, caule erecto, bracteis linearibus. S. N. 516. Fl. N. 792. Fl. D. 111. P. alp.
687. *R. illyricus*, foliis ternatis integerrimis lanceolatis. Fl. Sv. 500. Jacqu. Fl. Austr. III. t. 222. P.
688. *R. glacialis*, calycibus hirsutis, caule bifloro, foliis multifidis. Fl. Sv. 501. Fl. L. t. 3. f. 1. Fl. N. 160. Fl. D. 19. P. alp.
689. *R. nivalis*, calyce hirsuto, caule unifloro, foliis radicalibus palmatis: caulinis multipartitis sessilibus. Fl. Sv. 502. Fl. L. t. 3. f. 2. Fl. N. 627. *R. lapponicus* Fl. D. 144. P. alp.
 β. *pygmaeus*. Fl. L. 232. γ. t. 3. f. 3. N. A. Holm. 1789. p. 47.
690. *R. lapponicus*, foliis tripartitis lobatis obtusis, caule subnudo unifloro. Fl. Sv. 503. Fl. L. t. 3. f. 4. Fl. N. 542. P. alp.
- †691. *R. hyperboreus*, foliis profunde trilobis, lobis oblongis divaricatis, caule filiformi repente, A. H. X. p. 458. t. 4. f. 16. *R. Ammanni* Fl. N. 826. Fl. D. 331. Fl. Sib. T. IV. p. 204. t. 83. o). alp.

*) Varietatem prioris statuunt ill. Linné et rev. Gunnérus; distinctissimus caule repente.

132 POLYANDRIA. POLYGYNIA. *Ranunculus*.

692. *R. bulbosus*, calycibus retroflexis, pedunculis fulcatis, caule erecto multifloro, foliis compositis. Fl. Sv. 504. Fl. N. 490. Fl. D. 551. P. pr.
693. *R. Philonotis*, calycibus demum reflexis setosis, pedunculis fulcatis, foliis tripartitis inciso-lobatis hirsutis, fructu subglobofo. *R. Philonotis*. Ehrh. Beytr. 2. p. 145. Obs. Bot. Fasc. VI. p. 31. Bauh. Hist. Pl. III. p. 417. *R. hirsutus*. H. Kew. II. p. 268. p). P. pr. hum. a.
694. *R. repens*, calycibus patulis, pedunculis fulcatis, stolonibus repentibus, foliis compositis. Fl. Sv. 505. Obs. Bot. Fasc. I. p. 21. Fl. N. 825. Fl. D. 795. P. c. r.
695. *R. polyanthemus*, calycibus patulis, pedunculis fulcatis; caule erecto; foliis multipartitis. Fl. Sv. 506. Fl. N. 468. P. pr.
696. *R. acris*, calycibus patulis, pedunculis teretibus, foliis tripartito-multifidis: summis linearibus. Fl. Sv. 507. Fl. N. 160. P. pr. pasc.
- † 697. *R. lanuginosus*, calycibus patulis, pedunculis teretibus, caule petiolisque hirsutis, foliis trifidis lobatis crenatis holosericeis. S. N. 517. Weig. Fl. 365. Fl. D. 397. P. sylv.
698. *R. arvensis*, feminibus aculeatis, foliis superioribus decompositis linearibus. Fl. Sv. 508. Fl. D. 219. A. agr.
699. *R. hederaceus*, foliis subrotundis trilobis integerrimis, caule repente. S. N. 518. Obs. Bot. F. I. p. 21. Fl. D. 321. A. qu.

700. *R.*

p) Quomodo cum *R. scelerato* a Crantz, Hagen etc. comparari potuit non perspicio. *R. hirsutum* H. Kew. eundem esse credo, licet annum pronunciet.

POLYANDRIA. POLYGYNIA. Trollius. 133.

700. *R. aquatilis*, foliis submersis capillaceis, emarjis peltatis. Fl. Sv. 509. Fl. N. 646. P. aqu.

α. *anomalus*, foliis capillaceis nullis q).

β. *heterophyllus*, foliis difformibus.

γ. *millefolius*, foliis omnibus capillaceis, laciniis divergentibus.

δ. *peucedanifolius*, foliis omnibus capillaceis, laciniis parallelis subbifidis. Fl. D. 376.

TROLLIUS.

701. *T. europaeus*, corollis conniventibus, nectariis longitudine staminum. Fl. Sv. 510. Fl. N. 355. Fl. D. 133. P. pr.

HELLEBORUS.

702. *H. trifolius*, scapo unifloro, foliis ternatis. Sp. Pl. 784. Am. Acad. V. 2. p. 356. t. 4. f. 18. *Anemone groenlandica*. Fl. N. 1066. Fl. D. 566. P. frig.

CALTHA.

703. *C. palustris*. Fl. Sv. 511. Fl. N. 216. Fl. D. 668. P. ul.

q) In fossis fere exsiccatis haud raro occurrit.

XIV. DIDYNAMIA.

1. GYMNOSPERMA.

AJUGA. *Cal.* 5-fid. *Cor.* labium superius minimum emarginatum. *Stam.* labio superiore longiora. *Rad.* fibrosa. *Caul.* simplex, herbacea. *Fol.* simplex, opposita. *Flor.* verticillata. *Fulcr.* bracteae coloratae,

TEUCRIUM. *Cal.* 5-fid. *Cor.* labium superius (nullum) ultra basin bipartitum, divaricatum.

Rad. fibrosa. *Caul.* herbacea. *Fol.* simplex, opposita. *Flor.* varia. *Fulcr.* o. l. bracteae.

NEPETA. *Cal.* 5-fid. *Cor.* labium inferius crenulatum. *Faux* margine reflexo. *Stam.* approximata. *Rad.* ramosa. *Caul.* herbacea. *Fol.* simplex, opposita. *Flor.* verticillato-spicata. *Fulcr.* bracteae minutae.

MENTHA. *Cal.* 5-fid. *Cor.* 4-fid. subaequalis, lacinia latiore emarginata. *Stam.* erecta, distantia.

Rad. repens. *Caul.* herbacea. *Fol.* simplex, opposita. *Flor.* spicata l. verticillata. *Fulcr.* o.

GLECOMA. *Cal.* 5-fid. *Antherarum* singulum par in formam crucis connivens.

Rad. repens. *Caul.* herbacea, decumbens, radicans. *Fol.* simplex, opposita. *Flor.* subverticillata. *Fulcr.* o.

LAMIUM. *Cal.* 5-dent. *Cor.* labium superius fornicatum: inferius 2-lobum. *Faux* utrinque margine denticulo instructa.

Rad. fibrosa. *Caul.* herbacea. *Fol.* simplex. *Flor.* verticillata. *Fulcr.* bracteae subulatae.

GALOPSIS. *Cal.* 5-fid. *Cor.* labium superius subcrenatum, fornicatum: inferius supra 2-dentatum. *Rad.* ramosa. *Caul.* herbacea. *Fol.* simplex, opposita. *Flor.* verticillata. *Fulcr.* bracteae subulatae.

RETO.

- BETONICA.** *Cal.* 5-fid. aristatus. *Cor.* labium superius adscendens, planiusculum: tubus cylindricus. *Rad.* ramosa. *Caul.* herbacea. *Fol.* opposita, simplex, rugosa. *Flor.* spicata. *Fulcr.* bracteae.
- STACHYS.** *Cal.* 5-fid. *Cor.* labium superius fornicatum: inferius lateribus reflexum, lacinia media majore emarginata. *Stam.* deflorata versus latera reflexa. *Rad.* repens l. ramosa. *Caul.* herbacea. *Fol.* simplex, opposita. *Flor.* verticillata. *Fulcr.* bracteae.
- BALLOTA.** *Cal.* hypocrateriformis, 5-dent. 10-striat. *Cor.* labium superius crenatum, concavum. *Rad.* ramosa. *Caul.* herbacea. *Fol.* simplex, opposita. *Flor.* verticillata, secunda. *Fulcr.* bracteae setaceae.
- MARRUBIUM.** *Cal.* hypocrateriformis, rigidus, 10-striat. *Cor.* labium superius 2-fid., lineare, erectum. *Rad.* ramosa. *Caul.* herbacea. *Fol.* simplex, opposita, rugosa. *Flor.* globofo-verticillata. *Fulcr.* bracteae subulatae.
- LEONURUS.** *Cal.* 5-fid. *Cor.* labium superius integrum. *Antherae* punctis nitidis aspersae. *Rad.* ramosa. *Caul.* herbacea. *Fol.* simplex, lobata, opposita. *Flor.* verticillata. *Fulcr.* bracteae subulatae.
- CLINOPODIUM.** *Cal.* 2-labiat. *Involucrum* multifetum verticillo subiectum. *Rad.* ramosa. *Caul.* herbacea. *Fol.* simplex, opposita. *Flor.* verticillato-capitata. *Fulcr.* bracteae.
- ORIGANUM.** *Cal.* 2-labiat. *Strobilus* 4-gonus spicatus, calyces colligens. *Rad.* ramosa. *Caul.* herbacea. *Fol.* simplex, opposita. *Flor.* corymbosa, fastigiata. *Fulcr.* bracteae coloratae.

THYMUS. *Cal.* 2-labiat. fauce villis clausa.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex, opposita. *Flor.* capitata l. verticillata. *Fulcr.* bracteae minutae.

DRACOCEPHALUM. *Cal.* 2 labiat. *Cor.* faux inflata. *Lab. super.* concavum.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex, opposita. *Flor.* spicata. *Fulcr.* bracteae.

SCUTELLARIA. *Cal.* ore integro, post florescentiam clauso operculato.

Rad. repens. *Caul.* herbacea, ascendens. *Fol.* simplex, opposita. *Flor.* axillaris, solitaria. *Fulcr.* bracteae.

PRUNELLA. *Cal.* 2-lab. *Filamenta* 2-furca, altero apice antherifera. *Stigma* 2-fid.

Rad. repens. *Caul.* herbacea. *Fol.* simplex, opposita. *Flor.* capitato-oblonga. *Fulcr.* bracteae amplexicaules.

Verbena officinalis vid. Diandr. Monog.

2. ANGIOSPERMA.

BARTSIA. *Cal.* 2-lob. colorat. *Cor.* lab. super. longius. *Capsf.* 2-locul.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex, opposita. *Flor.* terminalis, subspicata. *Fulcr.* bracteae color.

RHINANTHUS. *Cal.* 4-fid. ventricos. *Capsf.* 2-locul. obtusa, compressa.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex, opposita. *Flor.* spicata. *Fulcr.* bracteae.

EUPHRASIA. *Cal.* 4-fid. cylindr. *Capsf.* 2-locul. ovato-oblonga. *Antherae* inferiores altero lobo basi spinosae.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex, opposita. *Flor.* subspicata, secunda. *Fulcr.* bracteae.

MELAM-

MELAMPYRUM. *Cal.* 4-fid. *Cor.* lab. super. compressum margine replicato. *Capsf.* 2-loc. obliqua. *Sem.* 2. gibba.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex, opposita. *Flor.* subspicata. *Fulcr.* bracteae coloratae.

LATHRAEA. *Cal.* 4-fid. *Glandula* depressa ad basin futurae germinis. *Capsf.* 1-locul.

Rad. dentato-squamosa. *Caul.* similis, subterrestris. *Fol.* crassa, squamosa. *Flor.* spicata, secunda. *Fulcr.* bracteae unilaterales, imbricatae, coloratae.

PEDICULARIS. *Cal.* 5-fid. *Cor.* lab. super. compressum. *Capsf.* 2-locul. mucronata, obliqua. *Sem.* tunicata.

Rad. subramosa. *Caul.* herbacea. *Fol.* pinnatifido-laciniata. *Flor.* terminalis, spicata. *Fulcr.* bracteae.

ANTIRRHINUM. *Cal.* 5-phyll. *Cor.* basis deorsum prominens nectarifera. *Capsf.* 2-locul.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex, alternata. *Flor.* varia. *Fulcr.* 0.

SCROPHULARIA. *Cal.* 5-fid. *Cor.* subglobosa, resupinata. *Capsf.* 2-locul.

Rad. ramosa l. tuberosa. *Caul.* herbacea, tetragona. *Fol.* simplex, opposita. *Flor.* axillaris, tristis. *Fulcr.* bracteae minutae.

DIGITALIS. *Cal.* 5-part. *Cor.* campanulata 5-fid. ventricosa. *Capsf.* 2-locul.

Rad. subfusiformis. *Caul.* herbacea. *Fol.* simplex, alternata. *Flor.* racemosa, secunda. *Fulcr.* bracteae.

LINNEA. *Cal.* 2-plex. *floris* 5-part. *fructus* 2-phyll. *Cor.* campanulata. *Bacca* sicca 3-locul.

Rad. repens. *Caul.* decumbens. *Fol.* simplex, opposita. *Flor.* geminata, cernua. *Fulcr.* bracteae minutae.

LIMOSELLA. *Cal.* 5-fid. *Cor.* 5-fid. aequalis. *Stam.*
per paria approximata. *Capsf.* 1-loc. 2-valv.
Rad. fibrosa. *Caul.* radicans. *Fol.* simplex, ra-
dicalis. *Flor.* radicalis, pedunculata. *Fulcr.* o.

OROBANCHE. *Cal.* 2-fid. *Cor.* ringens. *Capsf.* 1-locul.
2-valv. *Glandula* sub basi germinis.
Rad. bulbosa, squamata, parasitica. *Caul.* fim-
plex, squamosa. *Fol.* o. *Flor.* racemosa.
Fulcr. squamae.

XIV. DIDYNAMIA.

I. GYMNOSPERMIA.

AJUGA.

704. *A. pyramidalis*, tetragono-pyramidalis villosa, foliis radicalibus maximis. Fl. Sv. 512. Fl. N. 343. Fl. D. 185. B. pasc.
 †705. *A. alpina*, caule simplici, foliis caulinis radicalia aequantibus. S. N. 525. Fl. N. 1099. P. alp.
 †706. *A. reptans*, glabra, stolonibus repentibus. S. N. 525. Weig. Fl. 371. Fl. D. 925. P. nem.

TEUCRIUM.

- †707. *T. Chamaepithys*, foliis trifidis linearibus integerrimis, floribus sessilibus lateralibus solitariis, caule diffuso. S. N. 525. Fl. D. 733. A.
 708. *T. Scordium*, foliis oblongis sessilibus dentato-ferratis, floribus geminis axillaribus pedunculatis, caule diffuso. Fl. Sv. 513. Fl. D. 593. P. pr. pal.
 †709. *T. Scorodonia*, foliis cordatis ferratis petiolatis, racemis lateralibus secundis, caule erecto. S. N. 527. Fl. D. 485. P. sylv.

NEPETA.

710. *N. Cataria*, floribus spicatis: verticillis subpedicellatis, foliis pedicellatis cordatis dentato-ferratis. Fl. Sv. 514. Fl. D. 580. P. r. f.

MENTHA.

711. *M. sylvestris*, spicis oblongis, foliis oblongis tomentosis ferratis sessilibus, staminibus corolla longioribus. S. N. 532. Fl. D. 484. P. hum.
 712. *M. viridis*, spicis oblongis, foliis lanceolatis nudis ferratis sessilibus, staminibus corolla longioribus. S. N. 532. r). P.

713. *M.*

r) *Cel. Afzelius*.

140 DIDYNAMIA. GYMNOSPERMIA. Glecoma.

713. *M. hirsuta*, floribus capitatis, foliis ovatis ferratis subsessilibus pubescentibus, staminibus corolla longioribus. S. N. 532. Fl. N. 1110. Raji Syn. 3. t. 10. f. 1. 5). P. hum.
714. *M. aquatica*, floribus capitatis, foliis ovatis ferratis petiolatis, staminibus corolla longioribus. Fl. Sv. 517. Fl. N. 223. Fl. D. 673. P. riv.
- †715. *M. sativa*, floribus verticillatis, foliis ovatis acutiusculis ferratis, staminibus corolla longioribus. S. N. 512. Fl. D. 794. P.
716. *M. gentilis*, floribus verticillatis, foliis ovatis acutis ferratis, staminibus corolla brevioribus. S. N. 533. Weig. Suppl. 879. Fl. D. 673. ? P.
717. *M. arvensis*, floribus verticillatis, foliis ovatis acutis ferratis, staminibus corollam aequantibus. Fl. Sv. 516. Fl. N. 114. Fl. D. 512. P. c. hum.

GLECOMA.

718. *G. hederacea*, foliis reniformibus crenatis. Fl. Sv. 518. Fl. D. 789. 1). P. pr. c. f.

LAMIUM.

719. *L. album*, foliis cordatis acuminatis ferratis petiolatis, verticillis vigintifloris. Fl. Sv. 520. Fl. N. 937. Fl. D. 594. P. c. r. n.
720. *L. purpureum*, foliis cordatis obtusis petiolatis. Fl. Sv. 521. Fl. N. 466. Fl. D. 523. A. c.
721. *L. amplexicaule*, foliis floralibus sessilibus amplexicaulibus obtusis. Fl. Sv. 522. Fl. N. 938. Fl. D. 752. A. c. r.

GALEO-

5) An satis a sequente distincta?

7) Primo vere plerumque masculinis partibus destituitur.

GALEOPSIS.

722. *G. Ladanum*, internodiis caulinis aequalibus, verticillis omnibus remotis, calycibus inermibus. Fl. Sv. 524. Fl. N. 354. A. c. r.
 α. *angustifolia*. St. Helv. 266.
 β. *latifolia*. St. Helv. 267.
723. *G. Tetrahit*, internodiis caulinis superne incrassatis, verticillis summis subcontiguis, calycibus subpungentibus. Fl. Sv. 523. Fl. N. 75. A. c.
 α. *vulgaris*, flore rubro aut albo. St. Helv. 268.
 β. *cannabina*, floribus magnis flavis maculatis. St. Helv. 269. Fl. D. 929.
724. *G. Galeobdolon*, verticillis sexfloris, involucro tetraphyllo. Fl. Sv. 525. P. nem.

BETONICA.

725. *B. officinalis*, spica interrupta, corollarum labii lacinia intermedia emarginata. Fl. Sv. 515. Weig. Fl. 383. Fl. D. 726. P. pr.

STACHYS.

726. *S. sylvatica*, verticillis sexfloris, foliis cordatis petiolatis. Fl. Sv. 526. Fl. N. 70. P. sylv. n.
727. *S. palustris*, verticillis subsexfloris, foliis linearilanceolatis semiamplexicaulibus sessilibus. Fl. Sv. 528. Fl. N. 25. P. agr.
728. *S. alpina*, verticillis multifloris, foliorum serraturis apice cartilagineis, corollis labio plano. Fl. Sv. 527. Ros. Obs. p. 64. P. mont.
- † 729. *S. recta*, verticillis subspicatis, foliis cordato-ellipticis crenatis scabris, caulibus adscendentibus. S. N. 536. Jacqu. Fl. Austr. IV. t. 359. u). A. c.
 730. *S.*

*) *Cel. Afzelius*.

142 DIDYNAMIA. GYMNOSPERMIA. Ballota.

730. *S. arvensis*, verticillis sexfloris, foliis obtusis nudiusculis, corollis longitudine calycis, caule debili. S. N. 536. Fl. N. 1009. Fl. D. 587. Weig. Suppl. 878. *Glecoma arvensis*. Fl. Sv. 519. A. c.

BALLOTA.

731. *B. nigra*, foliis cordatis indivisis ferratis, calycibus acuminatis. Fl. Sv. 529. P. f.
732. *B. alba*, foliis cordatis indivisis ferratis, calycibus subtruncatis. Fl. Sv. 530. P. f.

MARRUBIUM.

733. *M. vulgare*, dentibus calycinis setaceis uncinatis. Fl. Sv. 531. Fl. N. 418. P. r.

LEONURUS.

734. *L. Cardiaca*, foliis caulinis lanceolatis trilobis. Fl. Sv. 532. Fl. N. 719. Fl. D. 727. P. r.

CLINOPODIUM.

735. *C. vulgare*, capitulis subrotundis hispidis, bracteis setaceis. Fl. Sv. 533. Fl. N. 527. Fl. D. 930. P. pr. mont.

ORIGANUM.

736. *O. vulgare*, spicis subrotundis paniculatis conglomeratis, bracteis calyce longioribus ovatis. Fl. Sv. 534. Fl. N. 112. Fl. D. 638. P. nem. mont.

THYMUS.

737. *T. Serpyllum*, floribus capitatis, caulibus repentibus, foliis planis obtusis basi ciliatis. Fl. Sv. 535. Fl. N. 233. P. sicc.
 α . *rubrum*.
 β . *album*.
 γ . *lanuginosum*.

738. *T.*

DIDYNAMIA. GYMNOSPERMIA. Dracocephalum. 143

738. *T. Acinos*, floribus verticillatis, pedunculis unifloris, caulibus erectis subramosis, foliis acutis serratis. Fl. Sv. 536. Fl. N. 158. Fl. D. 814. A. pr. sicc.

DRACOCEPHALUM.

739. *D. Ruyschiana*, floribus spicatis, foliis bracteisque lanceolatis indivisis muticis. Fl. Sv. 537. Fl. N. 519. Fl. D. 121. P. pr.
740. *D. thymiflorum*, floribus verticillatis, bracteis oblongis integerrimis, corollis vix calyce majoribus. S. N. 544. Fl. Sib. III. p. 233. t. 50. v). A.

SCUTELLARIA.

741. *S. galericulata*, foliis cordato-lanceolatis crenatis, floribus axillaribus. Fl. Sv. 538. Fl. N. 128. Fl. D. 637. P. l. lac. fluv. riv.
742. *S. hastifolia*, foliis integerrimis: inferioribus hastatis: superioribus sagittatis. Fl. Sv. 539. P. l. lac.

PRUNELLA.

743. *P. vulgaris*, foliis omnibus ovato-oblongis serratis petiolatis. Fl. Sv. 540. Fl. N. 156. Fl. D. 910. P. pr.
- β. *grandiflora*. Poll. 2. p. 174. w). Jacqu. Fl. Austr. IV. t. 377.

2. ANGIOSPERMIA.

BARTSIA.

744. *B. alpina*, foliis oppositis cordatis obtuse serratis. Fl. Sv. 541. N. Aët. Holm. 1789. p. 47. Fl. N. 100. Fl. D. 43. P. riv. alp.

RHI-

v) Cel. Afzelius.

w) Parum adeo differunt, ut saepe *grandifloram* et *parvifloram* ex una eademque radice vidi, nec credo *laciniatam* essentialiter differre.

RHINANTHUS.

745. *R. Christa Galli*, corollis labio superiore compresso brevior. Fl. Sv. 542. Fl. N. 305. A. pr. agr.
 α. *foemina*, calycibus glabris, auriculis galeae bifidis, nectario acuminato. *Scop. Rh. major*. Ehrh. Beytr. 6. p. 144. Fl. D. 981. ficc.
 β. *mas*, calycibus villosis, auriculis galeae truncatis, nectario rotundato. *Scop. Rh. minor*. Ehrh. l. c. hum.

EUPHRASIA.

746. *E. officinalis*, foliis ovatis lineatis argute dentatis. Fl. Sv. 543. Fl. N. 214. A. pr. pasc.
 747. *E. Odontites*, foliis linearibus, omnibus serratis. Fl. Sv. 544. Fl. D. 625. A. agr. hum. v.

MELAMPYRUM.

748. *M. cristatum*, spicis quadrangularibus, bracteis cordatis compactis denticulatis imbricatis. Fl. Sv. 545. A. pr. ster.
 749. *M. arvense*, spicis conicis laxis bracteis dentato-setaceis coloratis. Fl. Sv. 546. Fl. D. 911. A. agr.
 750. *M. nemorosum*, floribus secundis, bracteis dentatis: summis coloratis sterilibus, calycibus lanatis. Fl. Sv. 547. Fl. N. 718. Fl. D. 305. A. pr. nem.
 751. *M. pratense*, floribus secundis lateralibus, foliorum conjugationibus remotis, corollis clausis. Fl. Sv. 548. Fl. N. 146. A. pr.
 752. *M. sylvaticum*, floribus secundis lateralibus, foliorum conjugationibus remotis, corollis hiantibus. Fl. Sv. 549. Fl. N. 147. Fl. D. 145. A. sylv.

LATHRAEA.

753. *L. Squamaria*, caule simplicissimo, corollis pendulis, labio inferiore trifido. Fl. Sv. 550. Fl. D. 136. P. sylv.

PEDICULARIS.

754. *P. palustris*, caule ramoso, calycibus cristatis caloso-punctatis, corollis labio obliquis. Fl. Sv. 551. it. Westr. 40. Fl. N. 87. Fl. D. 225. ? x). A. ul.
755. *P. sylvatica*, caule ramoso, calycibus oblongis angulatis laevibus, corollis labio cordato. Fl. Sv. 552. it. Westr. 40. Fl. N. 703. A. pr. ul. sylv.
756. *P. Sceptum Carolinum*, caule simplici, floribus terno-verticillatis, corollis clausis, calycibus cristatis, capsulis regularibus. Fl. Sv. 553. Fl. N. 33. Fl. D. 26. P. sylv. ul.
757. *P. flammea*, caule simplici, foliis pinnatis recto imbricatis. Fl. Sv. 556. Fl. L. t. 4. f. 2. P. alp.
758. *P. hirsuta*, caule simplici, foliis dentato-pinnatis linearibus, calycibus hirsutis. Fl. Sv. 555. Fl. L. t. 4. f. 3. Fl. N. 469. y). P. alp.
759. *P. lapponica*, caule simplici, foliis pinnatifidis ferratis, calycibus bifidis obtusis. Fl. Sv. 554. Fl. L. t. 4. f. 1. Fl. N. 246. Fl. D. 2. P. alp.
- †760. *P. Groenlandica*, caule simplici, foliis pinnatis ferratis, calycibus oblongis glabris, corollis galea subulata arcuata longiore. *Ped. incarnata*. Pr. Fl. Scand. 680. Obs. Bot. Fasc. IV. 87. t. 1. z). P.
- †761. *P.*

x) Icon Fl. D. nomen *sylvaticae* habet, *palustrem* vero refert sine crista, quae variatio non multum rara.

y) Ad hanc, dubitantes licet, fig. 30. Fl. Dan. refert ill. a Linné, eandem figuram ad *flammeam* citat ill. Håller cum admonitione. Neutram referre certum est, an nova? *Ped. hirsuta*. All. Fl. Ped t. 2. f. 1. diversa omnino ut et *rosea* Jacq. Coeterum pace celeb. Haenke dictum sit Oederi figuram *P. flammeam* e Lapponia nec forma nec coloribus referre.

z) Communicavit Cel. Vahl vere distinctam speciem.

146 DIDYNAMIA. ANGIOSPERMIA. Antirrhinum.

- †761. *P. foliosa*, caule simplici, spica foliosa, corollis galea obtusissima integra, calycibus quinque dentatis. S. N. 553. Jacqu. Fl. Austr. II. t. 139. St. Helv. t. 11. f. 2. Fl. N. 1111. P. alp.

ANTIRRHINUM.

- †762. *A. Elatine*, foliis hastatis alternis, caulibus procumbentibus. S. N. 554. Fl. D. 426. A.
- †763. *A. spurium*, foliis ovatis alternis, caulibus procumbentibus. S. N. 554. Fl. D. 913. A.
764. *A. minus*, foliis plerisque alternis lanceolatis obtusis, caule ramosissimo diffuso. Fl. Sv. 558. Fl. D. 502. A. ficc.
765. *A. Linaria*, foliis lanceolato-linearibus confertis, caule erecto, spicis terminalibus sessilibus, floribus imbricatis. Fl. Sv. 557. Fl. N. 273. Fl. D. 982. P. f. coem. v. agr.
- β. *multiplicata*. Söderb. Diff. Florae Ryb.
766. *A. Orontium*, corollis ecaudatis, floribus subspicatis, calycibus corolla longioribus. Fl. Sv. 559. Fl. D. 941. A. c.

SCROPHULARIA.

767. *S. nodosa*, foliis cordatis trinervatis, caule obtusangulo. Fl. Sv. 560. Fl. N. 732. P. hum. pingu.
- †768. *S. aquatica*, foliis cordatis obtusis petiolatis decurrentibus, caule membranis angulato, racemis terminalibus. Fl. N. 560. Weig. Suppl. 880. Fl. D. 507. P. hum. umbr.
769. *S. vernalis*, foliis cordatis: caulinis ternis a) pedunculis axillaribus solitariis bifidis. S. N. 560. Fl. D. 411. P. c.

DIGI-

- a) Saepius tantum opposita.

DIDYNAMIA. ANGIOSPERMIA. Digitalis. 147,

DIGITALIS.

770. *D. purpurea*, calycinis foliolis ovatis acutis, corollis obtusis: labio superiore integro. S. N. 562. Fn. Sv. 557. Fl. D. 74. B. r.

LINNEA.

771. *L. borealis*. Fl. Sv. 562. c. icone. Fl. L. t. 12. f. 4. Fl. N. 67. Fl. D. 3. A&. Nidr. 4. t. 4. f. 6. P. sylv. acer.

LIMOSELLA.

772. *L. aquatica*, foliis lanceolatis. Fl. Sv. 563 Fl. N. 469. Fl. D. 69. A. qu. 1. lac.

OROBANCHE.

773. *O. major*, caule simplicissimo pubescente, staminibus subexsertis. Fl. Sv. 561. P. par. rad.

XV. TETRADYNAMIA.

I. SILICULOSA.

MYAGRUM. *Silic.* stylo conico terminata: loculo submonospermo.

Rad. simplex. *Caul.* herbacea. *Fol.* simplex, alterna. *Flor.* terminalis, subpaniculata. *Fulcr.* o.

SUBULARIA. *Silic.* integra, ovata; valvis concavis dissepimento contrariis. *Stylus* filicula brevior.

Rad. fibrosa. *Caul.* herbacea. *Fol.* simplex, graminea. *Flor.* alterna. *Fulcr.* o.

DRABA. *Silic.* integra ovali-oblonga: valvis planiusculis dissepimento parallelis.

Rad. simplex. *Caul.* herbacea. *Fol.* simplex, alterna. *Flor.* subracemosa. *Fulcr.* o.

LEPIDIUM. *Silic.* emarginata, cordata, polysperma: valvis carinatis contrariis.

Rad. simplex. *Caul.* herbacea. *Fol.* alterna, varia. *Flor.* subracemosa. *Fulcr.* o.

THLASPI. *Silic.* emarginata, obcordata, polysperma: valvis navicularibus marginato-carinatis.

Rad. simplex. *Caul.* herbacea. *Fol.* simplex. *Flor.* racemosa. *Fulcr.* o.

COCHLEARIA. *Silic.* emarginata, turgida, scabra: valvis gibbis obtusis.

Rad. varia. *Caul.* herbacea. *Fol.* simplex, carnosa. *Flor.* corymbosa. *Fulcr.* o.

IBERIS. *Cor.* irregularis: petalis 2 exterioribus majoribus. *Silic.* polysperma, emarginata.

Rad. simplex. *Caul.* scaposa. *Flor.* racemosa. *Fol.* radicalis. *Fulcr.* o.

AEYSSUM. *Filamenta* quaedam introsum denticulo notata. *Silic.* emarginata.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex.

Flor. corymbosa. *Fulcr.* o.

LUNARIA. *Silic.* integra, elliptica, compresso-plana, pedicellata, valvis dissepimento parallelis planis. *Cal.* foliolis faccatis.

Rad. subliguosa, ramosa. *Caul.* herbacea. *Fol.* simplex, cordata. *Flor.* subcorymbosa. *Fulcr.* o.

2. SILIQUOSA.

DENTARIA. *Sil.* elastice diffiliens valvulis revolutis. *Stigma* emarginatum. *Cal.* longitudinaliter connivens.

Rad. nodoso-squamosa. *Caul.* herbacea, axillis bulbiferis. *Fol.* alterna, varia. *Flor.* corymbosa. *Fulcr.* o.

CARDAMINE. *Sil.* elastice diffiliens, valvulis revolutis. *Stigm.* integrum. *Cal.* subhians.

Rad. fibrosa. *Caul.* herbacea. *Fol.* varia. *Flor.* subracemosa. *Fulcr.* o. l. stipulae.

SISYMBRIUM. *Sil.* dehiscens valvulis rectiusculis. *Cal.* patens. *Cor.* patens.

Rad. simplex l. fibrosa. *Caul.* herbacea. *Fol.* varia. *Flor.* racemosa l. axillaris. *Fulcr.* o.

ERYSIMUM. *Sil.* columnalis exacte tetraëdra. *Cal.* clausus.

Rad. simplex l. ramosa. *Caul.* herbacea. *Fol.* simplex. *Flor.* racemosa. *Fulcr.* o.

CHEIRANTHUS. *Germen* utrinque denticulo glandulato. *Cal.* clausus foliolis 2 basi gibbis. *Sem.* plana.

Rad. ramosa. *Caul.* herbacea. *Fol.* alterna, simplex. *Flor.* racemoso-spicata. *Fulcr.* o.

HESPERIS. *Petalis* oblique flexa. *Gland.* inter flamina breviora. *Sil.* stricta. *Stigma* basi bifurca apice connivente. *Cal* clausus.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex. *Flor.* subcorymb. *Fulcr.* o.

ARABIS. *Gland. nectar.* 4. singula intra calycis folium squamae instar reflexae. *Stigm.* simplex. *Sil.* linearis, torosa.

Rad. subramosa. *Caul.* herbacea. *Fol.* simplex, alterna. *Flor.* subpaniculata. *Fulcr.* o.

TURRITIS. *Sil.* longissima, stricta, angulata. *Cal.* connivens, erectus. *Cor.* erecta.

Rad. simplex. *Caul.* herbacea, simplicissima. *Fol.* simplex, alterna. *Flor.* racemoso-spicata. *Fulcr.* o.

BRASSICA. *Cal.* erectus, connivens. *Gland.* inter flamina breviora et pistillum interque longiora et calycem. *Sil.* septo valvis longiore.

Rad. fusiformis. *Caul.* herbacea. *Fol.* simplex. *Flor.* subracemosa. *Fulcr.* o.

SINAPIS. *Cal.* patens. *Cor.* ungues recti. *Gland.* Brassicae. *Sil.* septo longius producto cornuta.

Rad. subramosa. *Caul.* herbacea. *Fol.* simplex, sinuata, hispida. *Flor.* racemosa. *Fulcr.* o.

RAPHANUS. *Cal.* clausus. *Sil.* torosa, subarticulata, teres. *Gland.* Brassicae.

Rad. simplex. *Caul.* herbacea. *Fol.* simplex, sinuata. *Flor.* subracemosa. *Fulcr.* o.

BUNIAS. *Sil.* decidua, tetraëdra, angulis inaequalibus acuminatis muricatis,

Rad. fusiformis. *Caul.* herbacea. *Fol.* varia. *Flor.* varia. *Fulcr.* o.

ISATIS.

ISATIS. *Sil.* lanceolata. 1-loc. 1-sperma, decidua, 2-valvis: valvulis navicularibus.

Rad. fusiformis. *Caul.* herbacea, fastigiata.

Fol. simplex. *Flor.* racemosa. *Fulcr.* o.

CRAMBE. *Filamenta* 4 longiora apice bifurca: altero antherifero. *Bacca* ficca, globosa, decidua.

Rad. fusiformis. *Caul.* herbacea. *Fol.* simplex, sinuata. *Flor.* racemosa, axillaris. *Fulcr.* o.

XV. TETRADYNAMIA.

I. SILICULOSA.

MYAGRUM.

774. *M. sativum*, filiculis obovatis pedunculatis polyspermis. Fl. Sv. 564. Fl. N. 403. A. agr.
 775. *M. paniculatum*, filiculis lentiformibus orbiculatis punctato-rugosis. Fl. Sv. 565. Fl. N. 980. Fl. D. 204. A. agr.

SUBULARIA.

776. *S. aquatica*. Fl. Sv. 566. Fl. N. 786. Fl. D. 35. P. agr.

DRABA.

777. *D. alpina*, scapo nudo simplici, foliis lanceolatis integerrimis. Fl. Sv. 570. Fl. N. 435. Fl. D. 56. P. alp.
 778. *D. verna*, scapis nudis, foliis subserratis. Fl. Sv. 567. Fl. N. 2. Fl. D. 983. A. ficc.
 779. *D. muralis*, caule ramofo, foliis cordatis amplexicaulibus dentatis, racemo subpubescente, pedicellis filicula nuda vix longioribus. *Dr. muralis a.* Fl. Sv. 569. Ehrh. Beytr. 7. p. 153. B. ficc.
 780. *D. nemoralis*, caule ramofo, foliis ovatis sessilibus dentatis, racemo glabro, pedicellis filicula pubescente longioribus. *Dr. nemoralis* Fl. Sv. 569. β. Ehrh. Beytr. 7. p. 154. B. nem.
 781. *D. hirta*, scapo unifolio, foliis subhirsutis, filiculis obliquis pedicellatis. S. N. 585. Fl. Sv. 557. Fl. N. 697. Fl. D. 142. P. alp.
 β. *Dr. norwegica*, foliis radicalibus subcuneiformibus acutis dentatis congestis: caulinis paucis lanceolatis petiolatis integris. Fl. N. 846. Fl. D. 143. b). P. alp.

782. *D.*

b) An potius nova species ex sola icone et Fl. N. difficile dictu.

TETRADYNAMIA. SILICULOSA. *Lepidium*. 453

782. *D. incana*, foliis caulinis numerosis incanis, filiculis contortis pubescentibus. Fl. Sv. 568. Fl. N. 3. Fl. D. 130. *Dr. Confusa*. Ehrh. Beytr. 7. p. 155. c). B. ficc.
783. *D. contorta*, foliis caulinis numerosis hirsutis, filiculis contortis glabris. Ehrh. Beytr. 7. p. 155. d). B. ficc.

LEPIDIUM.

784. *L. alpinum*, foliis pinnatis integerrimis, scapo subradicato, filiculis lanceolatis mucronatis. S. N. 586. Fl. N. 1069. Fl. D. 569. e). P. alp.
785. *L. petraeum*, foliis pinnatis integerrimis, petalis emarginatis calyce minoribus. Fl. Sv. 573. Jacqu. Fl. Austr. II. t. 131. B. apr.
786. *L. latifolium*, foliis ovato-lanceolatis, integris serratis. Fl. Sv. 571. Fl. D. 557. P. a.
787. *L. ruderale*, floribus diandris apetalis, foliis radicalibus dentato-pinnatis: ramiferis linearibus integerrimis. Fl. Sv. 572. Fl. N. 549. Fl. D. 184. A. r.

THLASPI.

788. *T. arvense*, filiculis orbiculatis, foliis oblongis dentatis glabris. Fl. Sv. 574. Fl. N. 306. Fl. D. 793. A. agr. c.
789. *T. campestre*, filiculis subrotundis, foliis sagittatis dentatis incanis. Fl. Sv. 575. B. agr.
790. *T. Bursa Pastoris*, filiculis obcordatis, foliis radicalibus pinnatifidis. Fl. Sv. 576. Fl. N. 308. A. c. r.
 a. sinuata. Fl. D. 729.
 β. integrifolia.

K 5

COCH-

- e) Pili caulis, foliorum et fructus echinati.
 d) Haud vidi, auctoritate Cl. Ehrhardi adpositi.
 e) Cum interrogatione adducitur haec figura in S. N.

154 TETRADYNAMIA. SILICULOSA. Cochlearia.

COCHLEARIA.

791. *C. officinalis*, foliis radicalibus cordato-subrotundis, caulinis oblongis subsinuatis. Fl. Sv. 577. Fl. N. 196. Fl. D. 135. B. l. m.
792. *C. Danica*, foliis hastato-angulatis, omnibus deltoidibus. Fl. Sv. 578. Fl. N. 179. Fl. D. 100. B. l. m.
- †793. *C. Anglica*, foliis omnibus ovato-lanceolatis. S. N. 588. Fl. D. 329. B. l. m.
- †794. *C. Groenlandica*, foliis reniformibus integris carnosiss. S. N. 588. Fl. N. 491. B. l. m.
795. *C. Coronopus*, foliis pinnatifidis, caule depresso. Fl. Sv. 579. Fl. N. 800. Fl. D. 202. A. v. f. a.
796. *C. Armoracia*, foliis radicalibus lanceolatis crenatis; caulinis incisis. Fl. Sv. 580. P. hum.

IBERIS.

797. *I. nudicaulis*, herbacea, foliis sinuatis, caule nudo simplici. Fl. Sv. 581. Fl. D. 323. B. a.
- β. *integrifolia*, foliis omnibus orbiculatis.

ALYSSUM.

798. *A. incanum*, caule erecto, foliis lanceolatis incanis integerrimis, floribus corymbosis, petalis bifidis. Fl. Sv. 582. B. a.
- †799. *A. calycinum*, caulibus herbaceis, staminibus omnibus dentatis, calycibus persistentibus. S. N. 590. Jacqu. Fl. Austr. IV. t. 338. f). A.

LUNARIA.

800. *L. rediviva*, siliculis lanceolatis, radice perenni. Obs. Bot. Fasc. I. p. 68. Pr. Fl. Scand. 715. Fl. N. 592. Clus. H. Pl. I. p. 297. P. nem.

801. *L.*

f) *Cel. Kölpin.*

TETRADYNAMIA. SILIQUOSA. Dentaria. 155

801. *L. annua*, filiculis ellipticis, radice bienni. Obs.
Bot. F. I. 69. Fl. Sv. 583. g). B.

2. S I L I Q U O S A.

DENTARIA.

802. *D. bulbifera*, foliis inferioribus pinnatis, summis simplicibus. Fl. Sv. 584. Fl. D. 361. P. sylv.

CARDAMINE.

803. *C. bellidifolia*, foliis simplicibus ovatis integerrimis: petiolis longis. Fl. Sv. 590. Fl. L. t. 9, f. 2. Fl. D. 20. P. ? alp.

804. *C. petraea*, foliis simplicibus oblongis dentatis. Fl. Sv. 591. Fl. N. 744. Fl. D. 386. P. alp.

a. hirsuta.

β. glabra.

805. *C. trifolia*, foliis ternatis obtusis, caule subnudo. Fl. Sv. 589. Clus. H. Pl. II. p. 127. Jacqu. Fl. Austr. I. t. 27. P. alp.

806. *C. Impatiens*, foliis pinnatis incisiss stipulatis, floribus apetalis. Fl. Sv. 588. A. nem. hum.

807. *C. parviflora*, foliis pinnatis exstipulatis, foliolis lanceolatis obtusis, floribus corollatis. Fl. Sv. p. 464. *Card. Impatiens* Fl. D. 735. h). A. nem. hum.

808. *C. hirsuta*, foliis pinnatis, floribus tetrandris. Fl. Sv. 587. Fl. N. 195. i). P. pr.

809. *C. pratensis*, foliis pinnatis, foliolis radicalibus subrotundis: caulinis lanceolatis. Fl. Sv. 585. Fl. N. 10. P. pasc. pr. hum.

810. *C.*

g) Prior Scanica, haec Westrogothica, monente ill. a Linné in litt. unde femina accepit ad Hortum Upsal.

h) Ob defectum stipularum huc retuli.

i) Stamina 4, 5, 6 numeravit Pollich, quid ergo hunc a varietate hirsuta *C. amarae* distinguit?

156 TETRADYNAMIA. SILIQUOSA. *Sisymbrium*.

810. *C. amara*, foliis pinnatis, axillis stoloniferis. Fl. Sv. 586. Fl. N. 43. k). P. riv. umbr.

α. *glabra*.

β. *hirsuta*.

SISYMBRIUM.

811. *S. Nasturtium*, filiquis declinatis, foliis pinnatis: foliolis subcordatis. Fl. Sv. 592. Fl. N. 893. Fl. D. 690. A. riv. font.

812. *S. sylvestre*, filiquis declinatis oblongo-ovatis, foliolis lanceolatis ferratis. Fl. Sv. 594. Fl. N. 891. P. hum.

β. *S. islandicum*, filiquis declinatis, foliis ly-rato-pinnatis, foliolis ovalibus dentatis. Fl. D. 409. Fl. N. 850.

813. *S. amphibium*, filiquis declinatis oblongo-ovatis, foliolis pinnatifidis ferratis. Fl. Sv. 593. Fl. N. 892. P. hum. aqu.

α. *palustre*, foliis pinnatifidis ferratis. Fl. D. 931.

β. *aquaticum*, foliis simplicibus dentatis ferratis. Fl. D. 984.

γ. *terrestre*, foliis inferioribus simplicibus, superioribus pinnatis.

814. *S. supinum*, filiquis axillaribus subsessilibus solitariis, foliis dentato-sinuatis. S. N. 595. Fn. Sv. 557. A. r.

815. *S. arenosum*, caule subfolioso ramoso, foliis lyratis rectangulo-dentatis hispidis: pilis ramosis. Fl. Sv. 597. *Arabis arenosa* Scop. Fl. Carn. ed. 2. 837. t. 40. l). A. a.

816. *S.*

k) Huc potius pertinet Fl. D. t. 148. figura licet minus bona. Maxime ludit hirsutiae, glabritie, magnitudine et proceritate, non semper stolonifera.

l) Variat caule simplici et ramoso, corollis albis et violaceis.

TETRADYNAMIA. SILIQUOSA. *Erysimum*. 157.

816. *S. Sophia*, petalis calyce minoribus, foliis decomposito-pinnatis. Fl. Sv. 595. Fl. N. 277. Fl. D. 528. A. c. r.
817. *S. Irio*, foliis runcinatis dentatis nudis, caule laevi, filiquis erectis, Fl. Sv. 596. Fl. N. 511. A. c.

ERYSIMUM.

818. *E. officinale*, filiquis spicae appressis, foliis runcinatis. Fl. Sv. 598. Fl. N. 411. Fl. D. 560. A. r. coem.
819. *E. Barbarea*, foliis lyratis: extimo subrotundo. Fl. Sv. 599. Fl. N. 717. Fl. D. 985. B. c. f. v.
820. *E. Alliaria*, foliis cordatis. Fl. Sv. 600. Fl. D. 935. B. P. n. f.
821. *E. Cheiranthoides*, foliis lanceolatis integerrimis, filiquis patulis. Fl. Sv. 601. Obs. Bot. Fasc. I. 64. Fl. N. 587. Fl. D. 731. A. agr.
822. *E. Hieracifolium*, foliis lanceolatis ferratis. Fl. Sv. 602. Obs. Bot. F. I. 62. Fl. N. 513. Fl. D. 923. B. c.

CHEIRANTHUS.

823. *C. Erysimoides*, foliis lanceolatis dentatis, caule erecto simplicissimo, filiquis tetragonis. Fl. Sv. 603. Fl. N. 585. Fl. D. 229. B. c.

HESPERIS.

824. *H. inodora*, caule simplici erecto, foliis subhastatis petalis obtusis. S. N. 599. Weig. Fl. 429. Fl. D. 924. m). B. nem.

ARABIS.

825. *A. alpina*, foliis amplexicaulibus dentatis. Fl. Sv. 604. Fl. N. 413. Fl. D. 602. P. alp.
826. *A.*

m) Fl. D. t. 921. videtur mihi *Hesp. Sibirica* Sp. Pl. 927. quae varietas est *H. matronalis*.

158 TETRADYNAMIA. SILIQUOSA. Turritis.

826. *A. Thaliana*, foliis petiolatis integerrimis. Fl. Sv. 605. Fl. N. 495. A. ficc.
 β. foliis radicalibus subdentatis. Obs. Bot. F.I. p. 22.

TURRITIS.

827. *T. glabra*, foliis radicalibus dentatis hispidis: caulinis integerrimis amplexicaulibus glabris. Fl. Sv. 606. Fl. N. 348. Fl. D. 809. B. ficc.
 828. *T. hirsuta*, foliis omnibus hispidis: caulinis amplexicaulibus. Fl. Sv. 607. Fl. N. 344. B. fylv.
 829. *T. alpina*, foliis radicalibus dentatis hispidis: caulinis semiamplexicaulibus. S. N. 600. B.

BRASSICA.

830. *B. campestris*, radice cauleque tenui, foliis caulinis uniformibus cordatis sessilibus. Fl. Sv. 608. Fl. N. 301. Fl. D. 550. A. agr.
 831. *B. Napus*, radice caulescente fusiformi. Fl. Sv. 609. B. agr.

SINAPIS.

832. *S. arvensis*, siliquis multangulis toroso-turgidis laevibus, rostro ancipiti longioribus. Fl. Sv. 610. Fl. N. 302. Fl. D. 753. A. agr.
 833. *S. nigra*, siliquis glabris racemo appressis. Fl. Sv. 611. A. agr. f.

RAPHANUS.

834. *R. Raphanistrum*, siliquis teretibus articulatis laevibus unilocularibus. Fl. Sv. 612. Fl. N. 415. *Sinapis arvensis*. Fl. D. 678. A. agr.

BUNIAS.

835. *B. Cakile*, filiculis ovatis laevibus ancipitibus. Fl. Sv. 613. Fl. N. 21. A. l. m. a.
 836. *B. orientalis*, filiculis ovatis gibbis verrucosis. S. N. 603. Gmel. Fl. Sib. 3. t. 57. P. r. f.

ISATIS.

TETRADYNAMIA. SILIQUOSA. Isatis. 159

ISATIS.

837. *I. tinctoria*, foliis radicalibus crenatis: caulinis sagittatis, filiculis oblongis. Fl. Sv. 614. B. l. m.

CRAMBE.

838. *C. maritima*, foliis cauleque glabris. Fl. Sv. 615. Fl. N. 981. Fl. D. 316. P. l. m.

XVI. MONADELPHIA.

I. DECANDRIA.

GERANIUM. 1-gyn. *Cal.* 5-fid. l. 5-phyll. *Stigm.* 5.
Fruct. rostratus, 5 coccus. *Capsf.* feminum
 monospermae, arista demum spirali.
Rad. subfusiformis. *Caul.* herbacea, geniculata.
Fol. lobata l. pinnata. *Flor.* longe peduncu-
 lata. *Fulcr.* bracteae, stipulae.

2. POLYANDRIA.

ALTHAEA. *Cal.* duplex: *exterior* 9-fidus. *Arilli* plu-
 rimi monospermi.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex. *Flor.*
 axillaris. *Fulcr.* stipulae.

MALVA. *Cal.* duplex: *exterior* 3-phyllus. *Arilli* plu-
 rimi monospermi.

Rad. subfusiformis. *Caul.* herbacea. *Fol.* simplex,
 plicata. *Flor.* axillaris. *Fulcr.* stipulae.

LAVATERA. *Cal.* duplex: *exterior* 3-fidus. *Arilli* plu-
 rimi monospermi.

Rad. ramosa. *Caul.* herbacea. *Fol.* lobata. *Flor.*
 terminalis, axillaris. *Fulcr.* stipulae.

XVI. MONADELPHIA.

I. D E C A N D R I A,

GERANIUM.

839. *G. cicutarium*, pedunculis multifloris, floribus pentandris, foliis pinnatis incisus obtusis, caule ramoso. Fl. Sv. 625. Fl. N. 628. Fl. D. 986. A. c. r.
840. *G. phaeum*, pendunculis solitariis oppositifoliis, calycibus subaristatis, caule erecto, petalis undulatis. S. N. 616. Obs. Bot. F. I. p. 23. Fl. D. 987. P. pr. fylv.
- †841. *G. fuscum*, pedunculis bifloris oppositifoliis geminis, caule patulo, petalis integerrimis. S. N. 616. *G. montanum*. Fl. Isl. p. 9. P. alp.
842. *G. sylvaticum*, pedunculis bifloris, foliis subpeltatis quinquelobis inciso-ferratis, caule erecto, petalis emarginatis. Fl. Sv. 617. Fl. N. 73. Fl. D. 124. P. fylv.
843. *G. palustre*, pedunculis bifloris longissimis declinatis, petalis integris, foliis quinquelobis incisus. S. N. 617. Obs. Bot. F. I. p. 23. Weig. Suppl. 884. Fl. D. 596. P. pr. hum. n.
844. *G. pratense*, pedunculis bifloris, foliis subpeltatis multipartitis rugosis acutis, petalis integris. Fl. Sv. 618. Fl. N. 631. P. pr. fylv.
845. *G. robertianum*, pedunculis bifloris, calycibus pilosis decemangulatis. Fl. Sv. 619. Fl. N. 37. Fl. D. 694. B. fice, f.
846. *G. lucidum*, pedunculis bifloris, calycibus pyramidalis angulatis elevato-rugosis, foliis quinquelobis rotundatis. Fl. Sv. 620. Fl. N. 488. Fl. D. 218. B. mont.

L

847. *G.*

162 MONADELPHIA. POLYANDRIA. *Althaea*.

847. *G. molle*, pedunculis bifloris foliisque floralibus alternis, petalis bifidis, calycibus muticis, caule erectiusculo. Fl. Sv. 624. Vaill. Par. t. 15. f. 3. Fl. D. 679. A. c. r. f.
848. *G. Columbinum*, pedunculis bifloris folio longioribus, foliis quinquepartito-multifidis, arillis glabris, calycibus aristatis, Fl. Sv. 623. Vaill. Par. t. 15. f. 4. Fl. D. 932. A. c. r.
849. *G. dissectum*, pedunculis bifloris, foliis quinquepartito-trifidis, petalis emarginatis longitudine calycis, arillis villosis. Fl. Sv. 622. Obs. Bot. F. I. p. 72. Vaill. Par. t. 15. f. 2. Fl. D. 936. A. c.
850. *G. rotundifolium*, pedunculis bifloris, petalis subintegris longitudine calycis, caule prostrato, foliis reniformibus incis. Fl. Sv. 621. Cavan. Diff. IV. p. 214. t. 93. f. 2. A. r. c.
851. *G. sanguineum*, pedunculis unifloris, foliis quinquepartitis trifidis orbiculatis. Fl. Sv. 616. P. pr. sylv.

2. P O L Y A N D R I A.

ALTHAEA.

852. *A. officinalis*, foliis simplicibus tomentosis. S. N. 624. Fl. D. 530. P.

MALVA.

853. *M. rotundifolia*, caule prostrato, foliis cordato-orbiculatis obsolete quinquelobatis, pedunculis fructiferis declinatis. Fl. Sv. 626. Fl. N. 1030. Fl. D. 721. A. r. v. f.
854. *M. sylvestris*, caule erecto herbaceo, foliis septemlobatis acutis, pedunculis petiolisque pilosis. Fl. Sv. 637. B. r. f.

MONADELPHIA, POLYANDRIA. Lavatera. 162

855. *M. Alcea*, caule erecto pilis fasciculatis ^{a)} scabro, foliis multipartitis scabriusculis. Fl. Sv. 628. P. pr. n.
856. *M. moschata*, caule erecto, foliis radicalibus reniformibus: caulinis quinquepartitis pinnato-multifidis. Fl. Sv. 629. Fl. D. 905. A. P. pr. n. hum.

LAVATERA.

857. *L. thuringiaca*, caule herbaceo, fructibus denudatis, calycibus incis. S. N. 627. Fn. Sv. 558. Jacqu. Fl. Austr. IV. t. 311. P. f.

^{a)} A nostra diversa viderur *M. Alcea Cavan. Diff. 2. p. 75.* quæ caule glabro et foliis etiam in horto glabris est; nostra enim in hortis minime asperitatem nec caulis neque foliorum deponit.

XVII. DIADELPHIA.

I. HEXANDRIA.

- FUMARIA.** *Cal.* 2-phyll. minim. *Cor.* ringens *nectario* gibbo vel cornuto. *Filam.* 2. membranacea, singulo *Antherae* 3.
Rad. bulbosa, solida l. fibrosa. *Caul.* herbacea.
Fol. composita, multifida. *Flor.* subspicata.
Fulcr. bracteae.

2. OCTANDRIA.

- POLYGALA.** *Cal.* 5-phyll. foliolis 2 alaeformibus coloratis, *Legum.* obcordat. 2-locul.
Rad. ramosa. *Caul.* herbacea adscendens. *Fol.* simplex, integra. *Flor.* racemosa. *Fulcr.* bracteae.

3. DECANDRIA.

- SPARTIUM.** *Cal.* deorsum productus. *Filam.* germini adhaerentia. *Stigma* longitudinale supra villosum.
Rad. lignosa. *Caul.* fruticulosa, prostrata. *Fol.* varia, caduca. *Flor.* solitaria. *Fulcr.*
GENISTA. *Cal.* 2-lab. *Vexill.* oblongum a pistillo, staminibus carinam deprimentibus, deorsum reflexum.
Rad. lignosa. *Caul.* fruticulosa. *Fol.* varia.
Flor. varia. *Fulcr.* stipulae.
ULEX. *Cal.* 2-phyll. *Legum.* vix calyce longius.
Rad. lignosa. *Caul.* frutescens, prostrata, ramolissima, spinosa. *Fol.* simplex, acerosa.
Flor. subcorymbosa. *Fulcr.* o.

ONONIS. *Cal.* 5-part. laciniis linearibus. *Vexill.* striatum. *Legum.* sessile, turgidum. *Filam.* connata absque fissura.

Rad. lignosa. *Caul.* herbacea. *Fol.* varia. *Flor.* racemosa. *Fulcr.* bracteae, stipulae.

ANTHYLLIS. *Cal.* ventricosus. *Legum.* subrotundum, tectum.

Rad. ramosa. *Caul.* herbacea. *Fol.* pinnata, inaequalis: *Flor.* capitata. *Fulcr.* bracteae, involucrum:

PISUM. *Stylus* triangulus, supra carinatus pubescens. *Cal.* 2-lab. superne brevior.

Rad. ramosa. *Caul.* herbacea, debilis. *Fol.* pinnata. *Flor.* varia. *Fulcr.* stipulae, cirrhi.

OROBUS. *Stylus* linearis. *Cal.* basi obtusus: laciniis superior. profundius divisis, brevioribus.

Rad. ramosa l. tuberosa. *Caul.* herbacea, erecta. *Fol.* pinnata. *Flor.* axillaris, subracemosa. *Fulcr.* stipulae.

LATHYRUS. *Stylus* planus, supra villosus, superne latior. *Cal.* 2-lab. superne brevior.

Rad. ramosa. *Caul.* herbacea, debilis, alata. *Fol.* subpinnata. *Flor.* axillaris, secunda. *Fulcr.* stipulae, cirrhi.

VICIA. *Stigm.* latere inferiore transverse barbatum.

Rad. fibrosa, ramosa. *Caul.* herbacea, debilis. *Fol.* pinnata, multijuga. *Flor.* racemosa, secunda l. subsessilis, axillaris. *Fulcr.* stipulae, cirrhi.

ERVUM. *Cal.* longitudine corollae.

Rad. ramosa. *Caul.* herbacea, debilis. *Fol.* pinnata, multijuga. *Flor.* pedunculata, axillaris. *Fulcr.* stipulae, cirrhi.

CORONILLA. *Cal.* 2-lab. dentibus 2 superioribus connatis. *Vexillum* vix alis longius. *Legumen* isthmis interceptum.

Rad. ramosa. *Caul.* fruticosa l. herbacea, procumbens. *Fol.* pinnata. *Flor.* umbellata. *Fulcr.* stipulae deciduae.

ORNITHOPUS. *Legum.* articulatum, teres, arcuatum. *Rad.* fibrosa. *Caul.* herbacea, procumbens. *Fol.* pinnata, multijuga. *Flor.* terminalis, sessilis. *Fulcr.* stipulae, bracteae.

HEDYSARUM. *Cor.* carina transverse obtusa. *Legumen* articulis 1-spermis.

Rad. ramosa. *Caul.* herbacea, ascendens. *Fol.* pinnata, multijuga. *Flor.* spicata. *Fulcr.* stipulae.

PHACA. *Legumen* semibiloculare.

Rad. ramosa. *Caul.* herbacea. *Fol.* pinnata. *Flor.* racemosa, laxa. *Fulcr.* stipulae, bracteae.

ASTRAGALUS. *Legumen* biloculare, gibbum.

Rad. ramosa. *Caul.* herbacea, varia. *Fol.* pinnata. *Flor.* racemosa. *Fulcr.* stipulae, bracteae.

TRIFOLIUM. *Flora* subcapitati. *Legumen* vix calyce longius, non dehiscens, deciduum.

Rad. subramosa. *Caul.* herbacea. *Fol.* ternata. *Flor.* spicata, l. subcapitata. *Fulcr.* stipulae.

LOTUS. *Legumen* cylindricum, strictum. *Alae* sursum longitudinaliter conniventes. *Cal.* tubulosus.

Rad. ramosa. *Caul.* herbacea. *Fol.* ternata. *Flor.* varia. *Fulcr.* stipulae, bracteae.

MEDICAGO. *Legumen* compressum, cochleatum. *Carina* corollae a vexillo deflectens.

Rad. subfusiformis. *Caul.* herbacea, procumbens. *Fol.* ternata. *Flor.* varia. *Fulcr.* stipulae.

XVII. DIADELPHIA.

I. H E X A N D R I A.

FUMARIA.

858. *F. bulbosa*, caule simplici, calcare rubro, bracteis cuneatis fissis, foliolis acutis, radice solida. *F. bulbosa*. Fl. Sv. 631. α. *F. solida*. Ehrh. Beytr. 6. p. 146. P. n. m.
859. *F. fabacea*, caule simplici, foliolis obtusis, calcare recto, bracteis ovatis obtusis integris, radice solida. *F. bulbosa*. Fl. Sv. 631. β. Fl. D. 605. *F. intermedia*. Ehrh. Beytr. 6. p. 146. P. n.
860. *F. cava*, caule simplici, foliolis obtusiusculis, calcare recurvo, bracteis lanceolatis acutis, radice cava. *F. bulbosa*. Fl. Sv. 631. γ. *F. cava*. Ehrh. Beytr. 6. p. 145. δ. P. n.
861. *F. officinalis*, pericarpis monospermis racemosis, caule diffuso. Fl. Sv. 630. Fl. N. 161. Fl. D. 940. A. c.
- † 862. *F. capreolata*, pericarpis monospermis racemosis, scandentibus, S. N. 637. Fl. D. 340. A. c.

2. O C T A N D R I A.

POLYGALA.

863. *P. amara*, floribus cristatis racemosis, caulibus erectiusculis, foliis radicalibus obovatis majoribus. S. N. 638. Jacqu. Fl. Austr. V. t. 412. p). P. pr. L 4. 864. P.

o) Omni jure hæc tres distinguuntur, quod cultura earum per XXX neque annos docuit, nec *fabacea bulbosae* est varietas, multo minus *cavae*, ut voluit ill. Haller.

p) Cel. Afzelius.

168. DIADELPHIA. DECANDRIA. Spartium.

864. *P. vulgaris*, floribus cristatis racemosis, caulibus herbaceis simplicibus procumbentibus, foliis linearilanceolatis. Fl. Sv. 632. Fl. N. 188. Fl. D. 516. P. pr. pasc.

α. coerulea.

β. carnea.

γ. alba.

865. *P. major*, floribus cristatis racemosis, caulibus herbaceis simplicibus suberectis, foliis linearilanceolatis. S. N. 638. Jacqu. Fl. Austr. V. t. 413. Vaill. Par. t. 32. f. 1. q). *P. vulgaris major II.* Clus. Hist. I. p. 325. ? P. pr.

† *P. chamaebuxi*, nomen habet uterque index Florae Norvegicae, in ipso opere autem desideratur.

DECANDRIA

SPARTIUM.

866. *S. scoparium*, foliis ternatis solitariisque, ramis intermibus angulatis. Fl. Sv. 633. Fl. D. 313. S. a.

GENISTA.

867. *G. tinctoria*, foliis lanceolatis glabris, ramis striatis teretibus erectis. Fl. Sv. 634. S.
868. *G. pilosa*, foliis lanceolatis obtusis, caule tuberculato decumbente. Fl. Sv. 635. Jacqu. Fl. Austr. III. t. 208. S. a. eric.
869. *G. anglica*, spinis simplicibus, ramis floriferis intermibus, foliis lanceolatis. S. N. 645. Fl. D. 619. S. er. a.

870. G.

q) Cum dubitatione hanc addo. Plantam ad Ekerö Scaniae legi fere erectam pedalem, omnibus partibus majorem, at folia nonnihil latiora sunt, quam in specimine ex alpinis Peuninis allato, in quo flores haud ut in nostra pendentes videntur, sed sub expansione primi floris quoque lectum fuit. Flores praeterea rubri, qui in nostra coerulesci.

870. *G. germanica*, spinis compositis, ramis floriferis inermibus, foliis lanceolatis. S. N. 645. Obs. Bot. Fasc. I. p. 23. Fl. D. 526. r). S. er. a.

ULEX.

- † 871. *U. europaeus*, foliis villosis acutis, spinis sparsis. S. N. 648. Fl. D. 608. S.

ONONIS.

872. *O. arvensis*, floribus racemosis geminis, foliis ternatis: superioribus solitariis, ramis inermibus subvillosis. Aët. Phys. V. 1. p. 2. p. 130. *Ononis mitior* I. Clus. Hist. I. p. 97. *On. arvensis et hirtina* S. N. 651. 3 et 10. P. agr. pasc. v.
873. *O. spinosa*, floribus racemosis solitariis, foliis ternatis solitariisque ramis spinosis. Aët. Phys. V. 1. p. 2. pag. 131. Obs. Bot. Fasc. II. 68. Fl. D. 783. P. agr.

ANTHYLLIS.

874. *A. Vulneraria*, herbacea, foliis pinnatis inaequalibus, capitulo duplicato. Fl. Sv. 638. Fl. N. 88. Fl. D. 988. P. pr.
- α. *lutea*.
β. *alba*.
γ. *rubra*.
δ. *coccinea*.

PISUM.

875. *P. arvense*, petiolis tetraphyllis, stipulis crenatis, pedunculis unifloris. Fl. Sv. 639. Fl. N. 761. A. agr.
876. *P. maritimum*, petiolis supra planiusculis, caule angulato, stipulis sagittatis, pedunculis multifloris. Fl. Sv. 640. Fl. N. 698. Fl. D. 338. P. l. mar.

L 5

ORO-

r) Icon cit. Fl. Dan. certe huc pertinet, minus licet idonea.

OROBUS.

877. *O. vernus*, foliis pinnatis ovatis, stipulis semisagittatis integerrimis, caule simplici. Fl. Sv. 641. Fl. N. 236. P. pr. n.
878. *O. tuberosus*, foliis pinnatis lanceolatis, stipulis semisagittatis integerrimis, caule simplici. Fl. Sv. 642. Fl. N. 983. Fl. D. 781. P. pr.
- β. *grandis*. Obs. Bot. Fasc. I. 79.
879. *O. niger*, caule ramoso, foliis sexjugis ovato-oblongis. Fl. Sv. 643. P. pr. sylv. n.

LATHYRUS.

- † 880. *L. tuberosus*, pedunculis multifloris, cirrhis diphyllis: foliolis ovalibus, internodiis nudis. S. N. 663. s). P. agr.
881. *L. pratensis*, pedunculis multifloris, cirrhis diphyllis simplicissimis, foliolis lanceolatis. Fl. Sv. 647. Fl. N. 263. Fl. D. 527. P. pr. sylv.
882. *L. sylvestris*, pedunculis multifloris, cirrhis diphyllis: foliis ensiformibus, internodiis membranaceis. Fl. Sv. 644. Fl. D. 325. P. pr. n.
- β. *platyphyllus*. Fl. D. 785. pr. mar.
883. *L. latifolius*, pedunculis multifloris, cirrhis diphyllis: foliolis ellipticis, internodiis membranaceis. Fl. Sv. 645. P. pr. sylv.
884. *L. heterophyllus*, pedunculis multifloris, cirrhis diphyllis tetraphyllisque: foliolis lanceolatis, internodiis membranaceis. Fl. Sv. 646. 1). P. pr. n.
885. *L. palustris*, pedunculis multifloris, cirrhis polyphyllis, stipulis lanceolatis. Fl. Sv. 648. Obs. Bot. Fasc. I. p. 24. Fl. N. 982. Fl. D. 399. P. pr. sylv. hum.

VICIA.

s) Cf. Hellwig et Kölpin.

*) Semina e Westrogothia missa *Lathyrum sylvestrem* dedere.

VICIA.

886. *V. pisiformis*, pedunculis multifloris, petiolis polyphyllis, foliolis ovatis: infimis sessilibus. S. N. 663. Jacqu. Fl. Austr. IV. t. 364. u). - P.
887. *V. dumetorum*, pedunculis multifloris, foliolis reflexis ovatis mucronatis, stipulis subdentatis. Fl. Sv. 649. Obs. Bot. Fasc. I. 81. P. n.
888. *V. sylvatica*, pedunculis multifloris, foliolis ovalibus, stipulis denticulatis. Fl. Sv. 650. Fl. N. 16. Fl. D. 277. P. n.
889. *V. Cassubica*, pedunculis subsexfloris, foliolis denis ovatis acutis, stipulis integris. Fl. Sv. 653. Obs. Bot. Fasc. I. 82. Fl. D. 98. w). *V. Gerardi* S. N. 664. Jacqu. Fl. Austr. III. t. 229. P. n. a.
890. *V. Cracca*, pedunculis multifloris, floribus imbricatis, foliolis lanceolatis pubescentibus, stipulis integris. Fl. Sv. 652. Fl. D. 806. P. pr. f. agr.
891. *V. sativa*, leguminibus sessilibus subbinatis erectis, foliis retusis, stipulis notatis. Fl. Sv. 654. Fl. D. 522. A. agr.
892. *V. Lathyroides*, leguminibus solitariis erectis glabris, foliolis senis inferioribus obcordatis. S. N. 664. Obs. Bot. Fasc. I. 83. Fl. D. 58. A. pr. f. c.
893. *V. sepium*, leguminibus pedicellatis subquaternis erectis, foliolis ovatis integerrimis: exterioribus decrescentibus. Fl. Sv. 651. Fl. D. 699. P. pr. f. n.

ERVUM.

- † 894. *E. Lens*, pedunculis subbifloris, seminibus compressis convexis. S. N. 665. Weig. Fl. 831. A. agr.
895. *E. tetraspermum*, pedunculis subbifloris, seminibus globosis quaternis. Fl. Sv. 655. Fl. D. 95. A. c.

896. *E.*

u) *Cel. Afzelius*.

w) Folia in hac icone minus aucta picta sunt.

172 DIADELPHIA. DECANDRIA. Coronilla.

896. *E. hirsutum*, pedunculis multifloris, seminibus globosis binis. Fl. Sv. 656. Fl. N. 535. Fl. D. 639. A. c.

CORONILLA.

897. *C. Emerus*, fruticosa, pedunculis subtrifloris, corollarum unguibus calyce triplo-longioribus, caule angulato. Fl. Sv. 636. S. mont.
 † 898. *C. varia*, herbacea, leguminibus erectis teretibus torosis numerosis, foliolis plurimis glabris. S. N. 670. Clus. Hist. II. p. 237. x). P.

ORNITHOPUS.

- † 899. *O. perpusillus*, foliis pinnatis, leguminibus incurvatis. S. N. 670. Weig. Fl. 471. Fl. D. 730. A. er. sicc.

HEDYSARUM.

- † 900. *H. Onobrychis*, foliis pinnatis, leguminibus monospermis aculeatis, corollarum alis calycem aequantibus, caule elongato. S. N. 676. Weig. Fl. 472. P.

PHACA.

901. *P. alpina*, caulescens erecta glabra, leguminibus oblongis inflatis subpilosis. Fl. Sv. 657. Fl. N. 968. Fl. D. 856. P. alp.

ASTRAGALUS.

902. *A. pilosus*, caulescens erectus pilosus, floribus spicatis, leguminibus subulatis pilosis. S. N. 681. Fn. Sv. p. 558. Jacqu. Fl. Austr. I. t. 51. P.
 903. *A. glycyphyllus*, caulescens prostratus, leguminibus subtriquetris arcuatis, foliolis ovalibus pedunculo longioribus. Fl. Sv. 658. Fl. N. 984. P. pr. fylv.

904. *A.*

x) Cel. Kölpin.

904. *A. arenarius*, caulescens, ramosus, erectus, tomentoso-albus, foliolis complicatis, pedunculis subquadrifloris. Obs. Bot. Fasc. III. 87. t. 3. et II. 73. *A. arenarius*. Fl. Sv. 659. it. Scân. 108. P. a.
- † 905. *A. Danicus*, caulescens, decumbens, hirtus capitulis longe pedunculatis, leguminibus brevibus inflatis pilosis. Obs. Bot. Fasc. III. 86. *A. arenarius* Fl. D. 614. P.
906. *A. alpinus*, caulescens procumbens, floribus pendulis racemosis, leguminibus utrinque acutis pilosis. Fl. Sv. 661. Fl. L. t. 9. f. 1. Fl. N. 665. Fl. D. 51. P. alp.
907. *A. Uralensis*, acaulis, scapo erecto foliis longiore, leguminibus subulatis inflatis villosis erectis. Fl. Sv. 660. St. Helv. t. 14. P. a.
908. *A. Campestris*, acaulis, calycibus leguminibusque pilosis, foliolis lanceolatis acutis, scapo decumbente. Fl. Sv. 662. Stirp. Helv. t. 13. P. ficc.

TRIFOLIUM.

909. *T. Melilotus officinalis*, leguminibus racemosis nudis dispermis rugosis acutis, caule erecto. Fl. Sv. 683. Fl. D. 934. B. pr. v.
- † 910. *T. Melilotus Ornithopodioides*, leguminibus nudis octospermis subternis calyce duplo longioribus, caulibus declinatis. S. N. 687. Fl. D. 368. A.
911. *T. hybridum*, capitulis umbellaribus, leguminibus tetraspermis, caule ascendente. Fl. Sv. 664. P. pr. pasc.
912. *T. repens*, capitulis umbellaribus, leguminibus tetraspermis, caule repente. Fl. Sv. 665. Fl. N. 131. Fl. D. 990. P. pr. pasc.
913. *T. pratense*, spicis densis, corollis inaequalibus, dentibus calycinis quatuor aequalibus, stipulis aristatis, caulibus ascendentibus. A. Soc. Lin. I. Fl. Sv. 666. Fl. N. 130. Fl. D. 989. P. pr.

174 DIADELPHIA. DECANDRIA. Trifolium.

914. *T. flexuosum*, spicis laxis, corollis subaequalibus, stipulis subulatis conniventibus, caulibus flexuosis ramosis. A. Soc. Lin. I. Jacqu. Fl. Austr. IV. t. 386. *Tr. medium* Fn. Sv. 558. P. pr.
915. *T. alpestre*, spicis densis, corollis subaequalibus, stipulis setaceis divergentibus, foliolis lanceolatis, caulibus strictis simplicissimis. A. Soc. Lin. I. S.N. 688. Fl. D. 662. y). P. pr.
916. *T. arvense*, spicis villosis ovalibus, dentibus calycinis setaceis villosis aequalibus. Fl. Sv. 668. Fl. N. 357. Fl. D. 724. A. agr. pasc.
- β. *pumilum*. Raji Angl. 3. t. 14. f. 2.
917. *T. striatum*, capitulis sessilibus sublateralibus ovatis, calycibus striatis rotundatis. Fl. Sv. 669. Vaill. Par. t. 33. f. 2. A. agr. arg.
918. *T. fragiferum*, spicis subrotundis, calycibus inflatis bidentatis reflexis, caulibus repentibus. Fl. Sv. 670. Vaill. Par. t. 22. f. 2. P. pasc. agr. arg.
919. *T. montanum*, spicis subimbricatis subternis, vexillis subulatis marcescentibus, calycibus nudis, caule erecto. Fl. Sv. 667. Fl. N. 621. P. pr. sicc.
920. *T. agrarium*, spicis ovalibus imbricatis, vexillis deflexis persistentibus calycibus nudis, caule erecto. Fl. Sv. 671. Fl. N. 529. Fl. D. 558. A. pr. sicc.
921. *T. spadiceum*, spicis ovalibus imbricatis, vexillis deflexis persistentibus, calycibus pilosis, caule erecto. Fl. Sv. 672. A. pr. mont.
922. *T. procumbens*, spicis ovalibus imbricatis, vexillis deflexis persistentibus, caulibus procumbentibus. Fl. Sv. 673. Fl. D. 796. Raji Angl. 3. t. 14. f. 3. A. pr. pasc.
923. *T. filiforme*, spicis subimbricatis, vexillis deflexis persistentibus, calycibus pedicellatis, caulibus procumbentibus. Fl. Sv. 674. Raji Angl. 3. t. 4. f. 4. A. pasc.

LOTUS.

y) Etiam hae notae rix sufficiunt. Conf. *Gel. Afr.* Diff. p. 31.

LOTUS.

924. *L. maritima*, leguminibus solitariis membranaceo quadrangulis, foliis glabris, bracteis lanceolatis. Fl. Sv. 676. Fl. D. 800. P. l. mar.
925. *L. corniculata*, capitulis depresso, caulibus decumbentibus, leguminibus cylindricis patentibus. Fl. Sv. 675. Fl. N. 108. Fl. D. 991. P. pr. pasc.
 β . *procerior*. Rof. Obs. p. 38.
 γ . *tenuifolia erectiuscula*. it. Scan. p. 153.

MEDICAGO.

926. *M. falcata*, pedunculis racemosis, leguminibus lunatis, caule prostrato. Fl. Sv. 677. Fl. N. 648. Fl. D. 233. P. sicc. arg.
927. *M. Lupulina*, spicis ovalibus, leguminibus reniformibus monospermis, caulibus procumbentibus. Fl. Sv. 678. Fl. D. 992. B. pr. c.
- †928. *M. polymorpha*, leguminibus cochleatis, caule diffuso. S. N. 693. A.
 λ . *minima*, leguminibus subternis, aculeis uncinatis, stipulis integris. Fl. D. 211.

COMONILLA. *Cal.* 2-lab. dentibus 2 superioribus connatis. *Vexillum* vix alis longius. *Legumen* isthmis interceptum.

Rad. ramosa. *Caul.* fruticosa l. herbacea, procumbens. *Fol.* pinnata. *Flor.* umbellata. *Fulcr.* stipulae deciduae.

ORNITHOPUS. *Legum.* articulatum, teres, arcuatum.

Rad. fibrosa. *Caul.* herbacea, procumbens. *Fol.* pinnata, multijuga. *Flor.* terminalis, sessilis. *Fulcr.* stipulae, bracteae.

HEDYSARUM. *Cor.* carina transverse obtusa. *Legumen* articulis 1-spermis.

Rad. ramosa. *Caul.* herbacea, ascendens. *Fol.* pinnata, multijuga. *Flor.* spicata. *Fulcr.* stipulae.

PHACA. *Legumen* semibiloculare.

Rad. ramosa. *Caul.* herbacea. *Fol.* pinnata. *Flor.* racemosa, laxa. *Fulcr.* stipulae, bracteae.

ASTRAGALUS. *Legumen* biloculare, gibbum.

Rad. ramosa. *Caul.* herbacea, varia. *Fol.* pinnata. *Flor.* racemosa. *Fulcr.* stipulae, bracteae.

TRIFOLIUM. *Flores* subcapitati. *Legumen* vix calyce longius, non dehiscens, deciduum.

Rad. subramosa. *Caul.* herbacea. *Fol.* ternata. *Flor.* spicata, l. subcapitata. *Fulcr.* stipulae.

LOTUS. *Legumen* cylindricum, strictum. *Alae* sursum longitudinaliter conniventes. *Cal.* tubulosus.

Rad. ramosa. *Caul.* herbacea. *Fol.* ternata. *Flor.* varia. *Fulcr.* stipulae, bracteae.

MEDICAGO. *Legumen* compressum, cochleatum. *Carina* corollae a vexillo deflectens.

Rad. subfusiformis. *Caul.* herbacea, procumbens. *Fol.* ternata. *Flor.* varia. *Fulcr.* stipulae.

XVII. DIADELPHIA.

I. H E X A N D R I A.

FUMARIA.

858. *F. bulbosa*, caule simplici, calcare rubro, bracteis cuneatis fissis, foliolis acutis, radice solida. *F. bulbosa*. Fl. Sv. 631. α . *F. solida*. Ehrh. Beytr. 6. p. 146. P. n. m.
859. *F. fabacea*, caule simplici, foliolis obtusis, calcare recto, bracteis ovatis obtusis integris, radice solida. *F. bulbosa*. Fl. Sv. 631. β . Fl. D. 605. *F. intermedia*. Ehrh. Beytr. 6. p. 146. P. n.
860. *F. cava*, caule simplici, foliolis obtusiusculis, calcare recurvo, bracteis lanceolatis acutis, radice cava. *F. bulbosa*. Fl. Sv. 631. γ . *F. cava*. Ehrh. Beytr. 6. p. 145. δ . P. n.
861. *F. officinalis*, pericarpis monospermis racemosis, caule diffuso. Fl. Sv. 630. Fl. N. 161. Fl. D. 940. A. c.
- † 862. *F. capreolata*, pericarpis monospermis racemosis, scandentibus, S. N. 637. Fl. D. 340. A. c.

2. O C T A N D R I A.

POLYGALA.

863. *P. amara*, floribus cristatis racemosis, caulibus erectiusculis, foliis radicalibus obovatis majoribus. S. N. 638. Jacqu. Fl. Austr. V. t. 412. p). P. pr. L. 4. 864. P.

o) Omni jure hæ tres distinguuntur, quod cultura earum per XXX neque annos docuit, nec *fabacea bulbosae* est varietas, multo minus *cavae*, ut voluit ill. Haller.

p) Cel. Afzelius.

168. DIADELPHIA. DECANDRIA. *Spartium*.

864. *P. vulgaris*, floribus cristatis racemosis, caulibus herbaceis simplicibus procumbentibus, foliis linearilanceolatis. Fl. Sv. 632. Fl. N. 188. Fl. D. 516. P. pr. pasc.

α. *coerulea*.

β. *carnea*.

γ. *alba*.

865. *P. major*, floribus cristatis racemosis, caulibus herbaceis simplicibus suberectis, foliis lineari lanceolatis. S. N. 638. Jacqu. Fl. Austr. V. t. 413. Vaill. Par. t. 32. f. 1. q). *P. vulgaris major* II. Clus. Hist. I. p. 325. ? P. pr.

† *P. chamaebuxi*, nomen habet uterque index Florae Norvegicae, in ipso opere autem desideratur.

3. DECANDRIA

SPARTIUM.

866. *S. scoparium*, foliis ternatis solitariisque, ramis intermibus angulatis. Fl. Sv. 633. Fl. D. 313. S. a.

GENISTA.

867. *G. tinctoria*, foliis lanceolatis glabris, ramis striatis teretibus erectis. Fl. Sv. 634. S.

868. *G. pilosa*, foliis lanceolatis obtusis, caule tuberculato decumbente. Fl. Sv. 635. Jacqu. Fl. Austr. III. t. 208. S. a. eric.

869. *G. anglica*, spinis simplicibus, ramis floriferis intermibus, foliis lanceolatis. S. N. 645. Fl. D. 619. S. er. a.

870. G.

q) Cum dubitatione hanc addo. Plantam ad Ekerö Scaniae legi fere erectam pedalem, omnibus partibus majorem, at folia nonnihil latiora sunt, quam in specimine ex alpinis Peuninis allato, in quo flores haud ut in nostra pendentes videntur, sed sub expansione primi floris quoque lectum fuit. Flores praeterea rubri, qui in nostra coerulei.

870. *G. germanica*, spinis compositis, ramis floriferis inermibus, foliis lanceolatis. S. N. 645. Obs. Bot. Fasc. I. p. 23. Fl. D. 526. r). S. er. a.

ULEX.

- † 871. *U. europaeus*, foliis villosis acutis, spinis sparsis. S. N. 648. Fl. D. 608. S.

ONONIS.

872. *O. arvensis*, floribus racemosis geminis, foliis ternatis: superioribus solitariis, ramis inermibus subvillosis. Aët. Phys. V. 1. p. 2. p. 130. *Ononis mitior* I. Clus. Hist. I. p. 97. *On. arvensis et hirtina* S. N. 651. 3 et 10. P. agr. pasc. v.
873. *O. spinosa*, floribus racemosis solitariis, foliis ternatis solitariisque ramis spinosis. Aët. Phys. V. 1. p. 2. pag. 131. Obs. Bot. Fasc. II. 68. Fl. D. 783. P. agr.

ANTHYLLIS.

874. *A. Vulneraria*, herbacea, foliis pinnatis inaequalibus, capitulo duplicato. Fl. Sv. 638. Fl. N. 88. Fl. D. 988. P. pr.
- α. *lutea*.
β. *alba*.
γ. *rubra*.
δ. *coccinea*.

PISUM.

875. *P. arvense*, petiolis tetraphyllis, stipulis crenatis, pedunculis unifloris. Fl. Sv. 639. Fl. N. 761. A. agr.
876. *P. maritimum*, petiolis supra planiusculis, caule angulato, stipulis sagittatis, pedunculis multifloris. Fl. Sv. 640. Fl. N. 698. Fl. D. 338. P. l. mar.

L 5

ORO-

r) Icon cit. Fl. Dan. certe huc pertinet, minus licet idonea.

OROBUS.

877. *O. vernus*, foliis pinnatis ovatis, stipulis semigittatis integerrimis, caule simplici. Fl. Sv. 641. Fl. N. 236. P. pr. n.
878. *O. tuberosus*, foliis pinnatis lanceolatis, stipulis semilagittatis integerrimis, caule simplici. Fl. Sv. 642. Fl. N. 983. Fl. D. 781. P. pr.
- β. *grandis*. Obs. Bot. Fasc. I. 79.
879. *O. niger*, caule ramoso, foliis sexjugis ovato-oblongis. Fl. Sv. 643. P. pr. sylv. n.

LATHYRUS.

880. *L. tuberosus*, pedunculis multifloris, cirrhis diphyllis: foliolis ovalibus, internodiis nudis. S. N. 663. s). P. agr.
881. *L. pratensis*, pedunculis multifloris, cirrhis diphyllis simplicissimis, foliolis lanceolatis. Fl. Sv. 647. Fl. N. 263. Fl. D. 527. P. pr. sylv.
882. *L. sylvestris*, pedunculis multifloris, cirrhis diphyllis: foliis ensiformibus, internodiis membranaceis. Fl. Sv. 644. Fl. D. 325. P. pr. n.
- β. *platyphyllos*. Fl. D. 785. pr. mar.
883. *L. latifolius*, pedunculis multifloris, cirrhis diphyllis: foliolis ellipticis, internodiis membranaceis. Fl. Sv. 645. P. pr. sylv.
884. *L. heterophyllus*, pedunculis multifloris, cirrhis diphyllis tetraphyllisque: foliolis lanceolatis, internodiis membranaceis. Fl. Sv. 646. t). P. pr. n.
885. *L. palustris*, pedunculis multifloris, cirrhis polyphyllis, stipulis lanceolatis. Fl. Sv. 648. Obs. Bot. Fasc. I. p. 24. Fl. N. 982. Fl. D. 399. P. pr. sylv. hum.

VICIA.

s) Cf. Hellwig et Kölpin.

t) Semina e Westrogothia missa *Lathyrum sylvestrem* dedere.

VICIA.

886. *V. pisiformis*, pedunculis multifloris, petiolis polyphyllis, foliolis ovatis: infimis sessilibus. S. N. 663. Jacqu. Fl. Austr. IV. t. 364. u). - P.
887. *V. dumetorum*, pedunculis multifloris, foliolis reflexis ovatis mucronatis, stipulis subdentatis. Fl. Sv. 649. Obs. Bot. Fasc. I. 81. P. n.
888. *V. sylvatica*, pedunculis multifloris, foliolis ovalibus, stipulis denticulatis. Fl. Sv. 650. Fl. N. 16. Fl. D. 277. P. n.
889. *V. Cassubica*, pedunculis subsexfloris, foliolis denis ovatis acutis, stipulis integris. Fl. Sv. 653. Obs. Bot. Fasc. I. 82. Fl. D. 98. w). *V. Gerardi* S. N. 664. Jacqu. Fl. Austr. III. t. 229. P. n. a.
890. *V. Cracca*, pedunculis multifloris, floribus imbricatis, foliolis lanceolatis pubescentibus, stipulis integris. Fl. Sv. 652. Fl. D. 806. P. pr. f. agr.
891. *V. sativa*, leguminibus sessilibus subbinatis erectis, foliis retusis, stipulis notatis. Fl. Sv. 654. Fl. D. 522. A. agr.
892. *V. Lathyroides*, leguminibus solitariis erectis glabris, foliolis senis inferioribus obcordatis. S. N. 664. Obs. Bot. Fasc. I. 83. Fl. D. 58. A. pr. f. c.
893. *V. sepium*, leguminibus pedicellatis subquaternis erectis, foliolis ovatis integerrimis: exterioribus decrescentibus. Fl. Sv. 651. Fl. D. 699. P. pr. f. n.

ERVUM.

- † 894. *E. Lens*, pedunculis subbifloris, seminibus compressis convexis. S. N. 665. Weig. Fl. 831. A. agr.
895. *E. tetraspermum*, pedunculis subbifloris, seminibus globosis quaternis. Fl. Sv. 655. Fl. D. 95. A. c.

896. *E.*

u) Cel. Afzelius.

w) Folia in hac icone minus aucta picta sunt.

172 DIADELPHIA. DECANDRIA. Coronilla.

896. *E. hirsutum*, pedunculis multifloris, seminibus globosis binis. Fl. Sv. 656. Fl. N. 535. Fl. D. 639. A. c.

CORONILLA.

897. *C. Emerus*, fruticosa, pedunculis subtrifloris, corollarum unguibus calyce triplo-longioribus, caule angulato. Fl. Sv. 636. S. mont.
† 898. *C. varia*, herbacea, leguminibus erectis teretibus torosis numerosis, foliolis plurimis glabris. S. N. 670. Clus. Hist. II. p. 237. x) P.

ORNITHOPUS.

- † 899. *O. perpusillus*, foliis pinnatis, leguminibus incurvatis. S. N. 670. Weig. Fl. 471. Fl. D. 730. A. er. ficc.

HEDYSARUM.

- † 900. *H. Onobrychis*, foliis pinnatis, leguminibus monospermis aculeatis, corollarum alis calycem aequantibus, caule elongato. S. N. 676. Weig. Fl. 472. P.

PHACA.

901. *P. alpina*, caulescens erecta glabra, leguminibus oblongis inflatis subpilosis. Fl. Sv. 657. Fl. N. 968. Fl. D. 856. P. alp.

ASTRAGALUS.

902. *A. pilosus*, caulescens erectus pilosus, floribus spicatis, leguminibus subulatis pilosis. S. N. 681. Fn. Sv. p. 558. Jacqu. Fl. Austr. I. t. 51. P.
903. *A. glycyphyllus*, caulescens prostratus, leguminibus subtriquetris arcuatis, foliolis ovalibus pedunculo longioribus. Fl. Sv. 658. Fl. N. 984. P. pr. sylv.

904. *A.*

x) Cel. Kölpin.

904. *A. arenarius*, caulescens, ramosus, erectus, tomentoso-albus, foliolis complicatis, pedunculis subquadrifloris. Obs. Bot. Fasc. III. 87. t. 3. et II. 73. *A. arenarius*. Fl. Sv. 659. it. Scän. 108. P. a.
- † 905. *A. Danicus*, caulescens, decumbens, hirtus capitulis longe pedunculatis, leguminibus brevibus inflatis pilosis. Obs. Bot. Fasc. III. 86. *A. arenarius* Fl. D. 614. P.
906. *A. alpinus*, caulescens procumbens, floribus pendulis racemosis, leguminibus utrinque acutis pilosis. Fl. Sv. 661. Fl. L. t. 9. f. 1. Fl. N. 665. Fl. D. 51. P. alp.
907. *A. Uralensis*, acaulis, scapo erecto foliis longiore, leguminibus subulatis inflatis villosis erectis. Fl. Sv. 660. St. Helv. t. 14. P. a.
908. *A. Campestris*, acaulis, calycibus leguminibusque pilosis, foliolis lanceolatis acutis, scapo decumbente. Fl. Sv. 662. Stirp. Helv. t. 13. P. ficc.

TRIFOLIUM.

909. *T. Melilotus officinalis*, leguminibus racemosis nudis dispermis rugosis acutis, caule erecto. Fl. Sv. 683. Fl. D. 934. B. pr. v.
- † 910. *T. Melilotus Ornithopodioides*, leguminibus nudis octospermis subternis calyce duplo longioribus, caulibus declinatis. S. N. 687. Fl. D. 368. A.
911. *T. hybridum*, capitulis umbellaribus, leguminibus tetraspermis, caule adscendente. Fl. Sv. 664. P. pr. pasc.
912. *T. repens*, capitulis umbellaribus, leguminibus tetraspermis, caule repente. Fl. Sv. 665. Fl. N. 131. Fl. D. 990. P. pr. pasc.
913. *T. pratense*, spicis densis, corollis inaequalibus, dentibus calycinis quatuor aequalibus, stipulis aristatis, caulibus adscendentibus. A. Soc. Lin. I. Fl. Sv. 666. Fl. N. 130. Fl. D. 989. P. pr.

174 DIADELPHIA. DECANDRIA. Trifolium.

914. *T. flexuosum*, spicis laxis, corollis subaequalibus, stipulis subulatis conniventibus, caulibus flexuosis ramosis. A. Soc. Lin. I. Jacqu. Fl. Austr. IV. t. 386. *Tr. medium* Fr. Sv. 558. P. pr.
915. *T. alpestre*, spicis densis, corollis subaequalibus, stipulis setaceis divergentibus, foliolis lanceolatis, caulibus strictis simplicissimis. A. Soc. Lin. I. S. N. 688. Fl. D. 662. y). P. pr.
916. *T. arvense*, spicis villosis ovalibus, dentibus calycinis setaceis villosis aequalibus. Fl. Sv. 668. Fl. N. 357. Fl. D. 724. A. agr. pasc.
- β. *pumilum*. Raji Angl. 3. t. 14. f. 2.
917. *T. striatum*, capitulis sessilibus sublateralibus ovatis, calycibus striatis rotundatis. Fl. Sv. 669. Vaill. Par. t. 33. f. 2. A. agr. arg.
918. *T. fragiferum*, spicis subrotundis, calycibus inflatis bidentatis reflexis, caulibus repentibus. Fl. Sv. 670. Vaill. Par. t. 22. f. 2. P. pasc. agr. arg.
919. *T. montanum*, spicis subimbricatis subternis, vexillis subulatis marcescentibus, calycibus nudis, caule erecto. Fl. Sv. 667. Fl. N. 621. P. pr. sicc.
920. *T. agrarium*, spicis ovalibus imbricatis, vexillis deflexis persistentibus calycibus nudis, caule erecto. Fl. Sv. 671. Fl. N. 529. Fl. D. 558. A. pr. sicc.
921. *T. spadiceum*, spicis ovalibus imbricatis, vexillis deflexis persistentibus, calycibus pilosis, caule erecto. Fl. Sv. 672. A. pr. mont.
922. *T. procumbens*, spicis ovalibus imbricatis, vexillis deflexis persistentibus, caulibus procumbentibus. Fl. Sv. 673. Fl. D. 796. Raji Angl. 3. t. 14. f. 3. A. pr. pasc.
923. *T. filiforme*, spicis subimbricatis, vexillis deflexis persistentibus, calycibus pedicellatis, caulibus procumbentibus. Fl. Sv. 674. Raji Angl. 3. t. 4. f. 4. A. pasc.

LOTUS.

y) Etiam hae notae vix sufficiunt. Conf. *Gel. Afr.* Diff. p. 31.

LOTUS.

924. *L. maritima*, leguminibus solitariis membranaceo quadrangulis, foliis glabris, bracteis lanceolatis. Fl. Sv. 676. Fl. D. 800. P. l. mar.
 925. *L. corniculata*, capitulis depressis, caulibus decumbentibus, leguminibus cylindricis patentibus. Fl. Sv. 675. Fl. N. 108. Fl. D. 991. P. pr. pasc.
 β . *procerior*. Ros. Obs. p. 38.
 γ . *tenuifolia erectiuscula*. it. Scan. p. 153.

MEDICAGO.

926. *M. falcata*, pedunculis racemosis, leguminibus lunatis, caule prostrato. Fl. Sv. 677. Fl. N. 648. Fl. D. 233. P. ficc. arg.
 927. *M. Lupulina*, spicis ovalibus, leguminibus reniformibus monospermis, caulibus procumbentibus. Fl. Sv. 678. Fl. D. 992. B. pr. c.
 †928. *M. polymorpha*, leguminibus cochleatis, caule diffuso. S. N. 693. A.
 λ . *minima*, leguminibus subternis, aculeis uncinatis, stipulis integris. Fl. D. 211.

XVIII. POLYADELPHIA.

I. POLYANDRIA.

HYPERICUM. *Cal.* 5-part. *Pet.* 5. *Filamenta* multa in 5 phalanges basi connata. *Caps.* polysperma. *Rad.* ramosa. *Caul.* herbacea. *Fol.* simplex, opposita, punctata. *Flor.* corymbosa. *Fulcr.* bracteae.

XVIII. POLYADELPHIA.

I. P O L Y A N D R I A.

HYPERICUM.

929. *H. quadrangulare*, floribus trigynis, caule quadrato herbaceo. Fl. Sv. 679. Fl. N. 266. Fl. D. 640. P. f. pr.
930. *H. perforatum*, floribus trigynis, caule ancipiti, foliis obtusis pellucido punctatis. Fl. Sv. 680. Fl. N. 267. P. f. n.
931. *H. humifusum*, floribus trigynis axillaribus solitariis, caulibus ancipitibus prostratis filiformibus, foliis glabris. Fl. Sv. 681. Fl. D. 141. Weig. Fl. 486. P. pr. hum.
932. *H. montanum*, floribus trigynis, calycibus ferrato-glandulosis, caule tereti erecto, foliis ovatis glabris. Fl. Sv. 683. Fl. D. 173. P. pr. mont.
933. *H. hirsutum*, floribus trigynis, caulibus ferrato-glandulosis, caule tereti erecto, foliis ovatis subpubescentibus. Fl. Sv. 682. Fl. D. 802. P. pr. mont.
- †934. *H. pulchrum*, floribus tryginis calycibus ferrato-glandulosis, foliis cordatis glabris amplexicaulibus, caule tereti. S. N. 702. Fl. D. 75. P.

XIX. SYNGENESIA.

I. POLYGAMIA AEQUALIS.

a. flor. semiflosculosi.

TRAGOPOGON. *Rec. nudum. Cal. simplex. Papp. plumosus, stipitatus.*

Rad. fusiformis. Caul. herbacea. Fol. simplex, amplexicaulis. Flor. terminalis. Fulcr. o.

SCORZONERA. *Rec. nudum. Cal. imbricatus, squamis margine scariosis. Papp. plumosus, stipitatus.*

Rad. fusiformis. Caul. herbacea. Fol. simplex, radicalis. Flor. terminalis, solitaria. Fulcr. o.

PICRIS. *Rec. nudum. Cal. calyculatus. Papp. plumosus, stipitatus. Sem. transversim fulcata.*

Rad. subfusiformis. Caul. herbacea. Fol. sinuata, hispida. Flor. terminalis. Fulcr. bracteae.

SONCHUS. *Rec. nudum. Cal. imbricatus, ventricosus. Papp. pilosus, sessilis.*

Rad. subfusiformis. Caul. herbacea. Fol. sinuata l. integra. Flor. subcorymbosa. Fulcr. o.

LACTUCA. *Rec. nudum. Cal. imbricatus, cylindricus, margine membranaceo. Papp. simplex, stipitatus.*

Rad. subfusiformis. Caul. herbacea. Fol. subhastata, amplexicaulis. Flor. corymbosa. Fulcr. o.

PRENANTHES. *Rec. nudum. Cal. calyculatus. Papp. simplex. Flosculi simplici serie.*

Rad. fusiformis. Caul. herbacea. Fol. simplex, amplexicaulis. Flor. paniculata. Fulcr. o.

CEONDRILLA. *Rec. nudum. Papp. pilosus, stipitatus. Cal. calyculat. multiflorus.*

Rad. ramosa. Caul. herbacea. Fol. simplex, varia. Flor. paniculata. Fulcr. o.

LEONTO-

- LEONTODON.** *Rec.* nudum. *Cal.* imbricatus, squamis laxiusculis. *Papp.* capillaris.
Rad. ramosa. *Caul.* scaposa. *Fol.* simplex. *Flor.* solitaria. *Fulcr.* o.
- HIERACIUM.** *Rec.* nudum. *Cal.* imbricatus, ovatus. *Papp.* simplex, sessilis.
Rad. varia. *Caul.* herbacea, scaposa l. foliosa. *Fol.* simplex, subintegra. *Flor.* varia. *Fulcr.* o.
- CREPIS.** *Rec.* nudum. *Cal.* calyculat. squamis difformibus deciduis. *Papp.* pilosus.
Rad. ramosa. *Caul.* herbacea. *Fol.* runcinata. *Flor.* paniculata. *Fulcr.* o.
- HYOSERIS.** *Rec.* nudum. *Cal.* subaequalis. *Papp.* pilosus calyculatus.
Rad. ramosa. *Caul.* scaposa, ramosa. *Fol.* radicalis. *Flor.* solitaria. *Fulcr.* bracteae.
- LAPSANA.** *Rec.* nudum. *Cal.* calyculatus, squamis singulis interioribus canaliculatis.
Rad. ramosa. *Caul.* herbacea. *Fol.* sinuata. *Flor.* paniculata. *Fulcr.* o.
- APARGIA.** *Rec.* subnudum, alveolis marginatis. *Papp.* sessilis plumosus. *Cal.* imbricat. erectus.
Rad. ramosa. *Caul.* herbacea, varia. *Fol.* simplex. *Flor.* varia. *Fulcr.* o.
- CICHORIUM.** *Rec.* subpaleaceum. *Cal.* calyculatus. *Papp.* sub 5-dentatus, obsolete pilosus.
Rad. ramosa. *Caul.* herbacea, rigida. *Fol.* sinuata. *Flor.* sessilis, caulina. *Fulcr.* o.
- HYPOCHAERIS.** *Rec.* paleaceum. *Cal.* subimbricatus. *Papp.* plumosus.
Rad. subramosa. *Caul.* herbacea. *Fol.* radicalis fere. *Flor.* varia. *Fulcr.* o.

β. capitati.

ARCTIUM. *Cal.* globosus, squamis apice hamis inflexis.

Rad. fusiformis. *Caul.* herbacea. *Fol.* simplex, integra. *Flor.* terminalis et axillaris. *Fulcr.* o.

SERRATULA. *Cal.* subcylindricus, imbricatus, muticus.

Rad. ramosa. *Caul.* herbacea. *Fol.* varia. *Flor.* subcorymbosa. *Fulcr.* o.

CARDUUS. *Cal.* ovatus, imbricatus squamis spinosis, *Recept.* pilosum, *Papp.* pilosus.

Rad. varia. *Caul.* herbacea. *Fol.* decurrens, spinosa, rarius inermis sessilis. *Flor.* terminalis. *Fulcr.* o.

CNICUS. *Cal.* ovatus, imbricatus squamis spinosis. *Recept.* villosum. *Papp.* plumosus.

Rad. ramosa. *Caul.* herbacea. *Fol.* varia. *Flor.* terminalis. *Fulcr.* o. l. bracteae.

ONOPORDUM. *Cal.* squamis mucronatae. *Rec. favosum.*

Rad. fusiformis. *Caul.* herbacea, alata. *Fol.* decurrens, spinosa. *Flor.* subsolitaria. *Fulcr.* o.

CARLINA. *Cal.* radiatus squamis marginalibus longis coloratis.

Rad. fusiformis. *Caul.* herbacea. *Fol.* spinosa. *Flor.* corymbosa. *Fulcr.* o.

γ. discoidei.

BIDENS. *Rec.* paleaceum. *Papp.* aristis erectis scabris. *Cal.* imbricat.

Rad. fibrosa. *Caul.* herbacea. *Fol.* opposita. *Flor.* varia. *Fulcr.* o. nisi folia floralia.

EUPATORIUM. *Recept.* nudum. *Papp.* plumosus. *Cal.* imbricat. oblongus. *Styl.* longus semibifidus.

Rad. ramosa. *Caul.* herbacea. *Fol.* opposita, composita. *Flor.* corymbosa. *Fulcr.* o.

CHRYSOCOMA. *Rec.* nudum. *Papp.* simplex. *Cal.* hemisphaericus, imbricatus. *Stylus* vix flosculis longior.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex, sparsa. *Flor.* corymbosa. *Fulcr.* o.

2. POLYGAMIA SUPERFLUA.

a. discoidei.

TANACEUM. *Recept.* nudum. *Papp.* submarginatus, *Cal.* hemisphaer. imbricat. *Cor.* radii obsoletae, trifidae.

Rad. ramosa. *Caul.* herbacea. *Fol.* pinnata. *Flor.* corymbosa. *Fulcr.* o.

ARTEMISIA. *Recept.* subvillos. l. nudiuscul. *Papp.* o. *Cal.* imbricatus squamis rotundatis conniventibus. *Cor.* radii o.

Rad. ramosa. *Caul.* herbacea l. fruticulosa. *Fol.* multifido-composita. *Flor.* racemosa. *Fulcr.* o.

GNAPHALIUM. *Rec.* nudum. *Papp.* plumos. *Cal.* imbricat. squamis marginalibus rotundatis scariosis, coloratis.

Rad. fibrosa. *Caul.* herbacea. *Fol.* simplex. *Flor.* corymbosa l. racemosa. *Fulcr.* o.

CONYZA. *Rec.* nudum. *Papp.* simplex. *Cal.* imbricat, subrotund. *Cor.* radii trifidae.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex, sessilis. *Flor.* pedunculata. *Fulcr.* o.

β. radiati.

ERIGERON. *Rec.* nudum. *Papp.* pilosus. *Cor.* radii lineares, angustissimae.

Rad. subramosa. *Caul.* herbacea. *Fol.* simplex. *Flor.* varia. *Fulcr.* o.

TUSSILAGO. *Rec.* nudum. *Papp.* simplex. *Cal.* squamae aequales, discum aequantes, submembranaeae. *Cor.* radii interdum nullae.

Rad. repens. *Caul.* scaposa. *Fol.* simplex, radicalis, subcordata. *Flor.* solitaria l. thyrsoides. *Fulcr.* bracteae.

SENECIO. *Rec.* nudum. *Papp.* simplex. *Cal.* cylindricus, calyculatus, squamis apice sphacelatis. *Cor.* radii interdum o.

Rad. ramosa. *Caul.* herbacea. *Fol.* sinuata l. integra. *Flor.* subcorymbosa. *Fulcr.* o.

ASTER. *Rec.* nudum. *Papp.* simplex. *Cor.* radii plures 10. *Cal.* imbricat. squamis inferioribus patulis.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex, alterna. *Flor.* subpaniculata. *Fulcr.* o.

SOLIDAGO. *Rec.* nudum. *Papp.* simplex. *Cor.* radii circiter 5. *Cal.* squamae imbricat. clausae.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex. *Flor.* subracemosa. *Fulcr.* o.

CINERARIA. *Rec.* nudum. *Papp.* simplex. *Cal.* simplex, polyphyllus, aequalis.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex. *Flor.* corymbosa. *Fulcr.* bracteae.

INULA. *Rec.* nudum. *Papp.* simplex. *Antherae* basi in 2 fetas desinentes.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex, amplexicaulis. *Flor.* subpaniculata. *Fulcr.* o.

ARNICA. *Rec.* nudum. *Papp.* simplex. *Cor.* radii filamentis 5 absque antheris.

Rad. ramosa. *Caul.* herbacea, simplex. *Fol.* simplex, opposita. *Flor.* solitaria. *Fulcr.* o.

BELLIS. *Rec.* nudum, conicum. *Papp.* o. *Cal.* hemisphaer. squamis aequalibus. *Sem.* obovata.

Rad. fibrosa. *Caul.* scaposa. *Fol.* simplex, radicalis. *Flor.* solitaria. *Fulcr.* o.

CHRYSANTHEMUM. *Rec.* nudum. *Papp.* marginatus.
Cal. hemisphaericus imbricatus, squamis
 marginalibus membranaceis.

Rad. ramosa. *Caul.* herbacea. *Fol.* varia. *Flor.*
 terminalis ramorum. *Fulcr.* o.

MATRICARIA. *Recept.* nudum. *Papp.* o. *Cal.* hemi-
 sphaeric. imbricatus, squamis marginalibus
 acutiusculis solidis.

Rad. ramosa. *Caul.* herbacea. *Fol.* composita,
Flor. terminalis ramorum. *Fulcr.* o.

COTULA. *Rec.* subnudum. *Papp.* marginatus. *Cor.* disci
 4-fidae, radii fere nullae.

Rad. fibrosa. *Caul.* herbacea. *Fol.* simplex, am-
 plexicaulis. *Flor.* solitaria. *Fulcr.* o.

ANTHEMIS. *Rec.* paleaceum. *Papp.* o. *Cal.* hemisphaer.
 subaequal. *Flosculi* radii plures quam 5.

Rad. ramosa. *Caul.* herbacea. *Fol.* composita.
Flor. subpaniculata. *Fulcr.* o.

ACHILLEA. *Rec.* paleaceum. *Papp.* o. *Cal.* ovat. im-
 bricat. *Flosc.* radii circiter 4.

Rad. ramosa. *Caul.* herbacea. *Fol.* varia, den-
 ticulata. *Flor.* corymbosa. *Fulcr.* o.

3. POLYGAMIA FRUSTRANEA.

COREOPSIS. *Recept.* paleaceum. *Papp.* 2-cornis. *Cal.*
 erectus, polyphyllus basi radiis patentibus
 cinctus.

Rad. fibrosa. *Caul.* herbacea. *Fol.* simplex, op-
 posita. *Flor.* terminalis. *Fulcr.* folia floralia.

CENTAUREA. *Rec.* fetosum. *Papp.* simplex. *Cor.* radii
 infundibulif. longiores, irregulares.

Rad. ramosa. *Caul.* herbacea. *Fol.* varia. *Flor.*
 varia. *Fulcr.* o.

4. POLYGAMIA NECESSARIA.

CALENDULA. *Rec.* nudum. *Papp.* o. *Cal.* polyphyll. aequal. *Sem.* disci membranacea.

Rad. fusiformis. *Caul.* herbacea. *Fol.* simplex. *Flor.* pedunculata. *Fulcr.* o.

FILAGO. *Rec.* nudum. *Papp.* o. *Cal.* imbricat. *Floft.* *Foemin* inter squamas locati.

Rad. fibrosa. *Caul.* herbacea. *Fol.* simplex. *Flor.* subpaniculata. *Fulcr.* o.

4. MONOGAMIA.

JASIONE. *Cal.* commun. 10-phyll. *proprius* 5-dent. *Cor.* 5-pet. regulares. *Capsf.* inferae, 2-loc.

Rad. fusiformis. *Caul.* herbacea, simplex. *Fol.* simplex, subopposita. *Flor.* terminalis, solitaria, capitata. *Fulcr.* o.

LOBELIA. *Cal.* 5-fid. *Cor.* 1-pet. irregul. *Capsf.* infera, 2-locul.

Rad. fibrosa. *Caul.* scaposa. *Fol.* radicalis, simplex, 2-loc. *Flor.* racemosa. *Fulcr.* bracteae.

VIOLA. *Cal.* 5-phyll. *Cor.* 5-pet. irregul. postice cornuta. *Capsf.* supera 3-valv. 1-locul.

Rad. fibrosa. *Caul.* herbacea. *Fol.* subcordata. *Flor.* pedunculata, nutans. *Fulcr.* stipulae.

IMPATIENS. *Cal.* 2-phyll. minim. *Cor.* 5-pet. irregul. *nectario* cucullato. *Capsf.* supera, 5-valv. elastica.

Rad. ramosa. *Caul.* herbacea, succulenta. *Fol.* simplex. *Flor.* axillaris, pedunculata. *Fulcr.* o.

XIX. SYNGENESIA.

I. POLYGAMIA AEQUALIS.

TRAGOPOGON.

935. *T. pratense*, calycinis corollae radium aequantibus, foliis integris strictis. Fl. Sv. 684. Fl. N. 649. Fl. D. 906. B. pr.

† 936. *T. porrifolium*, calycibus corollae radio longioribus, foliis integris strictis, pedunculis superne incrassatis. S. N. 710. Fl. D. 797. B.

SCORZONERA.

937. *S. humilis*, caule subnudo unifloro, foliis lato-lanceolatis nervosis planis. Fl. Sv. 685. Fl. D. 816. P. pr.

† 938. *S. purpurea*, foliis lineari-subulatis integris canaliculato-triquetris, pedunculis cylindricis. S. N. 711. Jacqu. Fl. Austr. l. t. 35. 2). P.

PICRIS.

939. *P. Echioides*, perianthiis exterioribus pentaphyllis, interiore aristato majoribus. S. N. 711. A. agr.

940. *P. Hieracioides*, perianthiis laxis, foliis integris, pedunculis squamatis in calycem. Fl. Sv. 686. A. agr.

APARGIA.

941. *A. autumnalis*, caule ramoso, pedunculis squamatis, foliis lanceolatis dentatis glabris. *Leontodon autumnale* S. N. 715. Fl. Sv. 695. Obs. Bot. F. I. p. 89. Fl. N. 287. a). P. pr.

M 5

942. *A.*

z) *Cel. Kölpin.*

a) T. 501. Fl. Dan. hujus nomen gerit, sed aliam omnino plantam refert et quidem *Crepidem testorum* ad quem quodque iconem citavit ill. a Linné. Pappus sessilis appictus neminem fallere debet.

186 SYNGENESIA. POLYGAM. AEQUALIS. Sonchus.

942. *A. hispida*, foliis lanceolatis dentatis hispidis: setis furcatis. *Leontodon hispidum* S. N. 715. Fl. Sv. 694. Fl. D. 864. P. pr.

SONCHUS.

943. *S. maritimus*, pedunculo nudo, foliis lanceolatis amplexicaulibus indivisis retrorsum argute dentatis. S. N. 712. All. Fl. Pedem. 818. t. 16. f. 2. b). P.
 † 944. *S. palustris*, pedunculis calycibusque hispidis subumbellatis, foliis runcinatis basi aristatis. S. N. 712. Weig. Suppl. 890. P.
 945. *S. arvensis*, pedunculis calycibusque hispidis subumbellatis, foliis runcinatis basi cordatis. Fl. Sv. 687. Fl. N. 135. Fl. D. 606. P. agr.
 946. *S. oleraceus*, pedunculis tomentosis, calycibus glabris. Fl. Sv. 688. Fl. N. 140. A. c.
 α . *laevis*. Fl. D. 682.
 β . *asper*. Fl. D. 843.
 947. *S. lapponicus*, pedunculis squamosis, floribus racemosis, foliis runcinatis. *S. alpinus* Fl. Sv. 689. Smith. icon. I. t. 21. A. alp.
 948. *S. alpinus*, pedunculis bracteisque longioribus hirsutis, floribus racemosis, foliis runcinatis basi sagittatis. *Frölich* Usteri Ann. I. p. 20. Fl. N. 52. Fl. D. 182. P. alp.
 949. *S. Sibiricus*, pedunculis squamatis, foliis lanceolatis indivisis sessilibus. Fl. Sv. 690. Gmel. Fl. Sib. II. t. 3. P.

LACTUCA.

950. *L. quercina*, foliis runcinatis denticulatis acutis subtus laevibus, caule glabro. Fl. Sv. 691. P.
 951. *L. virofa*, foliis horizontalibus carina aculeatis dentatis. S. N. 713. Weig. Fl. 492. A. r.

CHON-

b) *Cel. Afzelius*.

CHONDRILLA.

- †952. *C. juncea*, foliis radicalibus runcinatis, caulinis linearibus integris. S. N. 713. Jacqu. Fl. Austr. V. t. 427. c). P.

PRENANTHES.

953. *P. muralis*, flosculis quinis, foliis runcinatis. Fl. Sv. 692. Fl. N. 945. Fl. D. 509. P. sylv.

LEONTODON.

954. *L. Taraxacum*, calyce squamis inferne reflexis, foliis runcinatis denticulatis laevibus. Fl. Sv. 693. Fl. N. 286. Fl. D. 574. P. c. pr. pasc.
 β. *palustre*, foliis integris lineari-lanceolatis. it. Oel. p. 62. d).

HIERACIUM.

955. *H. alpinum*, foliis oblongis integris dentatis pilosis, caule subnudo unifloro, calyce piloso. Fl. Sv. 696. Fl. N. 592. P. alp.
 α. *grandiflorum*, caule vix folioso, flore magno. Rarij Angl. 3. t. 6. f. 2.
 β. *parviflorum*, caule folioso altiore, flore minore. Fl. D. 27.
956. *H. Taraxaci*, foliis lanceolatis dentatis glabris, scapo subnudo unifloro, calyce hirsuto. S. N. 716. Obs. Bot. Fasc. IV. tab. 2. P. alp.
957. *H. Pilosella*, foliis integerrimis ovatis subtus tomentosis, stolonibus repentibus, scapo unifloro. Fl. Sv. 697. Fl. N. 283. P. pasc. pr. sicc.
958. *H. dubium*, foliis integris ovato-oblongis, stolonibus repentibus, scapo nudo multifloro. Fl. Sv. 698. P. pr.

959. *H.*

c) *Cl. Helwig.*

d) Huc pertinere videtur *Scorzonera Taraxaci* Roth. Beytr. 1. Th. p. 120. Folia licet runcinato-dentata sint.

188 SYNGENESIA. POLYGAMIA AEQUALIS. Crepis.

959. *H. Auricula*, foliis integerrimis lanceolatis, stolonibus repentibus, scapo nudo multifloro. Fl. Sv. 699. P. pr. pasc.
960. *H. cymosum*, foliis lanceolatis integris pilosis, scapo subnudo basi piloso, floribus subumbellatis. S. N. 716. Fl. D. 810. P.
961. *H. praemorsum*, foliis ovatis subdentatis, scapo nudo ramoso, floribus superioribus primoribus. Fl. Sv. 700. Fl. D. 942. P. pr. ficc.
- † 962. *H. aurantiacum*, foliis integris, caule subnudo simplicissimo piloso corymbifero. S. N. 716. Jacqu. Fl. Austr. V. t. 410. e). P.
963. *H. murorum*, caule ramoso, foliis radicalibus ovatis dentatis: caulino minore. Fl. Sv. 701. Fl. N. 396. P. pr. sylv.
- β. *sylvaticum*. Obs. Bot. Fasc. I. p. 90.
- γ. *laciniatum*, foliis ad basin longe dentatis.
- δ. *myophorum*, cynipis Hieracii.
964. *H. paludosum*, caule paniculato, foliis amplexicaulibus dentatis glabris, calycibus hispidis. Fl. Sv. 702. Fl. N. 566. Fl. D. 928. P. pr. n. hum.
965. *H. Sabaudum*, caule erecto multifloro, foliis ovato-lanceolatis dentatis semiamplexicaulibus. Fl. Sv. 703. Fl. N. 699. Fl. D. 872. P. pr.
966. *H. umbellatum*, foliis linearibus subdentatis sparsis, floribus subumbellatis. Fl. Sv. 704. Fl. N. 138. Fl. D. 680. P. pr. ficc. a.

CREPIS.

967. *C. tectorum*, foliis lanceolato-runcinatis sessilibus laevibus, inferioribus dentatis. Fl. Sv. 705. Fl. N. 525. *Leontodon autumnale* Fl. D. 501. A. pr. arv. tect.

a) Rever. Ström.

968. C.

SYNGENESIA. POLYGAM. AEQUALIS. Hyoseris. 189.

968. *C. biennis*, foliis runcinato-pinnatifidis scabris: basi superne dentatis, calycibus muricatis. Fl. Sv. 706. B. pr. r.

HYOSERIS.

969. *H. minima*, caule diviso nudo, pedunculis incrassatis. Fl. Sv. 707. Fl. D. 201. A. agr. arv. a.

HYPOCHAERIS.

970. *H. maculata*, caule subnudo, ramo solitario, foliis ovato-oblongis integris dentatis. Fl. Sv. 708. Fl. N. 397. Fl. D. 149. P. pr.
971. *H. glabra*, calycibus oblongis imbricatis, caule ramoso nudo, foliis dentato-sinuatis. S. N. 721. Fl. D. 424. Weig. Fl. p. 146. A. agr. a.
972. *H. radicata*, foliis runcinatis obtusis scabris, caule ramoso nudo laevi, pedunculis squamosis. Fl. Sv. 709. Fl. N. 398. Fl. D. 150. P. pr.

LAPSANA.

973. *L. communis*, calycibus fructus angulatis, pedunculis tenuibus ramosissimis. Fl. Sv. 710. Fl. N. 139. Fl. D. 500. A. c.

CICHORIUM.

974. *C. Intybus*, floribus geminis sessilibus, foliis runcinatis. Fl. Sv. 711. Fl. D. 907. P. pr. v. r. agr.

ARCTIUM.

975. *A. Lappa*, foliis cordatis inermibus petiolatis. Fl. Sv. 712. Fl. N. 258. B. P. r. v.
 α. capitulis glabris. *vulgaris*.
 β. capitulis tomentosis. Fl. D. 642. *arachnoidea*.

SERRATULA.

976. *S. tinctoria*, foliis lyrato-pinnatifidis: pinna terminali maxima, flosculis conformibus. Fl. Sv. 713. Fl. D. 281. P. pr.

190 SYNGENESIA. POLYGAM. AEQUALIS. *Carduus*.

977. *S. alpina*, calycibus subhirsutis ovatis, foliis indivisis. Fl. Sv. 714. Fl. N. 48. Fl. D. 37. P.
 978. *S. arvensis*, foliis dentatis spinosis. Fl. Sv. 715. Fl. N. 255. Fl. D. 644. P. agr.
 α . *glabra*. Obs. Bot. F. I. 91.
 β . *tomentosa*. l. cit.

CARDUUS.

979. *C. nutans*, foliis semidecurrentibus spinosis, floribus cernuis: squamis calycinis superna patentibus. Fl. Sv. 717. Fl. N. 565. Fl. D. 675. B. r.
 980. *C. Acanthoides*, foliis decurrentibus sinuatis margine spinosis, calycibus pedunculatis solitariis erectis villosis. Fl. Sv. 718. Jacq. Fl. Austr. III. t. 249. A. r. coem.
 981. *C. crispus*, foliis decurrentibus sinuatis margine spinosis, floribus aggregatis terminalibus, squamis inermibus subaristatis patulis. Fl. Sv. 719. Fl. N. 256. Fl. D. 621. A. c.

CNICUS.

982. *C. acaulis*, acaulis, calyce glabro. *Carduus acaulis* Fl. Sv. 722. P. pasc. pr.
 983. *C. lanceolatus*, foliis decurrentibus pinnatifidis hispidis: laciniis divaricatis, calycibus ovatis spinosis villosis, caule piloso. *Carduus lanceolatus*. Fl. Sv. 716. Fl. N. 720. f). B. arv. v. r.
 984. *C. palustris*, foliis decurrentibus dentatis margine spinosis, floribus racemosis erectis, pedunculis inermibus. *Carduus palustris*. Fl. Sv. 720. Fl. N. 598. P. pr. ul.

985. *C.*

f) Praeantibus *Haller*, *Scopoli*, *Allione*, *Schreber* Carduos pappo plumoso ad Cnici genus retuli, et nisi hoc adsumatur, nullus omnino character genera Cnici et Cardui distinguens. Bractaeae enim haud tutum characterem offerunt. Nec Cirsii genere opus est.

SYNGENESIA. POLYG. AEQUALIS. Onopordum. 191

985. *C. heterophyllus*, foliis amplexicaulibus lanceolatis ciliatis integris laciniatisque, caule subuni-
floro, calyce inermi. *Carduus heterophyllus*. Fl. Sv. 721.
St. Helv. t. 7. P. pr. fylv.
986. *C. helenioides*, foliis amplexicaulibus lanceolatis
dentatis: spinulis inaequalibus ciliatis, caule inermi.
Carduus helenioides. S. N. 726. g). *Carduus hetero-*
phyllus. Fl. N. 257. Fl. D. 109. P. pr. hum.
987. *C. oleraceus*, foliis pinnatifidis carinatis nudis,
bracteis concavis integris subcoloratis. Fl. Sv. 723.
Fl. D. 860. P. pr. n.

ONOPORDUM.

988. *O. Acanthium*, calycibus squarrosis: squamis pa-
tentibus, foliis ovato-oblongis sinuatis. Fl. Sv. 724.
Fl. D. 909. B. r.

CARLINA.

989. *C. vulgaris*, caule multifloro corymbofo, floribus
terminalibus, calycis radio albo. Fl. Sv. 725.
Fl. N. 966. B. pasc.

BIDENS.

990. *B. tripartita*, foliis trifidis, calycibus subfoliosis,
feminibus erectis. Fl. Sv. 726. Fl. N. 524. A. aqu.
β. *pumila*, foliis serratis, caule simplicissimo
uni-
floro. Obs. Bot. Fasc. I. 94.
991. *B. minima*, foliis lanceolatis sessilibus, floribus se-
minibusque erectis. S. N. 732. Obs. Bot. Fasc. I. 94.
Fl. D. 312. A. aqu.
992. *B. cernua*, foliis lanceolatis amplexicaulibus, flo-
ribus cernuis. Fl. Sv. 727. Fl. D. 841. A. aqu.
β. *radiata*, quae certe eadem est c. *Corropside*
Bidente.

993. *B.*

g) An revera diversus a heterophyllo?

192 SYNGENESIS. POLYG. AEQUALIS. *Eupatorium*.

993. *B. frondosa*, foliis pinnatis ferratis lineatis glabris, feminibus erectis, calycibus frondosis, caule laevi. S. N. 732. h). A. hum.

EUPATORIUM.

994. *E. cannabinum*, foliis digitatis. Fl. Sv. 728. Fl. D. 745. P. n. riv.

CHRYSOCOMA.

995. *C. Lynosyris*, herbacea, foliis linearibus glabris, calycibus laxis. Fl. Sv. 729. P.

2. POLYGAMIA SUPERFLUA.

TANACETUM.

996. *T. vulgare*, foliis bipinnatis incisis ferratis. Fl. Sv. 730. Fl. N. 68. Fl. D. 871. P. v. f.

ARTEMISIA.

997. *C. campestris*, foliis multifidis linearibus, caulibus procumbentibus virgatis. Fl. Sv. 732. Fl. N. 803. S. a.

α. *alba*, caulibus viridi-albis.

β. *rubra*, caulibus rubris.

998. *A. maritima*, foliis multipartitis tomentosis, racemis cernuis, flosculis foemineis ternis. Fl. Sv. 734. Fl. N. 801. S. l. m.

999. *A. rupestris*, foliis pinnatis, caulibus adscendentibus, floribus globosis cernuis, receptaculo paposo. Fl. Sv. 733. Lt. Gottl. t. 285. Fl. D. 801. i). P. ficc.

1000. *A. Absinthium*, foliis compositis multifidis, floribus subglobosis pendulis, receptaculo villosa. Fl. Sv. 735. Fl. N. 1081. P. r. coem.

1001. *A.*

h) *Cel. Afzelius*. Conf. Obs. Bot. Fasc. I. p. 93.

i) Omnino diversa ab *A. rupestris* Scop. Carn. ed. 2. 1038.

1001. *A. vulgaris*, foliis pinnatifidis planis incisfis subtus tomentosis, racemis simplicibus recurvatis, floribus radio quinquefloro. Fl. Sv. 731. Fl. N. 69. P. r. f. c.

α. alba, caulibus virescentibus.

β. rubra, caulibus rufis.

GNAPHALIUM.

1002. *G. arenarium*, herbaceum, foliis lanceolatis: inferioribus obtusis, corymbo composito, caulibus simplicissimis. Fl. Sv. 738. Fl. D. 641. B. a.

1003. *G. dioicum*, sarmentis procumbentibus, caule simplicissimo, corymbo simplici, floribus dioicis. Fl. Sv. 736. Fl. N. 632. P. pasc. pr.

α. mas, flore globofo.

β. foemina, flore oblongo.

Variat flore albo et rubro.

1004. *G. alpinum*, sarmentis procumbentibus, caule simplicissimo, capitulo aphylo, floribus oblongis. Fl. Sv. 737. Fl. N. 543. Fl. D. 332. k). P. alp.

1005. *G. sylvaticum*, caule herbaceo simplicissimo erecto, foliis linearibus, floribus sparsis. Fl. Sv. 739. Fl. N. 227. A. sylv.

- † 1006. *G. Norvegicum*, caule herbaceo simplicissimo erecto, foliis lanceolatis basi angustatis, floribus sparsis, calycibus acutis nigricantibus. Fl. D. 254. Fl. N. 841. Fl. Isl. 9. l). P. alp.

1007. *G. uliginosum*, caule herbaceo ramofo diffuso floribus confertis terminalibus. Fl. Sv. 740. Fl. N. 339. Fl. D. 859. A. in.

CONYZA.

k) Nonnisi flores foemineos vidit *Linne*, duplicis generis pinxit *Oeder*, paucos foemineos, androgynos plures observavit *Haller*, omnes hermaphroditos pronunciavit *Scopoli*. Quid ergo statuendum? An polygama trioica planta?

l) Magis a *sylvatico* differre videtur quam *supinum*.

194 SYNGENESIA. POLYG. SUPERFLUA. Conyza.

CONYZA.

- †1008. *C. squarrosa*, foliis lanceolatis acutis, caule herbaceo corymbofo, calycibus squarrosis. S. N. 752. Fl. D. 622. Weig. Fl. 531. B.

ERIGERON.

1009. *E. acre*, pedunculis alternis unifloris. Fl. Sv. 741. Fl. N. 504. P. pasc. v.
 †1010. *E. Dröbachtense*, caule floribusque pedunculato, terminali majore. Fl. D. 874. m). l. m.
 1011. *E. uniflorum*, caule unifloro, calyce piloso. Fl. Sv. 742. Fl. L. t. 9. f. 3. Fl. N. 665. Fl. D. 292. n). P. alp.

TUSSILAGO.

1012. *T. Farfara*, scapo unifloro imbricato, foliis subcordatis angulatis denticulatis. Fl. Sv. 743. Fl. N. 169. Fl. D. 595. P. arg. hum.
 1013. *T. frigida*, thyrsio fastigiato, floribus radiatis. Fl. Sv. 744. Fl. N. 81. Fl. D. 61. P. alp.
 1014. *T. alba*, thyrsio fastigiato, flosculis foemineis nudis paucis, foliis subcordato rotundis dentatis. Fl. Sv. 745. Obs. Bot. Fasc. I. p. 97. Fl. D. 524. P. umbr. hum.
 1015. *T. spuria*, thyrsio fastigiato, flosculis foemineis subpetaloideis plurimis, foliis triangulari cordatis denticulatis: lobis sinuatis. Prodr. Fl. Scand. 910. Obs. Bot. Fasc. I. 98. t. 2. P. l. lac. a.
 1016. *T. Petasites*, thyrsio ovato, flosculis foemineis nudis paucis (etiam nullis). Fl. Sv. 746. Fl. D. 842. P. pag.

SENECIO.

- m) Num nova species vel potius varietas *E. acris*? *E. canadense* diversum ab hoc viderur.
 n) Icon Fl. Dan. varietatem fittit grandifloram.

SENECIO.

1017. *S. vulgaris*, corollis nudis, foliis pinnato-finuatis amplexicaulibus, floribus sparsis. Fl. Sv. 747. Fl. N. 284. Fl. D. 513. A. c. r.

1018. *S. viscosus*, corollis revolutis, foliis pinnatifidis viscidis, squamis calycinis laxis longitudine perianthii. Fl. Sv. 749. A. r.

β. *S. denticulatus* Fl. D. t. 791. ?

1019. *S. sylvaticus*, corollis revolutis, foliis pinnatifidis denticulatis, caule corymbofo erecto. Fl. Sv. 748. Fl. N. 949. Fl. D. 869. A. r. sylv.

1020. *S. Jacobaea*, corollis radiantibus, foliis pinnato-lyratis: laciniis lacinulatis, caule erecto. Fl. Sv. 751. Fl. N. 950. Fl. D. 944. P. pr. v.

β. foliis non lacinulatis.

1021. *S. aquaticus*, corollis radiantibus, foliis dentatis: radicalibus ovatis, caulinis pinnatifidis: lacinia extrema majore. Hudf. Fl. Angl. 2. p. 366. Fl. D. 784. *S. erucifolius* Weig. Suppl. 895 o). P. pr. hum.

1022. *S. paludosus*, corollis radiantibus, foliis ensiformibus acute serratis subtus subvillosis, caule stricto. Fl. Sv. 752. Fl. D. 385. B. aqu.

ASTER.

1023. *A. Tripolium*, foliis lanceolatis integerrimis carnosif glabris, ramis inaequatis, floribus corymbosis. Fl. Sv. 753. Fl. N. 92. Fl. D. 615. P. l. m.

† 1024. *A. annuus*, foliis lanceolatis: inferioribus crenatis, caule corymbofo, pedunculis nudis, calycibus hemisphaericis. S. N. 762. Fl. D. 486. A.

N 2

SOLI-

*) Ad Parfile juxta Gothoburgum copiose. Forte *Jacobaea III laxifolia* I. Clus. Hist. 2. p. XXIII. quae eadem cum IV. videtur.

196 SYNGENESIA. POLYG. SUPERFLUA. Solidago.

SOLIDAGO.

1025. *S. Virgaurea*, caule subflexuoso angulato, racemis paniculatis erectis confertis. Fl. Sv. 754. Fl. N. 240. Fl. D. 663. P. pr.

CINERARIA.

1026. *C. palustris*, floribus corymbosis, foliis lato-lanceolatis dentato-finuatis, caule villoso. S. N. 765. Fl. D. 573. *Othonna palustris* Fl. Sv. 778. P. pal. pr. hum.
1027. *C. campestris*, floribus corymbosis, foliis radicalibus petiolatis ovatis tomentosis: caulinis sessilibus linearibus. Prodr. Fl. Scand. 922. Obs. Bot. F. I. 101. *Cin. alpina* γ. Spec. Plant. 1243. *Cin. integrifolia* β. S. N. 765. Gmel. Sib. 2. t. 71. f. A. *Jacob. altera*. Clus. Hist. II. p. 23. p). Jacqu. Fl. Austr. II. t. 180. apr.

INULA.

1028. *I. Helenium*, foliis amplexicaulibus ovatis rugosis subtus tomentosis calycum squamis ovatis. Fl. Sv. 755. Fl. D. 728. P. c.
1029. *I. Britannica*, foliis amplexicaulibus lanceolatis ferratis distinctis subtus villosis, caule ramoso erecto villoso. Fl. Sv. 756. Fl. D. 413. P. pr. hum.
1030. *I. dysenterica*, foliis amplexicaulibus cordato-oblongis, caule villoso paniculato, squamis calycinis fetaceis. Fl. Sv. 757. Fl. D. 410. P. r. hum.
1031. *I. Pulicaria*, foliis amplexicaulibus undulatis, caule prostrato, floribus subglobosis radio brevissimo. Fl. Sv. 758. Fl. D. 613. A. r. v.

1032. *I.*

p) Verum est iconem Clusianam multum ad nostrum accedere plantam, sed nostra minor est, nec quantum scio perennis, quia in Hortum introducta, absoluto semine periit, nec omni anno in locis ubi provenit observatur, quare biennem credo.

SYNGENESIA. POLYGAM. SUPERFLUA. Arnica. 197

1032. *I. falicina*, foliis lanceolatis recurvis ferrato-sca-
bris, ramis angulatis, floribus inferioribus altio-
ribus. Fl. Sv. 759. Fl. D. 786. P. pr. hum.

† 1033. *I. hirta*, foliis sessilibus lanceolatis recurvis sub-
ferrato-scabris, caule teretiufculo subpiloso, flo-
ribus inferioribus altioribus. S. N. 767. Jacqu. Fl.
Austr. IV. t. 358. g). P.

ARNICA.

1034. *A. montana*, foliis integris: caulinis geminis op-
positis. Fl. Sv. 760. Fl. N. 416. P.

α. *pratensis*, foliis ovatis. Fl. D. 63. pr.

β. *alpina*, foliis lanceolatis. Fl. L. 305. alp.

BELLIS.

1035. *B. perennis*, scapo nudo. Fl. Sv. 761. Fl. N. 1112.
Fl. D. 503. P. pasc. arg.

CHRYSANthemum.

1036. *C. segetum*, foliis amplexicaulibus: superne laci-
niatis, inferne dentato-ferratis. Fl. Sv. 762.
Fl. N. 622. Fl. D. 995. A. agr.

1037. *C. Leucanthemum*, foliis amplexicaulibus oblon-
gis: superne ferratis, inferne dentatis. Fl. Sv. 763.
Fl. N. 219. Fl. D. 994. P. pr.

1038. *C. inodorum*, foliis pinnatis multifidis, caule ra-
moso diffuso. S. N. 772. Fl. N. 634. Fl. D. 696.
Matricaria inodora. Fl. Sv. 765. A. r. c.

MATRICARIA.

1039. *M. Parthenium*, foliis compositis planis: foliolis
ovatis incis, pedunculis ramosis. S. N. 774.
A. Holm. 1769. p. 247. Weig. Fl. 548. Fl. D. 674.
P. r.

N 3

1040. *M.*

g) Cl. Hellwig.

198 SYNGENESIA. POLYGAM. SUPERFLUA. *Cotula*.

1040. *M. maritima*, receptaculis hemisphaericis, foliis bipinnatis subcarnosis supra convexis subtus carinatis. S. N. 774. Raji Angl. 3. t. 7. f. 1. N. Ad. Holm. 1787. p. 250. P. lit. m. mont.
1041. *M. suaveolens*, receptaculis conicis, radiis deflexis, squamis calycinis margine aequalibus. Fl. Sv. 766. r). A. c.
1042. *M. Chamomilla*, receptaculis conicis, radiis patentibus, squamis calycinis margine aequalibus. Fl. Sv. 764. Fl. N. 400. A. c.

COTULA.

- † 1043. *C. coronopifolia*, foliis lanceolato linearibus amplexicaulibus dentatis, floribus flosculosis. S. N. 774. Fl. D. 741. A. v. in.

ANTHEMIS.

1044. *A. maritima*, foliis pinnatis denticulatis carnosius nudis punctatis, caule prostrato, calycibus subtomentosis. S. N. 776. P. l. m.
1045. *A. arvensis*, receptaculis conicis: paleis setaceis, seminibus coronato-marginatis. Fl. Sv. 768. Fl. N. 589. A. c.
1046. *A. Cotula*, receptaculis conicis: paleis setaceis, seminibus nudis. Fl. Sv. 767. Fl. N. 220. A. r.
1047. *A. tinctoria*, foliis bipinnatis serratis subtus tomentosis, caule corymboso. Fl. Sv. 769. Fl. N. 534. Fl. D. 741. P. pr. agr.

ACHILLEA.

1048. *A. Ptarmica*, foliis lanceolatis acuminatis argute serratis. Fl. Sv. 771. Fl. N. 491. Fl. D. 643. P. pr. hum.
1049. *A. Millefolium*, foliis bipinnatis nudis: laciniis linearibus dentatis, caulibus superne sulcatis. Fl. Sv. 770. Fl. N. 194. Fl. D. 737. P. pasc. pr. f. 1050. A.

r) A sequente haud diversam judicat *Cel. Afzelius*.

SYNGENESIA, POL. FRUSTRANEA. Coreopsis. 199

1050. *A. nobilis*, foliis bipinnatis, inferioribus nudis planis, superioribus obtusis tomentosis, corymbis convexis confertissimis. S. N. 778. Obs. Bot. F. I. p. 87. Ros. Obs. 48. N. Act. Holm. 1787. p. 252. P. pr. n.

3. POLYGAMIA FRUSTRANEA.

COREOPSIS.

1051. *C. Bidens*, foliis lanceolatis ferratis oppositis amplexicaulibus. Fl. Sv. 772. 1). A. aqu.

CENTAUREA.

1052. *C. phrygia*, calycibus recurvato-plumosis, foliis indivisis oblongis scabris. Fl. Sv. 775. Fl. D. 520. P.
1053. *C. nigra*, calycibus ciliatis: squamula ovata: ciliis capillaribus erectis, foliis lyrato-angulatis, floribus flosculosis. S. N. 785. Fl. D. 996. P. pr.
1054. *C. Cyanus*, calycibus ferratis, foliis linearibus integerrimis: infimis dentatis. Fl. Sv. 776. Fl. N. 303. Fl. D. 993. A. agr.
1055. *C. Stabiosa*, calycibus ciliatis, foliis pinnatifidis: pinnis lanceolatis. Fl. Sv. 773. Fl. N. 510. P. pr. agr.
1056. *C. Jacea*, calycibus scariosis laceris, foliis lanceolatis: radicalibus sinuato-dentatis, ramis angulatis. Fl. Sv. 774. Fl. N. 368. Fl. D. 519. P. pr. agr.

4. POLYGAMIA NECESSARIA.

CALENDULA.

1057. *C. arvensis*, seminibus cymbiformibus muricatis incurvatis: extimis erectis protensis. Fl. Sv. 777. A. c. r.

N 4

FILA-

1) Sine dubio eadem cum *Bidente cernua* β , at heic repetita, ne incautum decipiet.

FILAGO.

1058. *F. germanica*, panicula dichotoma, floribus rotundatis, axillaribus hirsutis, foliis acutis. S. N. 795. Weig. Fl. 559. Fl. D. 997. *F. pyramidata* Fl. Sv. 779. Fl. N. 933. A. c.
 1059. *F. montana*, caule subdichotomo erecto, floribus conicis terminalibus axillaribusque. Fl. Sv. 780. A. a. ficc.
 1060. *F. arvensis*, caule paniculato. floribus conicis lateralibus. Fl. Sv. 781. A. ficc. a.

5. MONOGAMIA.

JASIONE.

1061. *J. montana*. Fl. Sv. 782. Fl. N. 823. Fl. D. 319. f). A. a. ficc.

LOBELIA.

1062. *L. Dortmanna*, foliis linearibus bilocularibus integerrimis, caule subnudo. Fl. Sv. 783. Fl. N. 522. Fl. D. 39. P. l. lac.

VIOLA.

1063. *V. hirta*, acaulis, foliis cordatis piloso-hispidis.
 • Fl. Sv. 788. Fl. N. 619. Fl. D. 618. P. pr. n.
 1064. *V. palustris*, acaulis, foliis reniformibus. Fl. Sv. 786. Fl. N. 170. Fl. D. 83. P. pr. hum.
 1065. *V. odorata*, acaulis, foliis cordatis, stolonibus reptantibus. Fl. Sv. 784. Fl. D. 309. P. pr. hort.
 1066. *V. canina*, caule adultiore adscendente, foliis oblongo-cordatis. Fl. Sv. 785. Fl. N. 172. P. pr. n.
 1067. *V. montana*, caulibus erectis, foliis cordatis oblongis. Fl. Sv. 787. Fl. N. 503. P. alp.
 1068. *V.*

*) Polygama Monoica observante Nob. Jacquin in Flor. Vindob.

1068. *V. mirabilis*, caule triquetro, foliis reniformi-cordatis, floribus caulinis apetalis. Fl. Sv. 789. Jacqu. Fl. Austr. I. t. 19. u). P. pasc. sylv.
 1069. *V. biflora*, caule bifloro, foliis reniformibus serratis. Fl. Sv. 790. Fl. N. 131. Fl. D. 46. P. alp.
 1070. *V. tricolor*, caule triquetro diffuso, foliis oblongis incisfis, stipulis pinnatifidis. Fl. Sv. 791. Fl. N. 171. Fl. D. 623. A. c.
β. arvensis, calyce corollam bicolorum parum excedente.

IMPATIENS.

1071. *I. noli tangere*, pedunculis multifloris solitariis, foliis ovatis, geniculis caulinis tumentibus. Fl. Sv. 792. Fl. N. 134. Fl. D. 582. w). A. umbr. riv.

u) In caulibus etiam saepius flores petalodeos observavit Cel. Vahl.

w) In ollis transpositae ex solo natali plantae florent fructum-que ferunt sine petalis.

XX. GYNANDRIA.

I. DIANDRIA.

ORCHIS. *Nectar.* corniforme pone florem.

Rad. carnosae geminatae vel fasciculatae. *Caul.* herbacea, simplicissima. *Fol.* simplex, vaginans. *Flor.* spicata. *Fulcr.* bracteae.

SATYRIUM. *Nectar.* scrotiforme vel inflato-didymum pone florem.

Rad. etc. Orchidis.

OPHRYS. *Nectar.* subtus subcarinatum.

Rad. etc. priorum.

SERAPIAS. *Nectar.* ovatum, gibbum, labio ovato.

Rad. etc. antecedentium.

CYPRIPEDIUM. *Nectarium* ventricosum, inflatum, cavum.

Rad. etc. congenerum, sed flores subsolitarii.

2. HEXANDRIA.

ARISTOLOCHIA. 6-gyna. *Cal.* o. *Cor.* 1-pet. linguata, integra. *Capsf.* 6-locul. infera.

Rad. fibrosa, repens. *Caul.* herbacea, flexuosa. *Fol.* simplex, caulina. *Flor.* axillaris. *Fulcr.* o.

3. POLYANDRIA.

ARUM. *Spatha* 1-phylla, cucullata. *Spadix* supra nudus, inferne foemineus, medio staminifer.

Rad. globosa. *Caul.* scaposa. *Fol.* simplex, radicalis. *Flor.* in spadice. *Fulcr.* o.

CALLA.

CALLA. *Spatha* 1-phylla, plana. *Spadix* tectus flosculis. *Cal.* o. *Cor.* o. *Bactae* polyspermae.
Rad. horizontalis, repens, reliqua *Ari.*

ZOSTERA. *Spadix* linearis altero latere fructifero.
Cor. o. *Cal.* o. *Stam.* alterna. *Sem.* solitaria, alterna.

Rad. fibrosa. *Cañl.* articulata, linearis, submersa. *Fol.* linearis, submersa. *Flor.* intra sinum folii. *Fulcr.* stipulae vaginantes.

XX. GYNANDRIA.

3. D I A N D R I A.

ORCHIS.

* *testiculatae.*

1072. *O. bifolia*, bulbis indivisis, nectarii labio lanceolato integerrimo: cornu longissimo, petalis patentibus. Fl. Sv. 793. Obs. Bot. F. I. 102. Fl. N. 32. Fl. D. 235. St. Helv. t. 35. P. pr.
1073. *O. pyramidalis*, bulbis indivisis, nectarii labio bicorni trifido aequali integerrimo, cornu longo, petalis subanceolatis. Fl. Sv. 798. St. Helv. t. 35. Jacqu. Fl. Austr. III. t. 226. P. pr.
1074. *O. Morio*, bulbis indivisis, nectarii labio quadrifido crenulato: cornu obtuso adscendente, petalis obtusis conniventibus. Fl. Sv. 794. Obs. Bot. F. I. p. 103. Fl. N. 745. Fl. D. 253. St. Helv. t. 33. P. pr.
1075. *O. mascula*, bulbis indivisis, nectarii labio quadrilobo crenulato: cornu obtuso, petalis dorsalibus reflexis. Fl. Sv. 795. Obs. Bot. Fasc. I. p. 104. Fl. 457. St. Helv. t. 33. P. pr. sylv.
1076. *O. ustulata*, bulbis indivisis, nectarii labio quadrifido punctis scabro, cornu obtuso, petalis distinctis. Fl. Sv. 797. Fl. D. 884. St. Helv. t. 28.
- Ophrys anthropophora*. Fl. D. 103. P. pr. n.
1077. *O. militaris*, bulbis indivisis, nectarii labio quinquefido punctis scabro: cornu obtuso, petalis confluentibus. Fl. Sv. 796. Fl. N. 694. St. Helv. t. 28. P. pr. n.
1078. *O. pallens*, bulbis indivisis, nectarii labio trifido integerrimo: cornu obtuso mediocri, petalis patentibus. S. N. 809. Mant. 2. p. 292. St. Helv. t. 30. Jacqu. Fl. Austr. I. t. 45. P. sylv.

** *palmatae.*

1079. *O. latifolia*, bulbis subpalmatis rectis, nectarii cornu conico: labio trilobo lateralibus reflexo, bracteis flore longioribus. Fl. Sv. 801. Fl. N. 656. Fl. D. 266. St. Helv. t. 32. P. pr.

1080. *O. mixta*, bulbis palmatis, nectarii cornu conico: labio trilobo subserrato, petalis dorsalibus reflexis. St. Helv. 1280. x). P. pr. subhum.

α. *O. incarnata*. Fl. Sv. 802, 803. β.

β. *O. sambucina*. Fl. Sv. 803. α.

1081. *O. maculata*, bulbis palmatis patentibus, nectarii cornu germinibus brevior: labio plano, petalis dorsalibus patulis. Fl. Sv. 800. Fl. N. 253. Fl. D. 933. St. Helv. t. 32. P. pr. hum.

1082. *O. odoratissima*, bulbis palmatis, nectarii cornu recurvo brevior, labio trilobo, foliis linearibus. Fn. Sv. 558. Fl. N. 696. St. Helv. t. 29. P. pr. sylv.

1083. *O. conopsea*, bulbis palmatis, nectarii cornu setaceo germinibus longior, labio trifido, petalis duobus patentissimis. Fl. Sv. 799. Fl. N. 707. Fl. D. 224. St. Helv. t. 29. P. pr. sylv.

† 1084. *O. cruenta*, bulbis palmatis, nectarii cornu germine brevior, labio indiviso subcordato crenato, petalis dorsalibus patulis. Fl. D. 876. P. pr. alp.

*** *fasciculatae.*

1085. *O. hyperborea*, bulbis fasciculatis, nectarii cornu longitudine germinis: labio lineari integerrimo truncato. S. N. 810. Fl. N. 828. Fl. Isl. 9. Obs. Bot. F. IV. t. 3. P. alp.

† 1086. 0.

* varietas β. pallide flava et praeter bulbos simillima *Or. pal-lenti*.

- † 1086. *O. abortiva*, bulbis fasciculatis filiformibus, nectarii labio ovato integerrimo, caule aphylo. S. N. 810. Fl. N. 695. Jacqu. Fl. Austr. II. t. 193. St. Helv. t. 36. P.
- † 1087. *O. Königii*, bulbis fibrosis, nectarii calcare germinibus brevior: labio tripartito, laciniis aequalibus integerrimis. Fl. N. 827. Fl. D. 333. P.

SATYRIUM.

1088. *S. viride*, bulbis palmatis, foliis oblongis obtusis, nectarii labio lineari trifido: lacinia intermedia obsoleta. Fl. Sv. 804. Fl. N. 278. Fl. D. 77. St. Helv. t. 26. P. pr.
1089. *S. nigrum*, bulbis palmatis, foliis linearibus, nectarii labio resupinato indiviso. Fl. Sv. 805. Fl. N. 445. Fl. D. 998. St. Helv. t. 27. P. pr.
1090. *S. albidum*, bulbis fasciculatis, foliis lanceolatis, nectarii labio trifido acuto: lacinia intermedia obtusa. Fl. Sv. 806. Fl. N. 446. Fl. D. 115. St. Helv. t. 26. P. pr.
1091. *S. repens*, bulbis fibrosis, foliis ovatis radicalibus, floribus secundis. Fl. Sv. 807. Fl. N. 321. t. 6. f. 1. Fl. D. 812. St. Helv. t. 22. P. pr. sylv.
β. *maculosum*.

OPHRYS.

1092. *O. Nidus Avis*, bulbis fibroso-fasciculatis, caule vaginato aphylo, nectarii labio bifido. Fl. Sv. 815. Fl. N. 963. Fl. D. 181. St. Helv. t. 37. P. sylv.
1093. *O. Corallorhiza*, bulbis ramosis flexuosis, caule vaginato aphylo, nectarii labio trifido. Fl. Sv. 816. Fl. N. 126. t. 6. f. 2. Fl. D. 451. P. sylv. hum.
- † 1094. *O. spiralis*, bulbis aggregatis oblongis, caule subfoliofo, floribus spirali-secundis, nectarii labio indiviso crenulato. S. N. 812. Fl. D. 387. St. Helv. t. 38. P. pr.

1095. *O.*

1095. *O. ovata*, bulbo fibroso, caule bifolio, foliis ovatis, nectarii labio bifido. Fl. Sv. 809. Fl. N. 127. Fl. D. 137. St. Helv. t. 37. P. pr. sylv.
1096. *O. cordata*, bulbo fibroso, caulē bifolio, foliis cordatis, nectarii labio bifido basi utrinquedentato. Fl. Sv. 810. Fl. N. 668. t. 7. f. 6. St. Helv. t. 22. P. sylv. acer.
1097. *O. liliifolia*, bulbo subrotundo, scapo nudo, foliis lanceolatis, nectarii labio integro, petalis dorsibus linearibus. Fl. Sv. 811. Fl. N. 962. P.
1098. *O. Loesfelii*, bulbo subrotundo, scapo nudo trigono, nectarii labio ovato. Fl. Sv. 812. *O. paludosa* Fl. D. 877. P. pr. ul.
1099. *O. paludosa*, bulbo subrotundo, scapo subnudo pentagono, foliis apice scabris, nectarii labio integro. Fl. Sv. 813. Fl. N. 965. *Malaxis paludosa* Swartz N. Act. Holm. 1799. p. 12. t. 6. P. pal.
1100. *O. monophylla*, bulbo subrotundo, scapo nudo, folio ovato, nectarii labio integro. Fl. Sv. 814. St. Helv. t. 36. P.
1101. *O. monorchis*, bulbo globoso, scapo nudo, nectarii labio trifido cruciato. Fl. Sv. 810. Fl. N. 644. Fl. D. 102. St. Helv. t. 22. P. pr. ul.
1102. *O. alpina*, bulbis ovatis, scapo nudo, foliis subulatis, nectarii labio indiviso obtuso utrinque unidentato. Fl. Sv. 817. Fl. N. 1098. Fl. D. 452. St. Helv. t. 22. Jacqu. Vind. t. 9. P. alp.
- † 1103. *O. Camtschatea*, bulbis - - - scapo filiformi vaginato, racemo laxo, nectarii labio lineari bifido. S. N. 814. Amoen. Acad. II. p. 361. t. 4. f. 24. Fl. Isl. 9. P.
1104. *O. myodes*, bulbis subrotundis, scapo folioso, nectarii labio oblongo quadripartito: laciniis duabus mediis elongatis lanceolatis. S. N. 813. *O. infestifera myodes* Fl. Sv. 818. Fl. N. 967. t. 5. f. 2. St. Helv. t. 24. P. pr. ul.

SERAPIAS.

1105. *S. latifolia*, bulbis fibrosis, foliis ovatis amplexicaulibus, floribus pendulis. S. N. 814. Fl. N. 970. t. 5. f. 3. Fl. D. 811. St. Helv. t. 40. P. pr. sylv.
1106. *S. longifolia*, bulbis fibrosis, foliis ensiformibus sessilibus, floribus pendulis: labio obtuso plicato. S. N. 815. Fl. N. 1103. Fl. D. 267. A. Svec. 1769. p. 248. Ros. Obs. p. 55. n. 46. St. Helv. t. 41. P. pr. sylv.
1107. *S. ensifolia*, bulbis fibrosis, foliis ensiformibus distichis, bracteis minutissimis, floribus erectis: nectarii labio obtuso petalis brevior. S. N. 815. Fl. D. 506. Ros. Obs. p. 54. n. 46. P. pr. sylv.
1108. *S. lancifolia*, bulbis fibrosis, foliis ovato-lanceolatis, bracteis longitudine capsulae, floribus erectis: nectarii labio obtuso petalis brevior. S. N. 815. St. Helv. t. 41. P. pr. sylv.
1109. *S. rubra*, bulbis fibrosis, foliis ensiformibus, floribus erectis: nectarii labio acuto. S. N. 816. Fl. D. 345. St. Helv. t. 42. P. pr. sylv.

CYPRIPEDIUM.

1110. *C. Calceolus*, radicibus fibrosis, foliis ovato-lanceolatis caulinis. Fl. Sv. 820. Fl. N. 1113. Fl. D. 999. St. Helv. t. 43. P. sylv.
1111. *C. bulbosum*, bulbo subrotundo, folio subrotundo radicali. Fl. Sv. 821. Fl. L. t. 12. f. 5. Gmel. Sib. L. t. 2. f. 1. P. alp.

2. H E X A N D R I A.

ARISTOLOCHIA.

1112. *A. Clematitis*, foliis cordatis, caule erecto, floribus axillaribus confertis. S. N. 825. Fn. Sv. 558. Weig. Suppl. 901. P. f.

3. P O L Y A N D R I A .

ARUM.

1113. *A. maculatum*, acaule, foliis hastatis integerri-
mis, spadice clavato. S. N. 328. Fl. D. 505.
P. umbr. hum.

CALLA.

1114. *C. palustris*, foliis cordatis, spatha planâ, spadice
undique hermaphrodito. Fl. Sv. 822. Fl. N. 727.
Fl. D. 422. P. pal.

ZOSTERA.

1115. *Z. marina*, pericarpis sessilibus. Fl. Sv. 823.
Fl. D. 15. Fl. N. 371. P. m.
1116. *Z. oceanica*, pericarpis pedicellatis oliviformibus.
S. N. 829. Fl. N. 1079. P. m.

XXI. MONOECIA.

I. MONANDRIA.

ZANNICHELLIA. Mas. Cal. o. Cor. o.

Foem. Cal. 1-phyll. Cor. o. *Germina* sub-
quaterna. *Sem.* totidem.*Rad.* fibrosa. *Caul.* articulata. *Fol.* graminea,
submersa. *Flor.* sessilis, axillaris. *Fulcr.* sti-
pulae vaginantes.CHARA. Mas. Cal. o. Cor. o. *Anthera* germini sub-
jecta.Foem. Cal. 4-phyll. Cor. o. *Stigma* 5-fid.
Sem. 1.*Rad.* fibrosa. *Caul.* articulata. *Fol.* subulata,
verticillata, florifera, submersa. *Flor.* solita-
ria., sessilis, epiphylla. *Fulcr.* aculei.

2. DIANDRIA.

LEMNA. Mas. Cal. 1-phyll. Cor. o.

Foem. Cal. 1-phyll. Cor. o. *Styl.* 1. *Capf.*
1-locul.*Rad.* filiformis, simplicissima. *Caul.* o. *Fol.* sim-
plex, natans. *Flor.* axillaris. *Fulcr.* vaginae
radicum.

3. TRIANDRIA.

TYPHA. Mas. *Amentum* cylindricum. Cal. obsoletus
3-phyll. Cor. o.Foem. *Ament.* cylindricum. Cal. capillo
villoso. Cor. o. *Sem.* 1. insidens pappo
capillari.*Rad.* repens, articulata. *Caul.* culmosa, farcta.
Fol. graminea. *Flor.* spicata, compacta.
Fulcr. o.

SPARGA.

SPARGANIUM. Mas. *Ament.* subrotundum. *Cal.* 3-phyll.
Cor. 6.

Foem. *Ament.* *Cal.* *Cor.* Maris. *Stigm.* 2-fid.
Drupa exfucca 1-sperma.

Rad. repens, articulata. *Caul.* culmosa, farcta,
flexuosa. *Fol.* graminea. *Flor.* subracemosa.
Fulcr. 6.

CAREX. Mas. *Amentum* imbricatum. *Cal.* 1-phyll.
Cor. 6.

Foem. *Ament.* *Cal.* *Cor.* Maris. *Nectar.* in-
flatum, 2-dentatum. *Stigm.* 2-3. *Semen*
3-quetrum intra nectarium.

Rad. fibrosa l. repens. *Caul.* culmosa, farcta,
saepius triquetra. *Fol.* graminea, aspera.
Flor. spicata. *Fulcr.* bracteae.

AXYRIS. Mas. *Cal.* 3-partit. *Cor.* 6.

Foem. *Cal.* 2-phyll. *Cor.* 6. *Styli* 2. *Sem.* 1.

Rad. subramosa. *Caul.* herbacea. *Fol.* simplex,
sparsa. *Flor.* axillaris et terminalis. *Fulcr.*

Amaranthus vide Monoecia Pentandria.

4 T E T R A N D R I A.

LITTORELLA. Mas. *Cal.* 4-phyll. *Cor.* 4-fida. *Stam.*
longa.

Foem. *Cal.* 6. *Cor.* sub 4-fida. *Styl.* longus.
Sem. Nux.

Rad. fibrosa. *Caul.* 6. *Fol.* radicalis, graminea.
Flor. axillaris, mascula pedunculata, foemi-
nea sessilis. *Fulcr.* squamae.

BETULA. Mas. *Amentum* cylindr. *Cal.* 1-phyll. 3-fid.
3-flor. *Cor.* 4-part.

Foem. *Amentum* cylindr. *Cal.* 1-phyll.
sub-3-fid. 2-flor. *Sem.* utrinque membrana
alatum.

Rad. lignosa. *Caul.* arborea. *Fol.* simplex, sparsa.
Flor. amentacea. *Fulcr.* stipulae caducae.

URTICA. Mas. *Cal.* 4-phyll. *Cor.* o. *Nectar.* centralis cyathiforme.

Foem. *Cal.* 2-valvis. *Cor.* o. *Sem.* 1.

Rad. ramosa. *Caul.* herbacea. *Fol.* opposita, simplex. *Flor.* amentacea. *Fulcr.* stipulae.

5. PENTANDRIA.

XANTHIUM. Mas. *Cal.* communis imbricat. *Cor.* 1-pet. 5-fida, infundib. *Rec.* paleaceum.

Foem. *Involut.* 2-phyll. 2-flor. *Cor.* o.

Drupa ficca, muricata, 2-fida. *Nucleus* 2-locul.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex, alternata. *Flor.* axillaris. *Fulcr.* stipulae.

AMARANTHUS. Mas. *Cal.* 3-5-phyll. *Cor.* o. *Stam.* 3-4.

Foem. *Cal.* maris. *Cor.* o. *Styli* 3. *Caps.* circumscissa. *Sem.* 1.

Rad. ramosa. *Caul.* herbacea. *Fol.* simplex, alternata. *Flor.* axillaris, glomerata. *Fulcr.* o.

6. POLYANDRIA.

GERATOPHYLLUM. Mas. *Cal.* multipartita. *Cor.* o. *Stam.* 16-20.

Foem. *Cal.* *Cor.* maris. *Pist.* 1. *Styl.* o. *Sem.* 1 - nudum.

Rad. fibrosa. *Caul.* herbacea. *Fol.* submersa, verticillata, dichotoma. *Flor.* axillaris, solitaria. *Fulcr.* o.

MYRIOPHYLLUM. Mas. *Cal.* 4-phyll. *Cor.* o. *Stam.* 8.

Foem. *Cal.* *Cor.* maris. *Pistilla* 4. *Styli* o. *Sem.* 4-nuda.

Rad. geniculata, repens. *Caul.* herbacea, articulata. *Fol.* submersa verticillata, composita. *Flor.* verticillato-spicata. *Fulcr.* squamae.

SAGITTARIA. Mas. *Cal.* 3-phyll. *Cor.* 3-pet. *Stam.* fere 24.

Foem. *Cal.* *Cor.* maris. *Pist.* multa. *Sem.* multa, nuda.

Rad. fibrosa, horizontalis. *Caul.* scaposa. *Fol.* simplex, radicalis. *Flor.* verticillato-subracemosa. *Fulcr.* bracteae.

POTERIUM. Mas. *Cal.* 4-phyll. *Cor.* 4-partit. *Stam.* multa.

Foem. *Cal.* *Cor.* maris. *Pist.* 2. *Bacca* et tubo corollae indurato.

Rad. ramofo-fibrosa. *Caul.* herbacea. *Fol.* pinnata. *Flor.* capitata. *Fulcr.* stipulae, bracteae.

QUERCUS. Mas. *Amentum* laxum. *Cal.* 5-fid. *Cor.* o. *Stam.* 5-10.

Foem. *Cal.* 1-phyll. integerr. *Cor.* o. *Styli* 2-5. *Sem.* 1. ovatum.

Rad. lignosa. *Caul.* arborea. *Fol.* simplex, alterna, sinuata. *Flor.* axillaris. *Fulcr.* o.

FAGUS. Mas. *Amentum.* *Cal.* 5-fid. campanul. *Cor.* o. *Stam.* 12.

Foem. *Cal.* 4-dent. *Cor.* 6-pet. *Styli* 3. *Caps.* muricata, 4-valv. *Sem.* 2.

Rad. lignosa. *Caul.* arborea. *Fol.* simplex, integerrima. *Flor.* axillaris. *Fulcr.* stipulae deciduae.

CARPINUS. Mas. *Ament.* squamis 1-flor. ciliatis. *Cor.* o. *Stam.* 20.

Foem. *Ament.* squamis 1-flor. villosis. *Cor.* o. *Germ.* 2. singulo *Styl.* 2. *Nux* ovata.

Rad. lignosa. *Caul.* arborea. *Fol.* simplex, integerrima, plicata. *Flor.* subterminalis. *Fulcr.* o.

CORYLUS. Mas. *Ament.* squamis 3-fid. *Cor.* o. *Stam.* 8.
Foem. *Cal.* 2-phyll. lacerus. *Cor.* o. *Styli* 2.
Nux ovata.

Rad. lignosa. *Caul.* fruticosa. *Fol.* simplex, rugosa. *Flor.* terminalis ramulorum. *Fulcr.* stipulae deciduae.

7. MONADELPHIA.

PINUS. Mas. *Ament.* *Cal.* 4-phyll. *Cor.* o. *Stam.* plurima. *Antherae* nudaе.
Foem. *Cal.* strobilus squamis bifloris. *Cor.* o.
Pist. 1. *Nux* alata.

Rad. lignosa. *Caul.* arborea ramis verticillatis.
Fol. acerosa, sempervirens. *Flor.* axillaris.
Fulcr. stipulae.

8. SYNGENESIA.

BRYONIA. Mas. *Cal.* 5-dent. *Cor.* 5 part. *Stam.* 3.
Foem. *Cal.* *Cor.* maris. *Stylus* 3-fid. *Bacca* globosa, polysperma.
Rad. fusiformis, crassa. *Caul.* herbacea, debilis.
Fol. simplex, palmata. *Flor.* axillaris. *Fulcr.* Cirrhi.

XXI. MONOECIA.

I. M O N A N D R I A.

ZANNICHELLIA.

1117. *Z. palustris*, Fl. Sv. 824. Fl. D. 67. aqu.

CHARA.

1118. *C. tomentosa*, aculeis caulinis ovatis. Fl. Sv. 1132. P. aqu.
 1119. *C. vulgaris*, caulibus laevibus, foliis interne dentatis. Fl. Sv. 1131. Fl. N. 751. P. qu.
 1120. *C. hispida*, caulium aculeis capillaribus confertis. Fl. Sv. 1133. Fl. N. 1118. Fl. D. 154. P. qu.
 1121. *C. flexilis*, caulium articulis inermibus diaphanis superne latioribus. Fl. Sv. 1134. P. l. m.

2. D I A N D R I A.

LEMNA.

1122. *L. trifulca*, foliis petiolatis lanceolatis. Fl. Sv. 828. Fl. N. 779. P. qu.
 1123. *L. minor*, foliis sessilibus utrinque planiusculis, radicibus solitariis. Fl. Sv. 827. Vaill. Par. t. 20. f. 3. Fl. N. 783. A. qu.
 1124. *L. gibba*, foliis sessilibus subtus hemisphaericis, radicibus solitariis. Fl. Sv. 826. Fl. N. 778. y). A. qu.
 1125. *L. polyrrhiza*, foliis sessilibus, radicibus confertis. Fl. Sv. 825. Vaill. Par. t. 20. f. 2. Fl. N. 777. A. qu.

O 4

3. TRI-

9) Varietatem antecedentis suspicatur *Cel. Afzelius* Diff. cit.

3. TRIANDRIA.

TYPHA.

1126. *T. latifolia*, foliis subensiformibus, spica mascula foemineaque approximatis. Fl. Sv. 831. Fl. D. 645. P. l. lac. fluv.
1127. *T. angustifolia*, foliis semicylindricis, spica mascula foemineaque remotis. Fl. Sv. 830. Fl. N. 342. Fl. D. 815. P. l. lac. fluv.

SPARGANIUM.

1128. *S. ramosum*, foliis ensiformibus triquetris, caule ramoso. H. Kew. III. 323. *Sparg. erectum* Fl. Sv. 832. α . Fl. N. 499. P. qu.
1129. *S. simplex*, foliis ensiformibus planis, caule simplici. H. Kew. III. 324. Fl. D. 932. *Sp. erectum* Fl. Sv. 832. β . P. qu.
1130. *S. natans*, foliis decumbentibus planis. Fl. Sv. 833. Fl. N. 500. Fl. D. 260. P. aqu. fl.

CAREX.

* *spica una simplici.*

1131. *C. dioica*, spica simplici dioica. Fl. Sv. 833. Fl. N. 715. Fl. D. 369. P. pal. sylv. St. 2.
1132. *C. capitata*, spica simplici androgyna ovata: superne mascula, foemineis digynis, capsulis imbricato-patulis. S. N. 842. Fl. D. 372. P. pal.
1133. *C. obtusata*, spica simplici androgyna superne mascula, foemineis trigynis, capsulis ovatis obtuse triquetris, foliis planis. N. A. Holm. 1793. p. 69. t. 4. P. ficc. ar.
1134. *C. pulicaris*, spica simplici androgyna, superne mascula, capsulis feminibus teretibus. S. N. 842. Fl. Sv. 834. Fl. N. 795. Fl. D. 166. fig. pess.
- C. psyllophora* Ehrh. Beytr. I. 186. P. pal. St. 2.

1135. *C. Leucoglochin*, spica simplici androgyna, superne mascula, capsulis reflexis, seminibus triquetris. Suppl. Plant. 413. Ehrh. Beytr. I. 186. *C. pauciflora* Ligthf. Fl. Scot. 543. t. VI. P. pal. 8. 3.

** *Spicis pluribus androgynis.*

- † 1136. *C. incurva*, spica composita ovata ebracteata: spiculis androgynis, culmo subtriquetro, foliis convolutis. Nat. H. Selfk. Skr. II. 1. p. 32. Ligthf. Fl. Scot. 544. t. 24. *C. maritima* Fl. N. 1017. Fl. D. 432. *C. capitata* β *maritima* Prodr. Fl. Scand. 1021. P. alp. St. 2.
1137. *C. chordorhiza*, spica composita: spiculis androgynis approximatis superne masculis, capsulis compressis, radice repente filiformi. Ehrh. Beytr. I. p. 186. Suppl. Pl. 414. *C. uliginosa* Sp. Pl. Ed. 1. 2. n. 5. P. pal.
1138. *C. arenaria*, spica composita: spiculis androgynis, inferioribus remotioribus foliolo longiore instructis, culmo triquetro. Fl. Sv. 835. Fl. N. 921. Fl. D. 425. P. a. St. 2.
1139. *C. leporina*, spica composita: spiculis ovatis sessilibus approximatis alternis androgynis nudis. Fl. Sv. 837. Fl. N. 813. a). P. pal. St. 2.
1140. *C. disticha*, spica composita subdisticha: spiculis ovatis imbricatis androgynis foliolo longiore instructis, culmo triquetro. Schreb. Lips. p. 63. *C. leporina* Prodr. Fl. Scand. 1025. a. y. *C. arenaria* L. Fl. Herb. t. 14. f. 2. P. pal. St. 2.

O 5

1141. C.

- z) Non vero Fl. Svec. 836. quae, confirmante jam *Cel. Smith* in litt. est *Scirpus Caricis* noster. Cfr. N. Act. Holm. 1793.
- e) *C. leporina* Fl. D. t. 294. forte huc pertinet, quamvis tunc minus bona icon.

1141. *C. vulpina* spica supradecomposita inferne laxiore: spiculis ovatis androgynis glomeratis superne masculis. Fl. Sv. 838. Fl. N. 818. Fl. D. 308. L. Fl. Herb. t. 14. f. 5. ? P. pal. St. 2.
 β . *C. intermedia*. b).
1142. *C. muricata*, spica composita, spiculis subovatis sessilibus remotis androgynis (superne masculis) capsulis acutis divergentibus bifido-spinosis. Fl. Sv. 839. Fl. N. 810. *C. canescens* Leers Fl. Herb. t. 14. f. 3. P. pr. St. 2.
1143. *C. echinata*, spica composita, spiculis subovatis sessilibus remotis androgynis (inferne masculis) capsulis acutis simpliciter mucronatis. *C. Leersii* Willden. Fl. 29. *C. muricata* Fl. D. 284. Leers Fl. Herb. t. 14. f. 8. P. pr. hum. St. 2 ?
1144. *C. loliacea*, spica composita, spiculis subovatis sessilibus remotis androgynis (inferne masculis) capsulis ovatis teretiusculis muticis divaricatis. Fl. Sv. 840. Fl. N. 922. P. pr. pal. St. 2.
1145. *C. Heleonastes*, spica composita, spiculis androgynis approximatis superne foemineis, capsulis imbricatis lateribus integerrimis Ehrh. Beytr. I. 136. Suppl. Pl. p. 414. P. pr. ul. St. 2.
1146. *C. microstachya*, spica composita, spiculis androgynis oblongis approximatis superne foemineis: terminali majore, capsulis adpressis obtectis. Ehrh. Beytr. III. 72. P. pr. ul. St. 2.
1147. *C. remota*, spicis ovatis subsessilibus remotis androgynis (superne foemineis) bracteis culmum aequantibus. S. N. 844. Fl. D. 370. P. sylv. hum. St. 2.

1148. *C.*

b) Est sola varietas *C. Vulpinae* spica haud supradecomposita bracteaque instructa. In icone Fl. D. nulla bractea, Leers vero plures bracteas adpingit.

1148. *C. elongata*, spiculis oblongis sessilibus remotis androgynis (superne foemineis) capsulis ovatis acutis. Fl. Sv. 841. Fl. N. 713. P. pr. pal. St. 2.
1149. *C. canescens*, spiculis subrotundis remotis sessilibus, obtusis androgynis (superne foemineis) capsulis ovatis obtusiusculis. Fl. Sv. 842. Fl. N. 776. Fl. D. 285. P. pr. ul. St. 2.
1150. *C. tenella*, spiculis ovatis remotis subsessilibus androgynis (superne foemineis) capsulis ovatis acutis. c), P. St. 2.
1151. *C. multiculmis*, spiculis oblongis sessilibus approximatis androgynis (superne foemineis) capsulis imbricatis depressis acutis amenti squamis longioribus. d), P. St. 2.
1152. *C. paniculata*, racemo composito, spiculis androgynis (apice foemineis). S. N. 844. Weig. Suppl. 909. Leers Fl. Herb. t. 14. f. 4. P. pal. St. 2.
 β. minor. *C. diandra* Schrank.
- † 1153. *C. norvegica*, spiculis ovatis obtusis congestis pedicellatis subternis androgynis, capsulis subrotundis. *Carex* Fl. D. 403. *C. Halleri* Fl. N. 849 e). P. alp. St. 3.

*** spi-

- e) Characterem nullibi adhuc dedit Cl. inventor! *Ehrharr*, mancum sine dubio ex specimine in Gram. et Cal. Dec. X. II. 98. obvio desumsi; simul vero fateri debeo, me hanc a *canescente* L. quam scil. pro hac in Herbario possideo, haud distinguere posse. Quam pro *Carice cinerea* Poll. dedit *Cel. Willdenow* huic persimilis, at manifeste diversa spiculis basi foemineis.
- d) Etiam hujus characterem e specimine in *Ehrh. Gr. et Cal.* Dec. IX. n. 88. desumsi.
- e) Suspicio hanc ad quartam divisionem pertinere et minutam spicam marem forte caducam. adesse, saltem facies et numerus stigmatum suadet, nec locum florum masculorum detegere potui in specimine a *Cel. Vahl*, dato. A *Carice Hall.* 1356 diversa omnino.

*** *spicis sexu distinctis: foemineis sessilibus.*

1154. *C. flava*, spicis confertis subsessilibus subrotundis: mascula lineari, capsulis acutis recurvis. Fl. Sv. 843. Fl. N. 526. Leers Fl. Herb. t. 15. f. 6. f). P. pal. St. 3.
1155. *C. pedata*, spicis foemineis sessilibus oblongis: inferiore axillari, foliis filiformibus. S. N. 844. Bauh. Prodr. 9. P. pal. alp.
1156. *C. digitata*, spicis linearibus erectis: mascula brevior inferioreque, bracteis aphyllis, capsulis distinctibus. Fl. Sv. 844. Fl. N. 1049. Leers Fl. H. t. 16. f. 4. P. sylv. St. 3.
1157. *C. montana*, spicis foemineis sessilibus subsolitariis ovatis masculae approximatis, culmo nudo, capsulis pubescentibus. Fl. Sv. 845. Fl. N. 708. Leers Herb. t. 16. f. 6. P. mont. St. 3.
1158. *C. globularis*, spica mascula oblonga, foeminea sessili ovata folio florali brevior approximata. Fl. Sv. 846. Fl. Lapp. 336. Nat. H. Selsk. Skr. II. p. 45. g). P. alp. St. 3.
1159. *C. filiformis*, spica mascula oblonga, foemineis sessilibus oblongis: inferiore foliolo proprio brevior. Fl. Sv. 847. Fl. N. 1048. Leers Fl. Herb. t. 16. f. 5. P. nem. sylv. St. 3.
1160. *C. pilulifera*, spicis terminalibus confertis subrotundis: mascula oblonga. S. N. 845. Fl. N. 583. *C. Oederi* Prodr. Fl. Scand. 1035. *C. decumbens* Ehrh. Gr. et Cal. 70. *Carex* Fl. D. 371. P. a. glar. St. 3.

1161. *C.*

f) Specimen misit *Cel. Weigel* ex Pomerania huic simillimum, at scripsit simul unicam remotissimam foemineam spicam prope radicem ex axilla imi folii prodire, an pedunculatam vel sessilem non dixit.

g) Fl. D. 443. si huc spectet, pessima.

1161. *C. saxatilis*, spicis tribus ovatis sessilibus alternis, mascula oblonga. Fl. Sv. 848. Fl. N. 449. Fl. D. 159. ? P. mont.

1162. *C. humilis*, spica mascula oblonga, foemineis serioribus remotis subpedunculatis paucifloris, culmo foliis brevior. Schreb. Lips. p. 65. 8). P.

**** *Spicis sexu distinctis: foemineis pedunculatis.*

1163. *C. atrata*, spicis androgynis terminalibus pedunculatis: florentibus erectis, fructiferis pendulis, Fl. Sv. 849. Fl. N. 451. Fl. D. 158. Nat. H. Selfk. Skr. II. 43. P. alp. St. 3.

1164. *C. limosa*, spicis ovatis pendulis: mascula longiore erectiore, radice repente. Fl. Sv. 850. Fl. N. 714. Fl. D. 646. Nat. H. Selfk. Skr. II. 41. *C. elegans* Willd. Fl. Ber. 104. t. 1. f. 4. P. pal. St. 3. β . *pauciflora*. Vahl. l. c. alp.

1165. *C. capillaris*, spicis pendulis: mascula erecta, foemineis oblongis, capsulis nudis acuminatis. Fl. Sv. 851. Fl. N. 796. Fl. D. 168. P. pr. ul. pal. St. 3.

1166. *C. pallescens*, spicis pendulis: mascula erecta: foemineis ovatis imbricatis, capsulis confertis obtusis. Fl. Sv. 852. Fl. N. 923. Fl. D. 444. L. Fl. Herb. t. 15. f. 4. *C. limosa pallida* Prodr. Fl. Scand. 1045 β . P. pr. pal. St. 3.

1167. *C. panicea*, spicis pedunculatis remotis: foemineis linearibus, capsulis obtusiusculis inflatis. Fl. Sv. 853. Fl. N. 325. Fl. D. 261. L. Fl. Herb. t. 15. f. 5. P. pal. St. 3.

1168. *C. pseudocyperus*, spicis pendulis, pedunculis subgeminatis, capsulis reversis biaristatis. Fl. Sv. 854. 1). P. fossa.

1169. *C.*

h) *Cel. Kölpin.*

i) Spicae saepius androgynae: mascula apice foeminea, foemineisque apice masculis.

- †169. *C. sylvatica*, spicis pendulis: mascula erecta, foemineis filiformibus pedunculo brevioribus, capsulis acutis distantibus. Schr. Lips. p. 62. Fl. D. 404.
C. Drymeja Ehrh. Beytr. I. 181. Suppl. Pl. 414.
C. capillaris L. Fl. Herb. 720. t. 15. f. 2. P. sylv. St. 3.
- †170. *C. difflans*, spicis remotissimis subsessilibus, bractea vaginante, capsulis angulatis mucronatis. S. N. 845. Ehrh. Cal. 40. P. pal. St. 3.
- †171. *C. stolonifera*, spicis erectis approximatis, foeminea inferiore pedunculata. k). P. pal. St. 3.
- †172. *C. caespitosa*, spicis erectis cylindricis ternis subsessilibus: mascula terminali, caule triquetro. Fl. Sv. 855. Fl. N. 582. P. pal. St. 2.

***** *spicis masculis pluribus.*

- †173. *C. maritima*, spicis foemineis teretibus remotis pendulis: calycibus aristatis, mascula erecta. Vahl. Nat. H. Selfk. Skr. II. p. 13. Fl. D. 704. l). P. litt. riv. St. 2.
- †174. *C. flacca*, spicis masculis geminis rectis, foemineis pedunculatis pendulis, capsulis ovato-triquetris confertis. *C. flacca* Schreb. Lips. app. 669. P. pr. hum. St. 3.
- †175. *C. lasiocarpa*, spicis masculis geminis: foemineis remotis subsessilibus, bractea longissima, capsulis hirsutis. Ehrh. Beytr. 3. p. 73. *C. tomentosa* Mant. Pl. 2. p. 123. m). *C. splendida* Willd. Fl. Ber. 103. t. 1. f. 3. P. pal. St. 3.

1176. C

k) E specimine sicco in Cal. Gr. n. 99. allata.

l) Praeter spicam superiorem androgynam saepe 2 masculae.

m) Certe multa suadent. Spica foeminea suprema haud raro apice mascula.

1176. *C. acuta*, spicis masculis pluribus, foemineis subsessilibus, capsulis obtusiusculis. Fl. Sv. 857. Fl. N. 38. n). P. St. 2.

α *nigra*, verna. pr. hum.

β. *rufa*, major. aqu.

γ. *gracilis*, H. Kew. III. 331.

1177. *C. crassa*, spicis masculis pluribus triquetris foemineis pedunculatis, calycibus linearibus acuminatis, capsulis subinflatis bicornibus. *C. crassa* Ehrh. Beytr. IV. p. 43. *C. riparia* Curtis. *C. vesicaria* a. Sp. Plant. 1388. P. riv. fylv. S. 3.

1178. *C. vesicaria*, spicis masculis pluribus: foemineis pedunculatis, capsulis inflatis acuminatis, culmo acute triquetro. Fl. Sv. 856. Fl. D. 647. P. qu. St. 3.

1179. *C. obtusangula*, spicis masculis pluribus: foemineis pedunculatis, capsulis inflatis acuminatis, culmo obsolete triquetro. *C. obtusangula* Ehrhart. Cal. Gr. Dec. V. n. 50. P. qu. St. 3.

1180. *C. hirta*, spicis remotis: masculis pluribus: foemineis subpedunculatis erectis, squamis longis aristatis, capsulis hirtis. Fl. Sv. 858. Fl. N. 740. Fl. D. 379. L. Fl. Herb. t. 16. f. 3. P. a. ficc. St. 3.

AXYRIS.

†1181. *A. prostrata*, foliis obovatis, caule subdiviso, floribus capitatis. S. N. 846. Gm. Sib. 3. t. 4. f. 2. Fl. N. 591. A.

4. TETRAN.

n) Haud raro diandra. Ehrh. Beytr. III. p. 113. et spicae foemineae apice masculae.

4. TETRANDRIA.

LITTORELLA.

1182. *L. lacustris*. S. N. 848. *L. juncea*. A. Holm. 1768.
p. 341. *Plantago uniflora*. Fl. Sv. 135. Fl. N. 833.
Fl. D. 170. P. l. lac.

BETULA.

1183. *B. alba*, foliis deltoideis acutis duplicato-ferratis glabris, strobilorum squamis lobis lateralibus rotundatis, petiolis glabris pedunculo longioribus. H. Kew. III. 337. Fl. Sv. 859. Fl. N. 324. Arb. sylv.

α. *vulgaris*, ramis adscendentibus.

β. *pendula*, ramis dependentibus.

γ. *laciniata*, foliis profunde laciniatis, laciniis acutis. *B. hybrida* N. Act. Svec. 1786.
p. 186. t. 6.

1184. *B. nana*, foliis orbiculatis crenatis. Fl. Sv. 860.
Fl. L. t. 6. f. 4. Am. Acad. I. t. 1. Fl. N. 419.
Fl. D. 91. Fr. pal. sylv. alp.

1185. *B. Alnus*, pedunculis ramosis, foliis subrotundo-cuneiformibus obtusissimis glutinosis: axillis venarum subtus villosis. H. Kew. III. 338. *B. Alnus* α.
Fl. Sv. 861. Fl. N. 151. Arb. hum.

β. *pumila*, heterophylla it. Westrog. 181.

1186. *B. incana*, pedunculis ramosis, foliis subrotundo-ellipticis acutis subtus pubescentibus: axillis venarum nudis, stipulis lanceolatis. H. Kew. III. 339.
S. N. 849. Suppl. Pl. 417. *B. Alnus* β. Fl. Sv. 861.
Arb. mont.

URTICA.

1187. *U. urens*, foliis oppositis ovalibus. Fl. Sv. 862.
Fl. N. 459. Fl. D. 739. A. c. r.

1188. *U.*

MONOECIA. PENTANDRIA. *Xanthium*.- 225.

1188. *U. dioica*, foliis oppositis cordatis, racemis geminis. Fl. Sv. 863. Fl. N. 372. Fl. D. 746. P. r. f. c.

5. PENTANDRIA.

XANTHIUM.

1189. *X. strumarium*, caule inermi, foliis cordatis trinervatis. Fl. Sv. 864. Fl. D. 970. A. r.

AMARANTHUS.

1190. *A. Blitum*, glomerulis lateralibus: floribus trifidis, foliis ovatis retusis, caule diffuso. Fl. Sv. 865. Willd. Amar. 15. A. c.

6. POLYANDRIA.

CERATOPHYLLUM.

1191. *C. demersum*, foliis dichotomo-bigemini, fructibus trispinosis. Fl. Sv. 866. P. qu.
1192. *C. submersum*, foliis dichotomo-trigemini, fructibus muticis. S. N. 856. Fl. D. 510. P. qu.

MYRIOPHYLLUM.

1193. *M. spicatum*, floribus masculis interrupte spicatis. Fl. Sv. 867. Fl. N. 386. Fl. D. 681. P. qu. lac.
1194. *M. verticillatum*, floribus verticillatis hermaphroditis. Fl. Sv. 868. P. qu.

SAGITTARIA.

1195. *S. sagittifolia*, foliis sagittatis acutis. Fl. Sv. 869. Fl. N. 875. Fl. D. 172. P. lac. foss.

POTERIUM.

- †1196. *P. Sanguisorba*, inerme, caulibus subangulosis.
S. N. 857. Weig. Suppl. 917. P.

QUERCUS.

1197. *Q. Robur*, foliis oblongis glabris sinuatis: lobis rotundatis, glandibus oblongis. H. Kew. III. 358. Fl. Sv. 870. Fl. N. 380. Arb. fylv.

α. *pedunculata*, Auct. *Formina*, foliis praecocioribus superne vix dilatatis, breviter petiolatis, fructibus longe pedunculatis. *Sommer - Ek.*

β. *sessilis*, Auct. *Mas*, foliis serioribus superne dilatatis longius petiolatis, fructibus breviter pedunculatis. *Winter - Ek.*

FAGUS.

1198. *F. sylvatica*, foliis ovatis obsolete ferratis. Fl. Sv. 871. Fl. N. 638. Arb. fylv.

CARPINUS.

1199. *C. Betulus*, squamis strobilorum planis. Fl. Sv. 872. Arb. nem. fylv.

CORYLUS.

1200. *C. Avellana*, stipulis ovatis obtusis. Fl. Sv. 873. Fl. N. 443. Fr. nem.

7. MONADELPHIA.

PINUS.

1201. *P. sylvestris*, foliis geminis: primordialibus solitariis glabris. Fl. Sv. 874. Fl. N. 337. Arb. fylv.

1202. *P.*

1202. *P. Abies*, foliis solitariis subulatis mucronatis
laevibus bifariam versis. Fl. Sv. 875. Fl. N. 39.
Fl. D. 193. Arb. sylv.

β. *viminalis*, ramulis elongatis pendentibus
flexuosis laevis. Aët. Holm. 1777. p. 313.
t. 8. 9.

8. SYNGENESIA.

BRYONIA.

1203. *B. alba*, foliis palmatis utrinque calloso-scabris,
floribus monoicis. Fl. Sv. 876. Fl. D. 813. P. f.

XXII. D I O E C I A.

I. M O N A N D R I A.

- NAJAS.** Mas. *Cal.* cylindr. 2-fid. *Cor.* 4-fid. *Fil.* o.
 Foem. *Cal.* et *Cor.* o. *Pist.* 1. *Caps.* ovata, 1-loc.
Rad. fibrosa. *Caul.* herbacea, dichotoma. *Fol.*
 opposita, dentata, submersa. *Flor.* axillaris,
 solitaria. *Fulcr.* o.
Salix purpurea vide ordin. sequ.

2. D I A N D R I A.

- VALLISNERIA.** Mas. *Spatha* 2-part. *Spadix* tectus
 • flosculis. *Cor.* 3-part.
 Foem. *Spatha* 2 fid. 1-flor. *Cal.* 3-part.
 super. *Cor.* 3-pet. *Styli* 3. *Caps.* 1-locul.
 polysperma.
Rad. articulata, stolonifera. *Caul.* scaposa, foe-
 minea spiralis. *Fol.* simplex, reticulata, de-
 mersa. *Flor.* foeminea solitaria: mascula ca-
 pitata. *Fulcr.* o.
SALIX. Mas. *Amentum* squama. *Cor.* o. *Stam.* 1-12.
Gland. baseos nectarifera.
 Foem. *Ament.* et *Cor.* maris. *Stylus* 2-fid.
Caps. 1-loc. 2-valv. *Sem.* papposa.
Rad. lignosa. *Caul.* fruticans l. arborea. *Fol.*
 simplex, alterna. *Flor.* amentacea. *Fulcr.*
 stipulae.

3. T R I A N D R I A.

- EMPETRUM.** Mas. *Cal.* 3-part. *Cor.* 3-pet. *Stam.* longa.
 Foem. *Cal.* et *Cor.* maris. *Styli* 9. *Bacca*
 9-sperma.
Rad. lignosa. *Caul.* fruticulosa. *Fol.* acerosa.
Flor. axillaris, sessilis. *Fulcr.* o.
Valeriana dioica vide Triandria Monogynia.
Carex dioica vide Monoecia Triandria.
Salix triandra vide Dioecia Diandria.

4. TETRAN

4. TETRANDRIA.

VISCUM. Mas. *Cal.* 4-part. *Cor.* o. *Antherae* calyci adnatae.

Foem. *Cal.* 4-phyll. super. *Cor.* o. *Styli* o.
Bacca 1-sperma. *Sem.* cordatum.

Rad. parasitica arborum. *Caul.* fruticosa, dichotoma. *Fol.* simplex, opposita. *Flor.* glomerata. *Fulcr.* o.

HIPPOPHAË. Mas. *Cal.* 2-part. *Cor.* o.

Foem. *Cal.* 2-fid. *Cor.* o. *Styl.* 1. *Bacca* 1-sperma.

Rad. lignosa, repens. *Caul.* arborea. *Fol.* simplex, sparsa. *Flor.* glomerata, sessilib. *Fulcr.* o.

MYRICA. Mas. *Amenti* squama lunata. *Cor.* o.

Foem. *Ament.* et *Cor.* o. *Styli* 2. *Bacca* 1-sperma.

Rad. lignosa. *Caul.* fruticosa. *Fol.* simplex, sparsa. *Flor.* varia. *Fulcr.* o.

Urtica dioica vide Monoecia Tetrandria.

5. PENTANDRIA.

CANNABIS. Mas. *Cal.* 5-part. *Cor.* o.

Foem. *Cal.* 1-phyll. integer, latere hians. *Cor.* o. *Styli* 2. *Nux* 2-valv. intra calycem clausum.

Rad. subramosa. *Caul.* herbacea. *Fol.* alterna, digitata. *Flor.* foeminea glomerata, mascula subracemosa. *Fulcr.* o.

HUMULUS. Mas. *Cal.* 5-phyll. *Cor.* o.

Foem. *Cal.* 1-phyll. integer, oblique patens. *Cor.* o. *Styli* o. *Sem.* 1. intra calycem foliatum.

Rad. repens. *Caul.* herbacea, volubilis. *Fol.* simplex, lobata. *Flor.* axillaris, glomerata. *Fulcr.* stipulae.

Salix pentandra vide Dioecia, Diandria.

H E X A N D R I A.

Rumex Acetosa et Acetosella vide Hexandria Trigynia.

6. O C T A N D R I A.

POPULUS. Mas. *Ament. Cal.* lamina lacera. *Cor* turbinate, obliqua, integra.

Foem. Ament. Cal. et Cor. maris. *Stigm.*

4-fid. *Capsf.* 2-locul. *Sem.* papposa.

Rad. lignosa. *Caul.* arborea. *Fol.* simplex, alternata. *Flor.* amentacea. *Fulcr.* o.

RHODIOLA. Mas. *Cal.* 4-part. *Cor.* 4-pet.

Foem. Cal. 4-part. *Cor.* o. *Nect.* 4. *Pist.* 4.

Capsf. 4.

Rad. carnosa, tuberosa. *Caul.* herbacea, simplex.

Fol. simplex, subsucculenta. *Flor.* cymosa.

Fulcr. o.

7. E N N E A N D R I A.

MERCURIALIS. Mas. *Cal.* 3-part. *Cor.* o. *Stam.* 9-12.

Antherae globosae didymae.

Foem. Cal. et Cor. maris. *Styli* 2. *Capsf.* 2-cocca. 2-locul.

Rad. varia, fibrosa. *Caul.* herbacea. *Fol.* simplex, opposita. *Flor.* axillaris, varia. *Fulcr.* stipulae.

HYDROCHARIS. Mas. *Spatha* 2-phylla. *Cal.* 3-fid.

Cor. 3-pet. *Filam.* 3 interiora stylifera.

Foem. Cal. et Cor. maris. *Styli* 6. *Capsf.* 6-locul.

Rad. fibrosa. *Caul.* repens, demersa. *Fol.* simplex, orbiculata, repens. *Flor.* longe pedunculata, emersa. *Fulcr.* bracteae.

D E C A N D R I A.

Lychnis dioica vide Decandria Pentagynia.

I C O S A N D R I A.

Rubus Chamaemorus vide Icosandria Polygynia.

8. POLYANDRIA.

STRATIOTES. Mas. *Spatha* 2-phyll. *Cal.* 3-part. *Cor.* 3-pet. *Nect.* 20. *Stam.* 12.

Foem. *Spatha Cal.* etc. ut in mare. *Germ.* 1. *Styl.* 6, bipartiti. *Bacca.*

Rad. fibrosa. *Caul.* scaposa. *Fol.* demersa, radicalis, simplex, crassa. *Flor.* terminalis. *Fulcr.* bracteae maris.

9. MONADELPHIA.

JUNIPERUS. Mas. *Ament.* *Cal.* squamae. *Cor.* 0. *Stam.* 3. Foem. *Cal.* 3-partit. *Cor.* 3-pet. *Styli* 3. *Bacca* 3-sperma.

Rad. lignosa. *Caul.* fruticosa. *Fol.* acerosa. *Flor.* axillaris, sessilis. *Fulcr.* 0.

TAXUS. Mas. *Cal.* 3-phyll. gemmae. *Stam.* multa. *Antherae* peltatae 8-fidae.

Foem. *Cal.* et *Cor.* maris. *Styl.* 0. *Sem.* 1. calyculo baccato integerrimo.

Rad. lignosa. *Caul.* arborea. *Fol.* acerosa, disticha. *Flor.* axillaris, pedunculata. *Fulcr.* 0.

XXII. DIOECIA.

1. MONANDRIA.

NAJAS.

1204. *N. marina*. Fl. Sv. 877. P. l. mar.

2. DIANDRIA.

VALLISNERIA.

1205. *V. spiralis*. Fl. L. 371. Fl. N. 1083. B. aqu.

SALIX.

* *foliis ferratis glabris*.

1206. *S. hermaphroditica*, foliis ferratis glabris, floribus hermaphroditis diandris. Fl. Sv. 878. Arb.

1207. *S. triandra*, foliis ferratis glabris, floribus triandris. S. N. 879. Hoffm. Sal. I. p. 45. t. 9. 10. t. 23. f. 2. F.

1208. *S. pentandra*, foliis ferratis glabris, floribus subpentandris. Fl. Sv. 879. Fl. N. 523. Fl. L. t. 8. f. 2. Fl. D. 943. F.

β. *S. polyandra*. Weig. Fl. 630. o).

1209. *S. phylicifolia*, foliis ferratis glabris lanceolatis, crenis undatis. Fl. Sv. 880. Fl. L. t. 8. f. D. Fl. N. 594. Arb.

β. sinibus crenarum nullis. Fl. L. t. 8. f. C.

1210. *S. Vitellina*, foliis ferratis ovatis acutis glabris, ferraturis cartilagineis, petiolis calloso-punctatis. S. N. 879. Weig. Fl. 619. Hoffm. Sal. I. p. 57. t. 11. 12. t. 24. f. 1. Arb.

1211. *S.*

o) Consentiente Nob. Weigel haec solum varietas est, licet constans *S. pentandrae*; nec numerus in hac adeo constans reperitur, ludit enim a quinario ad septenarium.

1211. *S. amygdalina*, foliis ferratis glabris lanceolatis petiolatis, stipulis trapeziformibus. Fl. Sv. 881. Fl. N. 821. Fr.
1212. *S. hastata*, foliis ferratis glabris lanceolatis subovatis acutis sessilibus, stipulis subcordatis. Fl. Sv. 882. Fl. L. t. 8. f. G. Fl. N. 597. F.
1213. *S. fragilis*, foliis ferratis glabris ovato-lanceolatis, petiolis dentato-glandulosis. Fl. Sv. 883. Fl. L. t. 8. f. B. Fl. N. 251. Arb.
1214. *S. purpurea*, foliis ferratis glabris lanceolatis: inferioribus oppositis, floribus monandris. Fl. Sv. 884. Weig. Fl. 622. *S. monandra* Hoffm. Sal. I. p. 18. t. 1. t. 5. f. 1. t. 23. f. 1. p). Fr.
1215. *S. myrsinites*, foliis ferratis glabris ovatis venosis. Fl. Sv. 885. Fl. L. t. 7. f. 6. t. 8. f. F. Fl. N. 595. Hoffm. Sal. I. p. 73. t. 17. 18. 19. 24. f. 2. F. alp.
1216. *S. arbuscula*, foliis ferratis glabris subdiaphanis subtus glaucis, caule suffruticoso. Fl. Sv. 886. Fl. N. 456. S. alp.
- α. foliis ovatis. Fl. L. 352. t. 8. f. E. Fl. D. 1052.
- β. foliis lanceolatis. Fl. L. 360. t. 8. f. M.
- γ. foliis integris. Fl. L. 356. Fl. D. 1055.
1217. *S. herbacea*, foliis ferratis glabris orbiculatis. Fl. Sv. 887. Fl. L. t. 8. f. H. t. 7. f. 3. 4. Fl. N. 111. Fl. D. 117. S. alp.

** *foliis integerrimis glabris.*

1218. *S. reticulata*, foliis integerrimis glabris ovatis obtusis. Fl. Sv. 888. Fl. N. 110. Fl. D. 212. Fl. L. t. 8. f. L. t. 7. f. 1. 2. Hoffm. Sal. II. p. 3. t. 25. 26. 27. S. alp.

P 5

1219. *S.*

p) *S. purpuream* et *Helicem* in unam speciem conjungit Nob. Hoffmann.

1219. *S. myrtilloides*, foliis integris glabris ovatis acutis.
Fl. Sv. 889. Fl. L. t. 8. f. I. K. Fl. N. 596. S. hum.
1220. *S. glauca*, foliis integerrimis subtus tenuissime
villosis ovato-oblongis. Fl. Sv. 890. Fl. L. t. 7.
f. 5. t. 8. f. P. Fl. N. 341. S. alp.

*** *foliis integerrimis villosis.*

1221. *S. aurita*, foliis integerrimis utrinque villosis, ob-
ovatis appendiculatis. Fl. Sv. 891. Fl. L. t. 8. f. Y.
Hoffm. Sal. I. p. 30. t. 4. f. 1. 2. t. 5. 3. t. 22. f. 1. F.
1222. *S. lanata*, foliis utrinque lanatis subrotundis acu-
tis. Fl. Sv. 892. Fl. L. t. 8. f. X. t. 7. f. 7. Fl. N.
431. *S. chrysanthos* Fl. D. 1057. F. alp.
β. *depressa*, foliis integris subtus villosis lan-
ceolato-ovatis utrinque acutis. Fl. Sv. 899.
Fl. L. t. 8. f. N. Fl. N. 991. F. alp.
1223. *S. Lapponum*, foliis integerrimis hirsutis lanceo-
latis. Fl. Sv. 893. Fl. L. t. 8. f. T. Fl. N. 626.
Fl. D. 1058. St. Helv. t. 14. F. alp.
1224. *S. arenaria*, foliis integris ovatis acutis supra
subvillosis, subtus tomentosis. Fl. Sv. 894. Fl. N.
439. Fl. D. 197. F. pr. sylv.
1225. *S. incubacea*, foliis integerrimis lanceolatis subtus
villosis nitidis, stipulis ovatis acutis. Fl. Sv. 895.
Fl. N. 533. F. pr.
1226. *S. repens*, foliis integerrimis lanceolatis utrinque
nudiusculis, caule repente. Fl. Sv. 896. Fl. N. 989.
S. depressa Hoffm. Sal. I. p. 63. t. 15. 16. S. pr. pal.
1227. *S. fusca*, foliis integerrimis ovatis subtus pube-
scentibus. Fl. Sv. 897. Fl. L. t. 8. f. R. Fl. N. 990.
Hoffm. Sal. II. p. 7. t. 28. 29.? S. pr. hum.
1228. *S. rosmarinifolia*, foliis integerrimis lanceolato-
linearibus strictis sessilibus subtus tomentosis. Fl.
Sv. 898. Fl. N. 820. S. pr. pal.

**** *foliis subserratis villosis.*

1229. *S. caprea*, foliis ovatis rugosis subtus tomentosis undatis superne denticulatis. Fl. Sv. 900. Fl. N. 252. Fl. D. 245. Hoffm. Sal. I. p. 25. t. 3. t. 5. f. 4. t. 21. f. 1. Arb. pr. n.

α. foliis omnibus crenatis. Fl. L. t. 8. f. S.

β. foliis inferioribus integris. Fl. L. t. 8. f. V.

1230. *S. viminalis*, foliis subintegerrimis lanceolato-linearibus longissimis acutis subtus sericeis, ramis virgatis. Fl. Sv. 901. Hoffm. Sal. I. p. 22. t. 2. t. 5. f. 2. t. 21. f. 2. Arb. f.

1231. *S. cinerea*, foliis subserratis oblongo-ovatis subtus subvillosis, stipulis dimidiato-cordatis. Fl. Sv. 902. Fl. N. 946. F. hum.

1232. *S. alba*, foliis lanceolatis acuminatis serratis utrique pubescentibus: serraturis infimis glandulosis. Fl. Sv. 903. Hoffm. Sal. I. p. 41. t. 7. 8. t. 24. f. 3.

3. TRIANDRIA.

EMPETRUM.

1233. *E. nigrum*, procumbens. Fl. Sv. 904. Fl. N. 127. Fl. D. 975. S. ster.

4. TETRANDRIA.

VISCUM.

† 1234. *V. album*, foliis lanceolatis obtusis, caule dichotomo, spicis axillaribus. Fl. Sv. 905. Fl. N. 444. F. sylv.

HIPPOPHAE.

1235. *H. Rhamnoides*, foliis lanceolatis. Fl. Sv. 906. Fl. N. 11. Fl. D. 265. Arb. l. mar.

MYRICA.

236 DIOECIA. TETRANDRIA. Myrica.

MYRICA.

1236. *M. Gale*, foliis lanceolatis subferratis, caule suffruticoso. Fl. Sv. 907. Fl. N. 142. Fl. D. 327. F. pal.

5. PENTANDRIA.

CANNABIS.

1237. *C. sativa*, S.N. 886. Weig. Fl. 633. A. c. r.

HUMULUS.

1238. *H. Lupulus*. Fl. Sv. 908. Fl. N. 358. P. f.

6. OCTANDRIA.

POPULUS.

1239. *P. alba*, foliis subrotundis dentato angulatis subtus tomentosis. Fl. Sv. 911. Arb.

1240. *P. tremula*, foliis subrotundis dentato angulatis utrinque glabris, petiolis compressis. Fl. Sv. 909. Fl. N. 137. Arb. nem. sylv.

1241. *P. nigra*, foliis deltoidibus acuminatis ferratis. Fl. Sv. 910. Arb.

RHODIOLA.

1242. *R. rosea*. Fl. Sv. 912. Fl. N. 103. Fl. D. 183. P. alp.

7. ENNEANDRIA.

MERCURIALIS.

1243. *M. perennis*, caule simplicissimo, foliis scabris. Fl. Sv. 913. Fl. D. 400. P. nem.

1244. *M. annua*, caule brachiato, foliis glabris, floribus spicatis. S.N. 890. Weig. Fl. 639. A. c.

HYDROCHARIS.

1245. *H. Morfus Ranae*. Fl. Sv. 914. Fl. D. 878. P. qu. lac.

8. POLY-

DIOECIA. POLYANDRIA. Stratiotes. 237

8. POLYANDRIA.

STRATIOTES.

1246. *S. Aloides*, foliis ensiformi-triangularis ciliato-aculeatis. Fl. Sv. 479. Fl. N. 653. Fl. D. 337. P. qu.

9. MONADELPHIA.

JUNIPERUS.

1247. *J. communis*, foliis ternis patentibus mucronatis bacca longioribus. Fl. Sv. 915. Fl. N. 280. F.
 a. minor,
 β. major, arborescens.

TAXUS.

1248. *T. baccata*, foliis approximatis. Fl. Sv. 916. Fl. N. 420. Arb. fylv.

XXIII. POLYGAMIA.

I. MONOECIA.

VERATRUM. Herm. Cal. o. Cor. 6 pet. Stam. 6. Pist. 3.
Caps. 3. polyspermae.

Mas. Pistillorum rudimenta.

Rad. ramosa, crassa. Caul. herbacea. Fol. simplex, nervosa, plicata. Flor. racemosa, terminalis. Fulcr. bracteae.

HOLCUS. Herm. Cal. gluma 1-2-flora. Cor. aristata.
Stam. 3. Styli 2. Sem. 1.

Mas. Cal. gluma 2-valv. Cor. o. Stam. 3.
Rad. fibrosa. Caul. culmosa, articulata. Fol. graminea. Flor. paniculata. Fulcr. o.

Avena elatior vide Triandria Digynia.

Rottboellia vide Triandria Digynia.

VALANTIA. Herm. Cal. o. Cor. 4-part. Stam. 4. Styl. 2-fid. Sem. 1.

Mas. Cal. o. Cor. 3-4 partit. Stam. 3. 4.
Pistill. obsolet.

Rad. fibrosa. Caul. herbacea, angulata, geniculata. Fol. verticillata. Flor. axillaris. Fulcr. o.

PARIETARIA. Herm. Cal. 4-fid. Cor. o. Stam. 4.
Styl. 1. Sem. 1. elongat. superum.

Foem. Cal. etc. Hermaphroditi. Stam. o.
Rad. ramosa. Caul. herbacea. Fol. simplex, integerrima. Flor. axillaris, conferta. Fulcr. o.

ATRIPLEX. Herm. Cal. 5-phyll. Cor. o. Stam. 5.
Stylus 2-part. Sem. 1. depressum.

Foem. Cal. 2-phyll. Cor. o. Stam. o.
Rad. ramosa. Caul. herbacea (praeter 1. speciem).
Fol. simplex, succulenta, farinosa. Flor. glomerata. Fulcr. o.

ACER.

ACER. Herm. *Cal.* 5-fid. *Cor.* 5-pet. *Stam.* 8. *Pist.* 1.
Capf. 2. alatae.

Mas. *Pistillum* o.

Rad. lignosa. *Caul.* arborea, opposita ramosa.

Fol. subpalmata. *Flor.* varia. *Fulcr.* stipulae
 caducae.

2. DIOECIA.

FRAXINUS. Herm. *Cal.* o. *Cor.* o. *Stam.* 2. *Pist.* 1.
 compressum. *Sem.* 1. lanceolatum.

Foem. *Stam.* o.

Rad. lignosa. *Caul.* arborea, ramis oppositis.

Fol. pinnata. *Flor.* racemosa. *Fulcr.* o.

Rhamnus catharticus vide Pentandria Monogynia.

Ilex Aquifolium vide Tetrandria Tetragynia.

Cucubalus Otites vide Decandria Trigynia.

XXIII. POLYGAMIA.

I. M O N O E C I A.

VERATRUM.

1249. *V. album*, racemo supradecomposito, corollis erectis. S. N. 902. Fl. N. 315. t. 1. P. alp.

HOLCUS.

1250. *H. mollis*, glumis bifloris nudiusculis, flosculo hermaphrodito mutico: masculo arista geniculata. S. N. 905. Schr. Gr. t. 20. f. 2. P. pr. sicc.
 1251. *H. lanatus*, glumis bifloris villosis, flosculo hermaphrodito mutico: masculo arista recurva. Fl. Sv. 917. Fl. N. 378. Fl. D. 963. P. ar. hum.
 1252. *H. odoratus*, glumis trifloris muticis acuminatis, flosculo hermaphrodito diandro. Fl. Sv. 918. Fl. N. 934. P. nem. hum.

VALANTIA.

- † 1253. *V. cruciata*, floribus masculis quadrifidis, pedunculis diphyllis. S. N. 908. Weig. Fl. 644. A. r.

PARIETARIA.

- † 1254. *P. officinalis*, foliis lanceolato-ovatis, pedunculis dichotomis, calycibus diphyllis. S. N. 908. Weig. Fl. 645. Fl. D. 521. P. r.

ATRIPLEX.

1255. *A. Portulacoides*, caule fruticoso, foliis obovatis. Fl. Sv. 919. S. l. m.
 1256. *A. laciniata*, caule herbaceo, foliis deltoideis dentatis subtus argenteis. Fl. Sv. 920. Fl. N. 1011. A. l. m.
 1257. *A. hastata*, caule herbaceo, calycum valvulis foemineis magnis deltoideis sinuatis. Fl. Sv. 921. Fl. N. 1012. A. c. r.

1258. *A.*

POLYGAMIA. MONOECIA. Acer. 241

1258. *A. patula*, caule herbaceo patulo, foliis subdeltoideo-lanceolatis, calycibus feminum disco dentatis. Fl. Sv. 922. Fl. N. 489. A. c. r.
1259. *A. littoralis*, caule herbaceo erecto, foliis omnibus linearibus integerrimis. Fl. Sv. 923. Fl. N. 725. A. l. m.
1260. *A. marina*, caule herbaceo erecto, foliis linearibus serratis. Mant. Pl. p. 300. Fl. N. 1107. A. l. m.
1261. *A. pedunculata*, caule herbaceo divaricato, foliis lanceolatis obtusis integris, calycibus foemineis pedunculatis. S. N. 910. Obs. Bot. Fasc. I. p. 31. Fl. D. 304. A. l. m.

ACER.

- † 1262. *A. Pseudoplatanus*, foliis quinquelobis inaequaliter serratis, floribus racemosis. S. N. 911. Weig. Fl. 650. Arb.
1263. *A. Platanoides*, foliis quinquelobis acuminatis acute dentatis glabris, floribus corymbosis. Fl. Sv. 924. Fl. N. 614. Arb.
1264. *A. campestre*, foliis lobatis obtusis emarginatis. Fl. Sv. 925. Fl. N. 637. F. sylv.

2. D I O E C I A.

FRAXINUS.

1265. *F. excelsior*, foliis serratis, floribus apetalis. Fl. Sv. 926. Fl. N. 423. Fl. D. 969. Arb.

XXIV. CRYPTOGAMIA.

I. FILICES.

EQUISETUM. *Spica* fructificationibus peltatis, basi multivalvi dehiscentibus.

Rad. repens. *Caul.* articulata, teres, striata.

Fol. o. *Flor.* spicata. *Fulcr.* o.

OPHIOGLOSSUM. *Spica* articulata, disticha, articulis transversim dehiscentibus.

Rad. repens. *Caul.* o. *Fol.* radicalis, simplicissima, solitaria. *Fruſt.* spicata, epiphylla. *Fulcr.* o.

OSMUNDA. *Spica* ramosa fructificationibus globosis.

Rad. fibrosa. *Caul.* simplex vel ramosa. *Fol.* pinnata. *Flor.* pinnata. *Fulcr.* o.

ACROSTICHUM. *Fructificationes* discum totum frondis tegentes.

Rad. nodosa. *Caul.* simplex. *Fol.* varia. *Flor.* expansa. *Fulcr.* o.

PTERIS. *Fructificationes* in linea marginem cingente.

Rad. nodosa, repens. *Caul.* ramosa. *Fol.* composita. *Flor.* marginalis. *Fulcr.* o.

ASPLENIUM. *Fructificat.* in lineolis disci frondis sparsis.

Rad. fibrosa. *Caul.* varia. *Fol.* varia. *Flor.* lineata. *Fulcr.* o.

POLYPODIUM. *Fructificat.* in punctis subrotundis sparsis per discum frondis.

Rad. varia. *Caul.* *Fol.* varia. *Flor.* punctata. *Fulcr.* o.

TRICHOMANES. *Fructificat.* solitariae, stylo setaceo terminatae, margini ipsi frondis insertae.

Rad. filiformis, repens. *Caul.* simplex. *Fol.* composita. *Fruſt.* solitaria. *Fulcr.* o.

PILULARIA. *Fructificat.* ad radicem, globus quadrilobularis.

Rad. fibrosa, repens. *Caul.* o. *Fol.* simplicissima, filiformis, inundata. *Flor.* radicalis. *Fulcr.* o.

ISOETES.

ISOËTES. *Frustrificat.* ad radicem intra basin frondis
2-locularis.

Rad. fibrosa. *Caul.* o. *Fol.* simplicissima, canaliculata, submersa. *Flor.* radicalis. *Fulcr.* o.

A M U S C I.

LYCOPODIUM. *Anthera* 2-valvis, sessilis. *Calyptra* o.
Rad. fibrosa. *Caul.* ramosa. *Fol.* imbricata, acerosa. *Flor.* sessilis, subspicata. *Fulcr.* o.

SPHAGNUM. *Anthera* operculata ore laevi. *Calyptra* o.
Rad. fibrosa. *Caul.* ramosa. *Fol.* imbricata.
Flor. breviter pedunculata. *Fulcr.* o.

PHASCUM. *Anthera* operculata ore ciliato. *Calyptra*
fugax.
Rad. fibrosa. *Caul.* o. *Fol.* radicalis. *Flor.* solitaria. *Fulcr.* o.

FONTINALIS. *Anthera* operculata, calyptrata, sessilis,
perichaetio inclusa.
Rad. fibrosa. *Caul.* ramosa, demersa. *Fol.* acerosa. *Flor.* sessilis. *Fulcr.* o.

BUXBAUMIA. *Anthera* operculata, hinc membranacea,
intra operculum sacculus pollinis. *Calyptra* caduca.
Rad. fibrosa. *Caul.* simplicissima. *Fol.* o. l. capsulam cingens. *Flor.* varia. *Fulcr.* o.

SPLACHNUM. *Anthera* insidens apophysi maximae coloratae. *Calyptra* caduca. *Stella* foeminea in distincto individuo.

Rad. fibrosa. *Caul.* rarius ramosa. *Fol.* acerosa.
Flor. pedunculata. *Fulcr.* apophysis.

POLYTRICHUM. *Anthera* operculata insidens apophysi minimae. *Calyptra* villosa. *Stella* foeminea distincta.

Rad. fibrosa. *Caul.* simplex l. ramosa. *Fol.* acerosa. *Flor.* pedunculata plerisque. *Fulcr.* o.

MINIUM. *Anthera* operculata. *Calyptra* laevis. *Foem.* capitulum nudum pulverulentum remotum.

Rad. etc. praecedentis.

BRYUM. *Anthera* operculata. *Calyptra* laevis. *Filam.* e tuberculo terminali ortum.

Rad. etc. praecedentis.

HYPNUM. *Anthera* *) operculata. *Calyptra* laevis. *Filamentum* laterale ortum e perichaetio.

Rad. etc. praecedentis.

3. ALGAE.

JUNGERMANNIA. *Anthera*, pedunculata quadrifida, nuda, e perichaetio tubuloso.

Rad. fibrosa. *Caul.* et *Fol.* varia. *Flor.* peltata.

Fulcr. stipulae foliiformes in nonnullis.

TARGIONIA. *Cal.* 2-valvis includens globum.

Rad. fibrillosa. *Caul.* o. *Fol.* simplicissima, sublanceolata. *Flor.* sessilis. *Fulcr.* o.

MARCHANTIA. *Mas.* *Cal.* peltatus subtus tectus corollis 1-petalis *Antherae* multifidae.

Foem. *Cal.* campanulatus, sessilis, polyspermus.

Rad. fibrosa. *Caul.* o. *Fol.* sinuosa, radicata.

Flor. sessilis et pedunculata. *Fulcr.* o.

BLASIA. *Mas.* *Anthera* subglobosa frondi immersa.

Foem. *Cal.* lagenaeformis. *Sem.* nuda per tubulum exeuntia.

Rad. etc. fere praecedentis.

RICCIA.

*) *Capsula* rectius dicitur haec Muscorum perperam nominata anthera.

RICCIA. *Granula* frondi innata in quibus *Cal. et Cor. o.*
Anthera cylindrica sessilis *germini* turbina-
 to imposita. *Stylus* filiformis anthe-
 ram perforans. *Pericarpium* sphaericum
 coronatum anthera emarcida. *Sem.* hemi-
 sphaerica pedicellata.

Rad. etc. fere antecedentis.

ANTHOCEROS. *Mas. Cal.* sessilis, cylindricus, integer.
Anthera subulata, longissima, 2-valvis.

Foem. Cal. 6-partit. *Sem.* 3.

Rad. etc. praecedentium.

LICHEN. *Receptacula* subrotunda, planiuscula, nitida.
Habitus diversissimus, subcoriaceus.

TREMELLA. *Frustrificatio* vix manifesta in corpore ge-
 latinoso.

Habitus rugosus, tremulus, auriformis, in
 humido tumidus.

FUCUS. *Frustrificatio* vesiculae cavae.

Habitus cartilagineus vel potius pergamenae
 similis, radicatione submersa marina.

ULVA. *Frustrificatio* in membrana diaphana.

Habitus vesicularis, tenuis.

CONFERVA. *Tubercula* inaequalia in fibris capillaribus
 longissimis.

Habitus filamentosus.

CORALLINA. *Stirps* dichotoma, articulata crusta cal-
 carea obducta.

Habitus filiformis lithophyti.

BYSSUS. *Lanugo* aut *Pulvis* simplicissimus.

Habitus capillaceus vel pulverulentus.

4. FUNGI.

AGARICUS. Fungus horizontalis subtus lamellosus.

Habitus spongiosus, fragilis.

BOLETUS. Fungus horizontalis subtus porosus.

Habitus inconstans, subpulvinatus.

- HYDNUM.** Fungus horizontalis subtus echinatus.
Habitus varius.
- PHALLUS.** Fungus subtus laevis, supra reticulatus.
Habitus pistillaris.
- CLATHRUS.** Fungus cancellatus.
Habitus diversus.
- HELVELLA.** Fungus turbinatus.
Habitus difformis plicatus.
- PEZIZA.** Fungus campanulatus vel cochleatus sessilis.
Habitus subcoriaceus patelliformis.
- CLAVARIA.** Fungus laevis oblongus.
Habitus simplex subcylindricus vel multum ramosus.
- LYCOPERDON.** Fungus subrotundus seminibus farinaceis repletus.
Habitus globosus.
- MUCOR.** Fungus vesiculis stipitatis.
Habitus diversus.
- LACHNUM.** Fungus undique hirtus.
Habitus Agarici.

XXIV. CRYPTOGAMIA.

I. FILICES.

EQUISETUM.

1266. *E. sylvaticum*, caule spicato, frondibus compo-
sitis. Fl. Sv. 927. Fl. N. 494. P. sylv.
1267. *E. arvense*, scapo fructificante nudo: sterili fron-
doso: ramis subdenis verticillatis tetragonis. *E. ar-*
vense Fl. Sv. 928. Fl. N. 363. P. pr. agr.
1268. *E. Telmateja*, scapo fructificante nudo: sterili
frondoso: ramis subquadrigenis verticillatis octo-
fulcatis. *E. Telmateja* Ehrh. Beytr. 2. p. 159. St.
Helv. 1675. P. pal. ul.
1269. *E. palustre*, caule angulato, frondibus simplici-
bus. Fl. Sv. 929. Fl. N. 410. P. aqu.
β, *polystachyon*. Raji Syn. 3. t. 5. f. 3.
1270. *E. fluviatile*, caule tereti striato subsimplici.
Fl. Sv. 930. Fl. N. 384. γ). P. aqu.
1271. *E. limosum*, caule subnudo laevi. Fl. Sv. 931.
Fl. N. 383. γ). P. pal.
1272. *E. hyemale*, caule basi subramoso fulcato, punctis
seriatis exasperato, vaginularum denticulis obso-
letis. *Equisetum hyemale* Fl. Sv. 931. Fl. N. 276.
P. pal. sylv.

OPHIOGLOSSUM.

1273. *O. vulgatum*, fronde ovata. Fl. Sv. 932. Fl. N.
693. Fl. D. 147. P. pr. sylv.

Q 4

OSMUN.

- γ) Varietas *E. fluviatile* Ehrh. Beytr. 1. p. 63.
r) Ramulorum fulcos ac dentes 4, 5, 6 numeravi, an ideo *E.*
Heleocharis Ehrh. Beytr. 2. p. 159. vere diversa?

OSMUNDA.

1274. *O. Lunaria*, scapo caulino solitario, fronde pinnata solitaria. Fl. Sv. 934. Fl. N. 185. Fl. D. 18. P. sylv.

β. *spica gemina*.

γ. *mutica*. Fl. D. 18. fig. min.

δ. *matricariaefolia*. Fl. D. 18. fig. supr.

1275. *O. regalis*, frondibus bipinnatis apice racemiferis. Fl. Sv. 932. Fl. N. 992. Fl. D. 217. P. sylv. hum. marg. fluv.

1276. *O. Struthiopteris*, frondibus pinnatis: pinnis pinnatifidis, scapo fructificante disticho. Fl. Sv. 935. Fl. N. 1. t. 1. f. 1-3. Fl. D. 169. P. sylv.

1277. *O. Spicant*, frondibus lanceolatis pinnatifidis, laciniis confluentibus integerrimis parallelis. Fl. Sv. 936. Fl. N. 213. Fl. D. 99. P. sylv. hum.

1278. *O. crispus*, frondibus supradecompositis: pinnis alternis subrotundis incis. S. N. 928. N. Act. Svec. 1789. p. 228. Fl. N. 1055. Fl. D. 496. P. alp.

ACROSTICHUM.

1279. *A. septentrionale*, frondibus nudis linearibus laciniatis. Fl. Sv. 937. Fl. N. 206. Fl. D. 60. P. mont.

1280. *A. hyperboreum*, fronde pinnata, pinnis rotundatis 3-5-fidis. N. Act. Svec. 1793. P. alp.

1281. *A. ilvense*, frondibus subbipinnatis: pinnis opposito coadunatis obtusis subtus hirsutis basi integerrimis. Fl. Sv. 938. Fl. N. 635. Fl. D. 391. P. mont.

PTERIS.

1282. *P. aquilina*, frondibus supradecompositis, foliis pinnatis pinnis lanceolatis: infimis pinnatifidis: superioribus minoribus. Fl. Sv. 939. Fl. N. 30. P. sylv.

ASPLE.

ASPLENIUM.

1283. *A. Scolopendrium*, frondibus simplicibus cordato-lingulatis integerrimis, stipitibus hirsutis. Fl. N. 941. P. sylv.
1284. *A. Trichomanes*, frondibus pinnatis, pinnis subrotundis crenatis. Fl. Sv. 942. Fl. N. 281. Fl. D. 119. P. mont.
1285. *A. Breynii*, frondibus subbipinnatis, foliolis oblongulatis alternis apice ferrulatis. Prodr. Fl. Scand. 1155. Obs. Bot. Fasc. I. 111. *A. Germanicum* Weiff. Crypt. p. 299.
1286. *A. Ruta muraria*, frondibus alternatim decompositis: foliolis cuneiformibus crenulatis. Fl. Sv. 943. Fl. N. 282. Fl. D. 190. P. muri.
1287. *A. Adiantum nigrum*, frondibus subtripinnatis, foliis alternis: pinnis lanceolatis inciso-ferratis. S. N. 934. Act. Svec. 1769, p. 248. Fl. D. 250. P. sylv.

POLYPODIUM.

1288. *P. vulgare*, frondibus pinnatifidis: pinnis oblongis subferratis obtusis, radice squamata. Fl. Sv. 944. Fl. N. 184. P. arb. lap.
1289. *P. Lonchitis*, frondibus pinnatis: pinnis lunulatis, ciliato-ferratis declinatis, stipitibus strigosis. S. N. 936. Fl. N. 49. Act. Nidros. 4. t. 5. f. 2. 3. Fl. D. 497. P. mont.
- † 1290. *P. fontanum*, frondibus pinnatis lanceolatis, foliolis subrotundis argute incis, stipite laevi. S. N. 937. Koem. Fl. Isl. II. P.
- † 1291. *P. montanum*, frondibus subbipinnatis: pinnis alternis, pinnulis integerrimis lanceolatis obtusiusculis, glomerulis marginalibus. Wild. Fl. Berol. 883. *P. Oreopteris* Ehrh. s). P. sylv. hum.

Q 5

1292, P.

s) Cel. Vahl.

250 CRYPTOGRAMIA. FILICES. Polypodium.

1292. *P. Phegopteris*, frondibus subbipinnatis, foliolis infimis reflexis: paribus pinnula quadrangulati coadunatis. Fl. Sv. 945. Fl. N. 50. P. sylv.
1293. *P. cristatum*, frondibus subbipinnatis, foliolis ovato-oblongis: pinnis obtusiusculis apice acute ferratis. S. N. 937. N. Act. Svec. 1787. p. 256. t. 9. *P. Callipteris* Ehrh. Beytr. III. p. 77. P. pal.
1294. *P. Filix mas*, frondibus bipinnatis: pinnis obtusis crenulatis, stipite paleaceo. Fl. Sv. 946. Fl. N. 4. t. 1. f. 4. N. Act. Svec. 1787. p. 257. P. sylv.
1295. *P. spinulosum*, frondibus bipinnatis: pinnis pinnatifidis ferratis, ferraturis spinulosis. *P. spinulosum* Fl. D. 707. *P. Filix foemina spinosa* Weis Crypt. p. 316. Müll. Fridr. 841. t. 2. f. 2. P. pr. sylv.
1296. *P. molle*, frondibus subbipinnatis, foliolis lanceolatis incisissimis ferratis basi pinnatifidis, ferraturis acutis spinulosis. *P. molle* Schreb. Lips. p. 70. *P. Filix foemina crenata* Weis Crypt. p. 313. f. Moris. III. f. 14. t. 3. f. 7. P. pr. sylv.
1297. *P. Filix foemina*, fronde bipinnata: pinnulis linearibus lanceolatis pinnatifidis acutis. Fl. Sv. 947. Fl. N. 29. Plukn. Phyt. t. 180. f. 4. Moris. III. f. 14. t. 3. f. 8. P. sylv.
1298. *P. Thelypteris*, fronde bipinnata: pinnis pinnatifidis integerrimis subtus undique polline tectis. S. N. 937. Fl. N. 625. Fl. D. 760. *Acrostichum Thelypteris* Fl. Sv. 939. P. sylv. pal.
- † 1299. *P. aculeatum*, frondibus bipinnatis: pinnis lunulatis ciliato-dentatis, stipite strigoso. S. N. 937. Fl. N. 976. P. sylv.

1300. *P.*

*) Descriptio Cel. Weiß haud bene in omnibus respondet nostrae filici, quae tamen illa Morisoni et Schreberi est. Num haec sit *P. molle* Ehrh. Pl. Crypt. adhuc ignoro. A *Polypodie molli* Jacq. ic. rar. certe diversissima.

CRYPTOGAMIA. FILICES. Trichomanes. 251

1300. *P. fragile*, frondibus bipinnatis: foliolis remotis: pinnis subrotundis incisfis. Fl. Sv. 948, Fl. N. 341, Fl. D. 401. P. lap. muri.
 1301. *P. Dryopteris*, frondibus supradecompositis: foliolis ternis bipinnatis, Fl. Sv. 949, Fl. N. 31. Fl. D. 759. P. sylv.

TRICHOMANES.

- † 1302. *T. tunbrigense*, frondibus pinnatis: pinnis oblongis dichotomis decurrentibus dentatis. S. N. 941. Nat. H. Selfk. Skr. II. p. 68, Fl. D. 954. v). P. alp.

PILULARIA.

1303. *P. globulifera*. Fl. Sv. 950. Fl. D. 223. Vaill. Par. t. 15, f. 6. P. inund.

ISOËTES.

1304. *I. lacustris*, Fl. Sv. 951. it. Scan. 417, c. icone. Fl. N. 607, Fl. D. 191. v). P. fl. lac.

M U S C I.

LYCOPODIUM.

1305. *L. clavatum*, foliis sparsis filamentosis, spicis tetretibus pedunculatis geminis. Fl. Sv. 952. Fl. N. 224, Fl. D. 126. P. sylv.
 1306. *L. Selaginoides*, foliis sparsis ciliatis lanceolatis, spicis solitariis terminalibus foliosis. Fl. Sv. 953. Fl. N. 226, Fl. D. 70. P. sylv. pal.
 1307. *L.*

u) Trichomanis genus certe in plura divelli patitur.

v) Utramque Dillenii speciem Calamariae in Norvegia observavit *Cel. Müller*. cfr. Reysse til Telemark p. 40. In nostris lacubus quoque occurrunt. An specie vel varietate tantum diversae? *Coromandeliana* quam misit opt. *König* foliis longis fere filiformibus c. fig. 2. tab. 80. *Dill. Musc. convenit*.

252 CRYPTOGRAMIA. MUSCI. Sphagnum.

1307. *L. isundatum*, foliis sparsis integerrimis, spicis terminalibus foliosis. Fl. Sv. 954. Fl. D. 336. P. l. lac.
1308. *L. Selago*, foliis sparsis octofariis, caule dichotomo erecto fastigiato, floribus sparsis. Fl. Sv. 955. Fl. N. 205. Fl. D. 104. P. sylv. acer.
1309. *L. annotinum*, foliis sparsis quinquefariis subserratis, furculis annotino-articulatis, spicis terminalibus glabris erectis. Fl. Sv. 956. Fl. N. 225. Fl. D. 127. P. sylv. acer.
1310. *L. alpinum*, foliis quadrifariam imbricatis acutis, caulibus erectis bifidis, spicis sessilibus teretibus. Fl. Sv. 957. Fl. N. 272. Fl. D. 79. P. alp.
1311. *L. complanatum*, foliis bifariis connatis: superficialibus solitariis, spicis geminis pedunculatis. Fl. Sv. 958. Fl. N. 271. Fl. D. 78. P. sylv.
- † 1312. *L. dubium*, furculis simplicissimis erectis compressis, foliis complicatis carinatis acutis alternis distiche imbricatis. Fl. Isl. II. w).

SPHAGNUM.

1313. *S. palustre*, ramis deflexis. Fl. Sv. 959. Fl. N. 202. Fl. D. 474. P. pal.

PHASCUM.

1314. *P. acaulon*, acaule, capsula sessili, foliis ovatis acutis. Fl. Sv. 960. Fl. D. 249. f. sin. *Ph. cuspidatum* Schreb. Phasc. 3. t. 1. f. 1, 2. Fl. N. 509. terra nuda.
1315. *P. muticum*, acaule, capsula sessili, foliis ovatis muticis concavis conniventibus. Schreb. Phasc. 4. t. 1. f. 11, 12. terra nuda.

1316. *P.*

w) De genere dubius ex habitu huc retulit D. König.

1316. *P. subulatum*, acaule, capsula sessili, foliis subulato-setaceis patulis. S. N. 944. Schreb. Phasc. I. t. 1. f. 15-17. Fl. D. 249. f. supr. terra nuda.

FONTINALIS.

1317. *F. antipyretica*, foliis complicato-carinatis trifariis acutis, capsulis lateralibus. Fl. Sv. 961. Fl. N. 988. Dill. Musc. t. 33. f. 1. fluv.
 1318. *F. minor*, foliis ovatis concavis trifariis acutis passim geminis, capsulis terminalibus. S. N. 945. Fl. N. 969. t. 3. f. 2. Dill. Musc. t. 33. f. 2. fluv.
 1319. *F. squamosa*, foliis imbricatis subulato-lanceolatis, capsulis lateralibus. Hedw. Crypt. III. 2. p. 32. t. 12. x). fluv.
 1320. *F. capillacea*, foliis lineari-setaceis. Fl. Sv. 962. Ehrh. Beytr. 3. p. 158. Dill. Musc. t. 33. f. 5. fluv.
 1321. *F. secunda*, ramosa, capsulis secundis lateralibus. N. Aët. Ups. p. 243. *Sphagnum arboreum* S. N. 944. Fl. N. 1107. Dill. Musc. t. 32. f. 6. Arb.

BUXBAUMIA.

1322. *B. aphylla*, aphylla. Fl. Sv. 1293. Fl. N. 571. Fl. D. 44. Am. Acad. 5. t. 1. Schmied. Diff. p. 25. t. 1. f. 1-25. Dill. Musc. t. 68. terra glar.
 1323. *B. foliosa*, acaulis, capsula subsessili foliis cincta. Am. Acad. 10. p. 112. t. 4. f. 4. *Phascum Halleri* Prodr. Fl. Scand. 1185. Müll. Fridr. 854. Aët. Svec. 1764. t. 2. f. 6-10. Hall. St. Helv. 1725. t. 46. f. 3. *Buxbaumia sessilis* Schmied. Diff. p. 26. t. 1. f. 1-14. terra hum.

SPLACHNUM.

1324. *S. rubrum*, umbraculo orbiculari hemisphaerico. Fl. Sv. 963. Amoen. Acad. 2. t. 3. f. 2. Fl. N. 381. Dill. Musc. t. 83. f. 9. pal.

1325. *S.*

x) *Cel. Acharius.*

254 CRYPTOGRAMIA. MUSCI. Polytrichum.

1325. *S. luteum*, umbraculo orbiculari plano. Fl. Sv. 964. Am. Acad. 2. t. 3. f. 1. pal.
1326. *S. ampullaceum*, umbraculo ampullaceo obovato. Fl. Sv. 965. Fl. N. 798. Fl. D. 822. Dill. Musc. t. 44. f. 3. pal.
1327. *S. vasculosum*, umbraculo ampullaceo subgloboso. S. N. 945. Fl. D. 192. Fl. Isl. p. 11. pal.
1328. *S. angustatum*, caulescens, foliis piliferis, pedunculo brevissimo. Amoen. Acad. X. p. 113. *S. Bryoides* Prodr. Fl. Scand. 1193. Fl. Isl. p. 12. pal. alp.
1329. *S. mnioides*, subacaule, receptaculo oblongo. Amoen. Acad. X. p. 99. y). pal. alp.
1330. *S. sphaericum*, receptaculo globoso. Am. Acad. X. p. 113. t. 4. f. 1. N. Act. Upf. 4. p. 240. pal. alp.

POLYTRICHUM.

1331. *P. commune*, caule simplici, capsula parallelipeda, foliis ferrulatis acutis. Schr. Lips. 73. Fl. Sv. 966. α. β. Fl. N. 203. pr. sylv. hum.
α. *ferrulatum* Dill. Musc. t. 54. f. 1.
β. *ciliatum*. Dill. Musc. t. 54. f. 2. Fl. D. 295.
1332. *P. piliferum*, caule simplici, capsula parallelipeda, foliis integerrimis piliferis. Schr. Lips. 74. Fl. Sv. 966. γ. Dill. Musc. t. 54. f. 3. pr. sylv. sicc.
1333. *P. alpinum*, caule ramosissimo, pedunculis terminalibus. S. N. 946. Dill. Musc. t. 55. f. 4.
P. ramosum Fl. N. 814. Fl. D. 297. pal. alp.
1334. *P. urnigerum*, caule ramosissimo, pedunculis axillaribus. Fl. Sv. 967. Fl. N. 704. Dill. Musc. t. 55. f. 5. Fl. D. 296. pr. pal. alp.

MNIUM.

y) Speciem non vidi, at si diversa, certe ad illam non pertinet fig. 192. Fl. Dan. quae *S. vasculosum* refert. At *Cel. Afzelius* in Diff. p. 34. contendit *vasculosum* et *mnioides* varietates ex aetate esse.

MNIUM.

1335. *M. pellucidum*, caule simplici, foliis ovatis. Fl. Sv. 968. Fl. N. 817. Fl. D. 300. Dill. Musc. t. 31. f. 2. hum.
1336. *M. androgynum*, caule ramoso, androgynum. Fl. Sv. 971. Fl. N. 816. Fl. D. 299. Dill. Musc. t. 31. f. 1. pal.
1337. *M. fontanum*, caule simplici geniculis inflexo. Fl. Sv. 969. Fl. N. 673. Fl. D. 298. Dill. Musc. t. 44. f. 2. aqu.
1338. *M. palustre*, caule dichotomo, foliis subulatis. Fl. Sv. 970. Fl. N. 682. Dill. Musc. t. 31. f. 3. pal.
1339. *M. hygrometricum*, acaule, capsula nutante, calyptra reflexa tetragona. Fl. Sv. 972. Fl. N. 464. Fl. D. 648. f. 2. Dill. Musc. t. 52. f. 75. terra nuda.
1340. *M. purpureum*, caule dichotomo, axillis pedunculiferis, capsula erecta, foliis carinatis. Fl. Sv. 973. Fl. N. 662. Dill. Musc. t. 49. f. 51. pr.
1341. *M. setaceum*, capsulis erectis, operculis filiformibus longitudine capsularum. Fl. Sv. 974. Dill. Musc. t. 48. f. 47. lign. aqu.
1342. *M. cirrhatum*, foliis arefactione revolutis. Fl. Sv. 975. Fl. D. 538. f. 4. Dill. Musc. t. 48. f. 42. sylv.
1343. *M. annotinum*, foliis ovatis acuminatis pellucidis, pedunculis subradicalibus, capsulis nutantibus. Fl. Sv. 976. Fl. N. 336. Dill. Musc. t. 50. f. 65. hum.
1344. *M. hornum*, capsulis pendulis, pedunculo curvato, furculo simplici, foliis margine scabris. Fl. Sv. 977. Dill. Musc. t. 51. f. 71. sylv.
1345. *M. capillaceum*, capsulis erectiusculis oblongis, operculo conico, foliis capillaceis. N. Act. Upf. 4. p. 241. Dill. Musc. t. 49. f. 57. ficc. alp.

1346. *M. capillare*, capsulis pendulis, foliis ovatis fetiferis carinatis, pedunculis longissimis. Fl. Sv. 982. Dill. Musc. t. 59. f. 67. sicc. lap.
1347. *M. crudum*, capsulis pendulis, calyptra recurvata, foliis pellucidis. Fl. Sv. 979. Fl. N. 674. Dill. Musc. t. 51. f. 70. ul.
1348. *M. pyriforme*, capsulis pendulis turbinatis, stipite filiformi, rosularum foliis fetiferis. Fl. Sv. 980. Dill. Musc. t. 50. f. 60. hum.
1349. *M. Polytrichoides*, calyptra villosa. Fl. Sv. 983. Dill. Musc. t. 55. f. 6. eric. hum.
 β . Dill. Musc. t. 55. f. 7. et 12.
1350. *M. Serpyllifolium*, pedunculis aggregatis, foliis patentibus pellucidis. Fl. Sv. 982. Fl. N. 661. hum.
 α . *punctatum*, foliis integris obtusis punctatis. Dill. Musc. t. 53. f. 81.
 β . *cuspidatum*, foliis serratis acutis. Dill. Musc. t. 53. f. 79.
 γ . *proliferum*, foliis rosaceo-congestis lanceolatis acutis. Dill. Musc. t. 52. f. 77.
 δ . *undulatum*, foliis oblongis undulatis. Dill. Musc. t. 52. f. 76.
1351. *M. triquetrum*, foliis trifariis patulis subulato-lanceolatis carinatis. Fl. Sv. 978. Fl. N. 1102. pal.
 α . Dill. Musc. t. 51. f. 72.
 β . *ibid.* f. 73.
 γ . *ibid.* f. 74. x).

BRYUM,

x) Negat huc pertinere icones Dill, Cl. Ehrhart Beytr. 1. p. 137. affirmat Cel. Hudson.

BRYUM.

1352. *B. apocarpum*, capsulis sessilibus terminalibus, calyptra minima. Fl. Sv. 986. Fl. N. 997. lap. arb.

α. aureum. Dill. Musc. t. 32. f. 5.

β. coccineum. Fl. D. 480. Dill. Musc. t. 32. f. 4.

1353. *B. striatum*, capsulis sessilibus sparsis, calyptris striatis sursumve pilosis. Fl. Sv. 987. Fl. N. 558. arb. lap. mr.

α. minus. Dill. Musc. t. 55. f. 9.

β. majus. ibid. f. 8. Fl. D. t. 537. f. 3.

γ. tenerius. ibid. f. 10.

δ. capsula subsessili. ibid. f. 11. *Polytrichum arboreum* Fl. D. 648. f. 1.

1354. *B. pomiforme*, capsulis erectis sphaericis. Fl. Sv. 988. Fl. N. 1008. Dill. Musc. t. 44. f. 1. lap.

1355. *B. pyriforme*, capsulis erectis obovatis, calyptra subulata acaulibus, foliis ovatis muticis. Fl. Sv. 989. Fl. N. 561. Fl. D. 537. f. 1. Dill. Musc. t. 44. f. 6. pr.

1356. *B. exstinctorium*, capsula erecta oblonga minore, calyptra laxa aequali. Fl. Sv. 990. Fl. N. 947. Fl. D. 1001. f. 2. arv. mr.

α. minus. Dill. Musc. t. 45. f. 8.

β. majus. ibid. f. 9.

1357. *B. subulatum*, capsulis erectis subulatis, furculis acaulibus. Fl. Sv. 991. Fl. D. 1000. f. 2. Dill. Musc. t. 45. f. 10. hum.

1358. *B. rurale*, capsulis erectiusculis, foliis piliferis recurvis. Fl. Sv. 992. Fl. N. 576. Dill. Musc. t. 45. f. 12. test. mr. arb.

1359. *B. murale*, capsulis erectis, foliis piliferis erectiusculis, furculis simplicibus caespitosus. Fl. Sv. 993. Dill. Musc. t. 45. f. 14.

1360. *B. scoparium*, capsulis rectiusculis, pedunculis aggregatis, foliis secundis recurvatis, caule declinato. Fl. Sv. 995. Fl. D. 821. f. 1. Dill. Musc. t. 46. f. 16. eric. sylv.
1361. *B. undulatum*, capsulis erectiusculis, pedunculis subsolitariis; foliis lanceolatis carinatis undulatis patentibus serratis. Fl. Sv. 996. Fl. D. 477. Dill. Musc. t. 46. f. 18. pr.
1362. *B. glaucum*, capsulis erectiusculis: operculo arcuato, foliis erectis imbricatis, furculis ramosis. Fl. Sv. 994. Fl. D. 824. f. 2. Dill. Musc. t. 46. f. 20. eric.
1363. *B. pellucidum*, capsulis erectiusculis, foliis recurvis acutis, caule hirsuto. Fl. Sv. 997. Dill. Musc. t. 46. f. 23. pal.
1364. *B. flexuosum*, capsulis erectis, foliis setaceis, pedunculis flexuosis. Fl. Sv. 998. Dill. Musc. t. 47. f. 33. sylv.
1365. *B. unguiculatum*, capsulis erectis oblongis, pedunculis axillaribus, furculis in summitate crassioribus. S. N. 948. Dill. Musc. t. 48. f. 47. mr.
1366. *B. capillaceum*, capsulis erectis oblongis, pedunculis terminalibus, foliis setaceis patulis decurrentibus. Dickf. Fasc. 4. t. 1. f. 6. Fl. D. 1000. f. 1. alp.
1367. *B. aciculare*, capsulis erectis: operculo aciformi, foliis erectis subsecundis. S. N. 948. Fl. D. 1000. f. 1. Dill. Musc. t. 46. f. 26. mont. hum.
1368. *B. heteromallum*, capsulis erectis, foliis setaceis secundis. Fl. Sv. 999. Fl. D. 479. Dill. Musc. t. 47. f. 37. eric. pasc.
1369. *B. tortuosum*, capsulis erectis, foliis setaceis imberbibus: arefactione retortis. Fl. Sv. 1000. Fl. D. 880. f. 2. Dill. Musc. t. 48. f. 40. mont.

1370. *B. truncatulum*, capsulis erectis subrotundis: operculis mucronatis. Fl. Sv. 1001. Fl. N. 781. Fl. D. 537. f. 2. Dill. Musc. t. 45. f. 7. nem. pr. mr.
1371. *B. viridulum*, capsulis erectis ovatis, foliis lanceolatis acuminatis imbricato-patulis. Fl. Sv. 1002. Fl. N. 660. Dill. Musc. t. 48. f. 43. pr. sylv.
1372. *B. paludosum*, acaule, foliis setaceis, capsulis obtusissimis patulis. Fl. Sv. 1004. Dill. Musc. t. 49. f. 53. pr. pal.
1373. *B. laterale*, capsulis erectis ovalibus, pedunculis brevissimis lateralibus, foliis setaceis subsecundis. N. Act. Upf. 4. p. 241. Fl. D. 538. f. 3. *Bryum Norvegicum* Fl. N. 1064. Prodr. Fl. Scand. 1245. alp. subhum.
1374. *B. hypnoides*, capsulis erectis, furculo rectiusculo, ramis lateralibus brevibus fertilibus. Fl. Sv. 1003. Fl. N. 559. lap.
 α. *hirsutum*. Dill. Musc. t. 47. f. 32.
 β. *pilosum*. ibid. f. 27.
 γ. *virescens*. ibid. f. 28.
 δ. *microcarpos*. ibid. f. 29. Fl. D. 476.
 ε. *obtusum*. ibid. f. 30.
 ζ. *barbatum*. ibid. f. 31.
1375. *B. verticillatum*, capsulis erectis, pedunculis arefactione confertis, foliis piliferis, furculis fastigiatis. S. N. 949. Dill. Musc. t. 47. f. 35. mont. hum.
1376. *B. aestivum*, capsulis erectis subrotundis axillariibus, foliis subulatis distantibus. S. N. 949. Dill. Musc. t. 47. f. 39. mont. hum.
1377. *B. Celsii*, capsulis erectiusculis, pedunculis longissimis, foliis setaceis, furculis acaulibus. Fl. Sv. 1005. Fl. N. 574. Dill. Musc. t. 49. f. 54. terr.
1378. *B. trichodes*, capsulis erectiusculis: ore ciliato absque annulo, pedunculo longissimo. Fl. Sv. 1006. Dill. Musc. t. 49. f. 58. pal.

1379. *B. squarrosum*, capsulis oblongis, foliis quinquefariis imbricatis recurvis, caule ferrugineo tomentoso. Fl. Sv. 1007. Buxbaum. Cent. 4. t. 35. f. 1. pal. turf.
1380. *B. argenteum*, capsulis pendulis, furculis cythricis imbricatis laevibus. Fl. Sv. 1008. Fl. D. 880. f. 1. Dill. Musc. t. 50. f. 62. test. mr. lap.
1381. *B. pulvinatum*, capsulis subrotundis, pedunculis reflexis, foliis piliferis. Fl. Sv. 1009. Dill. Musc. t. 50. f. 65. mr. lap.
- † 1382. *B. pendulum*, capsulis pendulis, furculis procumbentibus, pedunculo brevissimo, foliis setaceis flexuosis. Fl. D. 823. f. 1. mont. hum.
1383. *B. bimum*, capsulis pendulis ovato-oblongis obtusis, pedunculis longis, foliis lanceolatis alternis. N. Act. Ups. 4. p. 242. Schreb. Lips. 1047. Dill. Musc. t. 51. f. 73. pal.
1384. *B. caespitium*, capsulis pendulis, foliis lanceolatis acuminato-setaceis, pedunculis longissimis. Fl. Sv. 1010. Fl. N. 570. Dill. Musc. t. 50. f. 66. test. mr.
1385. *B. virens*, capsulis pendulis incurvis obliquis, foliis lanceolatis acutis carinatis. N. A. Ups. 4. 241. Dill. Musc. t. 50. f. 61. ? mont.
1386. *B. carneum*, capsulis pendulis subglobosis, foliis acutis alternis. Fl. Sv. 1011. Dill. Musc. t. 50. f. 69. ul.
1387. *B. simplex*, capsula nutante oblonga, foliis subulatis, furculo simplicissimo: medio pedunculifero. S. N. 949. Dill. Musc. t. 50. f. 59. ar.
1388. *B. alpinum*, capsula pendula oblonga, foliis ovatis acuminatis carinatis, furculis ramosis, axillis pedunculiferis. S. N. 949. Mant. alt. p. 309. Fl. N. 1063. Dill. Musc. t. 50. f. 64. alp.
- † 1389. *B. hyperboreum*, capsulis ovalibus erectis, pedunculis brevibus, foliis capillaribus. Fl. N. 1060. Fl. D. 538. f. 1. alp.

† 1390. *B.*

- † 1390. *B. Schisti*, capsulis erectis obtuse ovatis: capitulo oblique rostrato, apophysi capitulo subiecta, foliis setaceis curvatis. Fl. N. 1061. Fl. D. 538. f. 2. terr. schist.
- † 1391. *B. Oederi*, capsulis sphaericis subnutantibus maturitate striatis, foliis striatis acuminatis reflexis. Fl. N. 1005. Fl. D. 478. alp. hum.

HYPNUM.

* *frondibus pinnatis.*

1392. *H. taxifolium*, fronde simplicissima patente lanceolata, basi pedunculifera. Fl. Sv. 1013. Fl. D. 473. f. 2. Dill. Musc. t. 34. f. 2. sylv.
1393. *H. denticulatum*, fronde pinnata simplici: pinnis duplicatis, basi pedunculifera. Fl. Sv. 1014. Dill. Musc. t. 34. f. 5. sylv.
1394. *H. Bryoides*, fronde simplicissima pinnata lanceolata, apice pedunculifera. Fl. Sv. 1012. Fl. D. 473. f. 1. Dill. Musc. t. 34. f. 1. sylv.
1395. *H. Adianthoides*, fronde pinnata ramosa erecta medio pedunculifera. S. N. 950. Act. Holm. 1769. p. 248. Dill. Musc. t. 34. f. 3. lap. pal.
1396. *H. Trichomanoides*, fronde pinnata ramosa: foliis distichis imbricatis acinaciformibus obtusis. Schr. Lips. 88. N. Act. Ups. 4. p. 242. Dill. Musc. t. 34. f. 8. arb.
1397. *H. complanatum*, fronde pinnata ramosa, foliis imbricatis acutis complicatis compressis. Fl. Sv. 1015. Dill. Musc. t. 34. f. 7. arb.
1398. *H. sylvaticum*, fronde pinnata ramosa procumbente, foliolis acutis, medio pedunculifera. S. N. 950. Dill. Musc. t. 34. f. 6. sylv.

** *furculis vagis.*

1399. *H. lucens*, furculis ramosis, frondibus subpinnatis: foliolis punctatis. S. N. 950. Dill. Musc. t. 34. f. 10. sylv. ul.

1400. *H. rugosum*, furculis vagis erectiusculis, foliis secundis recurvatis basi rugosis. S. N. 950. Mant. Fl. p. 131. Dill. Musc. t. 37. f. 24. montos.
1401. *H. undulatum*, furculis ramosis, frondibus subpinnatis, foliis undulatis complicatis. S. N. 950. Dill. Musc. t. 36. f. 11. arb.
1402. *H. crispum*, furculis ramosis, frondibus subpinnatis, foliis undulatis planis. S. N. 950. Dill. Musc. t. 36. f. 12. sylv. eric.
1403. *H. triquetrum*, ramis vagis recurvis, foliis ovatis recurvatis patulis. Fl. Sv. 1016. Fl. N. 871. Dill. Musc. t. 38. f. 28. sylv.
1404. *H. rutabulum*, ramis vagis subrepentibus, foliis ovatis mucronatis imbricatis. S. N. 950. Fl. D. 824. f. 2. Dill. Musc. t. 38. f. 29. 30. 31. sylv.

*** *furculis pennatis.*

1405. *H. filicinum*, furculis pennatis: ramulis distantibus, foliis imbricatis incurvis acutis secundis. Fl. Sv. 1017. Dill. Musc. t. 26. f. 19. pr. hum.
1406. *H. proliferum*, furculis proliferis plano-pinnatis, pedunculis aggregatis. Fl. Sv. 1018. Fl. N. 373. Dill. Musc. t. 35. f. 14. pr.
1407. *H. delicatulum*, furculis subproliferis plano-pinnatis cuspidatis, pedunculis aggregatis. Fl. Sv. 1019. Fl. N. 865. Dill. Musc. t. 83. f. 6. pr.
1408. *H. parietinum*, furculis plano-pinnatis continuatis, pedunculis aggregatis. Fl. Sv. 1020. Fl. N. 306. Dill. Musc. t. 35. f. 13. pasc. pr.
1409. *H. praelongum*, furculis subpinnatis decumbentibus, ramulis remotis, foliis ovatis, capsulis cernuis. Fl. Sv. 1021. Dill. Musc. t. 35. f. 15. pr. arb.
1410. *H. Crista castrensis*, furculis pinnatis, ramulis approximatis apicibus recurvis. Fl. Sv. 1022. Fl. N. 867. Dill. Musc. t. 36. f. 20. sylv.

1411. *H.*

1411. *H. abietinum*, furculis pinnatis teretiusculis remotis inaequalibus. Fl. Sv. 1023. Fl. N. 866. Dill. Musc. t. 35. f. 17. sylv.

1412. *H. plumosum*, furculis pinnatis repentibus, ramis confertis, foliis imbricatis subulatis, capsulis erectis. S. N. 951. Dill. Musc. t. 35. f. 16. arb. terr.

**** *foliis recurvis.*

1413. *H. cupressiforme*, furculis subpinnatis, foliis secundis recurvis apice subulatis. Fl. Sv. 1024. Fl. N. 868. Fl. D. 535. f. 2. Dill. Musc. t. 37. f. 23. sylv.

1414. *H. aduncum*, furculis erectiusculo-subramosis, foliis secundis recurvis subulatis, ramulis recurvatis. Fl. Sv. 1025. Dill. Musc. t. 37. f. 26. pr. hum.

1415. *H. Scorpioides*, ramis vagis procumbentibus recurvis, foliis secundis acuminatis. Fl. Sv. 1026. Dill. Musc. t. 37. f. 25. a). pal.

1416. *H. viticulosum*, furculis repentibus, ramis vagis teretibus, foliis patulis acuminatis. S. N. 951. Dill. Musc. t. 39. f. 43. pr. arb.

1417. *H. aquaticum*, furculis procumbentibus submersis, foliis linearibus acuminatis recurvis secundis, capsulis oblongis. S. N. 951. N. Act. Upf. 4. p. 243. Jacqu. Fl. Austr. t. 291. Dill. Musc. t. 43. f. 70. lap. sub. aqu.

1418. *H. squarrosum*, ramis vagis, foliis ovatis apice subulatis undique reflexis. Fl. Sv. 1027. Fl. N. 870. ul.

α. majus. Dill. Musc. t. 39. f. 38.

β. minus. ibid. f. 39. Fl. D. 535. f. 1. 648. f. 3. ? b).

1419.

a) Varietates tantum esse *H. Scorpioides* et *rugosum* cum Webro contendit Cl. Afzelius Dill. cit. p. 34.

b) Folia compilato-carinata quinquefariam imbricata quae in S. N. huic tribuuntur, ego nondum vidi. Figuræ Fl. D. diversas species depingere videntur.

1419. *H. palustre*, furculis repentibus, ramis confertis, foliis ovatis secundis, capsulis erectiusculis. S. N. 951. Dill. Musc. t. 37. f. 27. pal.

1420. *H. loreum*, furculis reptantibus, ramis vagis erectis, foliis secundis, capsulis subrotundis. S. N. 951. Dill. Musc. t. 39. f. 40. mont.

***** *furculis dendroidibus vel fasciculatis.*

1421. *H. dendroides*, furculo erecto, ramis fasciculatis terminalibus simpliciusculis, capsulis erectis. Fl. Sv. 1028. Fl. N. 580. Fl. D. 823. Dill. Musc. t. 40. f. 48. pr. pasc.

1422. *H. alopecurum*, furculo erecto, ramis fasciculatis terminalibus subdivisis, capsulis subnutantibus. Fl. Sv. 1029. Fl. N. 863. Dill. Musc. t. 41. f. 49. sylv.

***** *furculis teretiusculis.*

1423. *H. curtispendulum*, furculis vagis teretibus, foliis ovatis acutis patulis, capsulis pendulis. Fl. Sv. 1030. Dill. Musc. t. 43. f. 69. pasc.

1424. *H. purum*, furculis pinnato-sparsis subulatis, foliis ovatis obtusis conniventibus. Fl. Sv. 1031. Fl. N. 862. Fl. D. 706. f. 2. Dill. Musc. t. 40. f. 45. pasc.

1425. *H. illecebrum*, furculis ramisque vagis teretibus erectiusculis obtusis. Fl. Sv. 1032. Fl. N. 861. Fl. D. 706. f. 1. Dill. Musc. t. 40. f. 46. terr.

1426. *H. riparium*, furculis teretibus ramosis, foliis acutis patulis distantibus. Fl. Sv. 1033. Fl. D. 649. f. 1. Dill. Musc. t. 40. f. 44. aqu.

1427. *H. cuspidatum*, furculis vagis; apice foliis convolutis acuminatis. Fl. Sv. 1035. sylv. pal.

a. pungens. Weis Crypt. 251. Dill. Musc. t. 39. f. 34.

β. inerme. Weis Crypt. 253. Dill. Musc. t. 40. f. 47.

***** *fur-*

***** *furculis confertis.*

1428. *H. sericeum*, furculo repente, ramis confertis erectis, foliis subulatis, capsulis erectis. Fl. Sv. 1036. Fl. N. 860. mr. terr. arb.
α. curvatum. Weis Crypt. 254. Dill. Musc. t. 42. f. 59.
β. praelongum. Weis Crypt. 255. Dill. Musc. t. 42. f. 60.
1429. *H. velutinum*, furculo repente, ramis confertis erectis, foliis subulatis, capsulis subnutantibus. Fl. Sv. 1037. Fl. D. 475. Dill. Musc. t. 42. f. 61. arb.
1430. *H. serpens*, furculis repentibus, ramis filiformibus, foliis oblitteratis. Sp. Pl. 1596. Dill. Musc. t. 42. f. 64. lap. terr. trunci hum.
1431. *H. sciuroides*, furculis erectis ramosis incurvatis. Sp. Pl. 1596. Dill. Musc. t. 41. f. 54. arb.
1432. *H. gracile*, furculis repentibus, ramis fasciculatis teretibus erectiusculis, capsulis erectis ovatis. S. N. 952. Dill. Musc. t. 41. f. 55. *H. ornithopodioides* Fl. D. 649. f. 2. arb. lap.
1433. *H. myosuroides*, furculis ramosissimis, ramis subulatis utrinque attenuatis teretibus. S. N. 952. Dill. Musc. t. 41. f. 51. mr. lap.
1434. *H. clavellatum*, repens, ramis erectis confertissimis, capsulis incurvis: operculis inflexis. S. N. 952. Dill. Musc. t. 85. f. 17. arb.
1435. *H. Halleri*, ramulis approximatis brevibus, foliis planis recurvatis. S. N. 952. alp.
1436. *H. nitens*, ramis brevissimis, foliis lanceolatis subulatis nitentibus. S. N. 952. N. Act. Upf. 4. p. 242. Dill. Musc. t. 39. f. 37. ul.

3. ALGAE.

JUNGERMANNIA.

* *frondibus pinnatis secundis.*

1437. *J. Sphagni*, frondibus reptantibus simpliciter pinnatis latere floriferis, foliolis subrotundis integerrimis imbricatis secundis. Dickf. Crypt. 1. p. 6. t. 1. f. 10. c). musc.
1438. *J. asplenoides*, frondibus simpliciter pinnatis, foliis pinnatis ovatis subciliatis. S. N. 953. Fl. D. 1061. Dill. Musc. t. 69. f. 5. sylv.
1439. *J. viticulosa*, frondibus simpliciter pinnatis lanceolatis apice floriferis, foliis integerrimis. S. N. 953. Fl. Fr. 928. Dill. Musc. t. 70. f. 10. terr. hum.
1440. *J. polyanthos*, frondibus simpliciter pinnatis, foliis integerrimis imbricatis convexis. S. N. 953. Dill. Musc. t. 69. f. 7. terr. hum.
1441. *J. lanceolata*, frondibus simpliciter pinnatis lanceolatis apice floriferis, foliis integerrimis. S. N. 953. Fl. Fr. 928. Dill. Musc. t. 70. f. 10. terr. hum.
1442. *J. bidentata*, frondibus simpliciter pinnatis apice floriferis, foliis bidentatis. S. N. 953. Dill. Musc. t. 70. f. 11. arb. lap.
1443. *J. bicuspidata*, frondibus simpliciter pinnatis medio floriferis, foliis bidentatis. S. N. 953. Fl. Fr. 930. Dill. Musc. t. 70. f. 13. arb.
1444. *J. fissa*, frondibus simpliciter pinnatis, foliolis bifidis, capsulis radicalibus. N. Act. Ups. 4. p. 244. *J. bicornis* Fl. D. 888. *J. globulifera* Poll. Fl. III. n. 1061. *Minium fissum* S. N. 947. terr. hum.

1445. *J.*o) *Cel. Acharius.*

1445. *J. quinquedentata*, frondibus pinnatis ramosis apice floriferis, foliolis quinquedentatis. S. N. 953. Dill. Musc. t. 71. f. 23. terr. ul.

** *frondibus pinnatis: pinnulis auriculatis superne testis.*

1446. *J. nemorea* frondibus supra bipinnatis apice floriferis, foliis ciliatis. S. N. 953. Fl. N. 675. Dill. Musc. t. 71. f. 18. sylv.

1447. *J. resupinata*, frondibus supra bipinnatis inferne floriferis, foliis crenulatis imbricatis rotundis, Fl. Sv. 1038. Fl. N. 676. Dill. Musc. t. 71. f. 19. lap.

1448. *J. albicans*, frondibus supra bipinnatis apice floriferis, foliis linearibus recurvatis. Fl. Sv. 1039. Dill. Musc. t. 71. f. 20. terr.

1449. *J. trilobata*, frondibus subtus bipinnatis, foliis quadratis obsolete trilobis, Fl. Sv. 1040. Dill. Musc. t. 71. f. 22. terr.

1450. *J. reptans*, frondibus subtus bipinnatis apice reptantibus, foliis quadridentatis. S. N. 954. Fl. Fr. 932. Dill. Musc. t. 71. f. 24. terr.

1451. *J. multiflora*, fronde repente ramosa, foliolis alternis geminis setaceis aequalibus. S. N. 954. Dill. Musc. t. 69. f. 4. terr.

1452. *J. purpurea*, pinnis subtus auriculatis, foliolis subrotundis convexo - concavis amplexicaulibus, capsulis terminalibus. N. Act. Upf. 4. p. 244. *Mnium Jungermannia* Fl. Sv. 984. rup. aqu.

*** *frondibus imbricatis.*

1453. *J. complanata*, furculis repentibus, foliis inferne auriculatis duplicato - imbricatis, ramis aequalibus. Fl. Sv. 1041. Fl. N. 521. Fl. D. 1062. Dill. Musc. t. 72. f. 26. arb.

1454. *J. dilatata*, furculis repentibus, foliis inferne auriculatis duplicato - imbricatis, ramis apice latioribus, Fl. Sv. 1042. Dill. Musc. t. 72. f. 27. arb.

1455. *J.*

1455. *J. tamariscifolia*, foliis imbricatis serie duplici: superioribus subrotundis convexis obtusis quadruplo majoribus. Fl. Sv. 1043. Fl. N. 857. Dill. Musc. t. 72. f. 31. arb. lap.

1456. *J. platyphylla*, furculis procumbentibus subtus imbricatis, foliis cordatis acutis. S. N. 954. Fl. Fr. 936. Dill. Musc. t. 72. f. 32. arb.

1457. *J. ciliaris*, furculis repentibus, foliis duplicato-imbricatis: inferne auriculatis ciliatis. Fl. Sv. 1044. Fl. N. 855. d) lap.

1458. *J. tomentella*, furculis repentibus foliisque integris tomentosis imbricatis, vaginis tomentosis. *J. tomentella* Ehrh. Beytr. 2. p. 150. terr.

1459. *J. varia*, furculis erectiusculis bifariam imbricatis, foliis bipartitis. S. N. 954. Fl. Fr. 937. Dill. Musc. t. 73. f. 36. terr.

1460. *J. pulcherrima*, furculis erectiusculis confertis, foliis ovatis imbricatis bifidis ciliatis. N. Act. Ups. 4. p. 244. Dill. Musc. t. 69. f. 3. arb.

**** *furculis undique imbricatis: foliolis sparsis.*

1461. *J. julacea*, furculis teretibus erectis, foliis undique imbricatis bidentatis, floribus pedunculatis. S. N. 954. N. Act. Holm. 1789. p. 50. Fl. D. 1002. f. 2. Dill. Musc. t. 73. f. 38. alp.

1462. *J. cincinnata*, furculis teretibus subcompressis dense imbricatis compactis, foliis convexo-concavis glabris, floribus pedunculatis. Lightf. Fl. Scot. 2. p. 786. Dill. Musc. t. 73. f. 38. e) rup.

1463. *J. rupestris*, furculis teretibus, foliis subulatis secundis. Fl. Sv. 1045. Fl. N. 869. Dill. Musc. t. 73. f. 40. lap.

1464.

d) Cl. Ehrhart hanc cum *J. pulcherrima* eandem facit, quo jure dijudicare nequeo, at Dill. icon. t. 73. f. 35. merito ad *tomentellam* suam refert. Cel. Dickson *ciliarem* et *pulcherrimam* distinguit.

e) Cel. Acharius.

1464. *J. trichophylla*, furculis teretibus, foliis capilla-
ceis aequalibus. Fl. Sv. 1046. Fl. N. 859. Dill.
Musc. t. 73. f. 37. lap.
1465. *J. alpina*, furculis teretibus, foliis ovatis paten-
tibus, calycibus imbricatis. S. N. 954. Fl. N. 858.
Fl. D. 1002. f. 1. Dill. Musc. t. 73. f. 39. lap.
1466. *J. Sertularioides*, furculis repentibus, ramulis
undique floriferis, foliis capillaceis aequalibus ar-
ticulatis. Am. Acad. X. p. 116. t. 4. f. 6. N. Aët.
Upf. 4. p. 243. eric. ul.
- ***** *acaules, frondibus simplicibus.*
1467. *J. epiphylla*, acaulis, foliolo frondi innato. Fl.
Sv. 1047. Fl. N. 338. Fl. D. 359. Dill. Musc. t. 74.
f. 41. ul.
1468. *J. pinguis*, acaulis, fronde oblonga sinuata pingui.
Fl. Sv. 1048. Dill. Musc. t. 73. f. 42. pal.
1469. *J. multifida*, acaulis, fronde bipinnatifida. S. N.
955. Dill. Musc. t. 74. f. 43. terr. ul.
1470. *J. furcata*, acaulis, fronde lineari-ramosa: ex-
tremitatibus obtusiusculis furcatis. Fl. Sv. 1049.
Fl. N. 856. Dill. Musc. t. 74. f. 45. lap. lign.
1471. *J. pusilla*, acaulis, fronde subpinnatifida: lobis
imbricatis, perichaetio plicato. Fl. Sv. 1050. Dill.
Musc. t. 74. f. 46. lap.

TARGIONIA.

1472. *T. hypophylla*. S. N. 955. Fl. Isl. 14. Dill. Musc.
t. 78. f. 9. terr.

MARCHANTIA.

1473. *M. polymorpha*, calyce communi decemfido. Fl.
Sv. 1051. Fl. N. 19. hum. umbr.
- α. *stellata*, radiis teretibus. Dill. Musc. t. 76.
f. 6. E. F. t. 77. f. 7. B. C.
- β. *umbellata*, calyce octocrenato. Dill. Musc.
t. 76. f. 6. A. C. D. t. 77. f. 7. D.

1474. *M. hemisphaerica*, calyce communi quinquefido hemisphaerico, perichaetio nullo. Fl. Sv. 1052. Fl. N. 506. Fl. D. 762. Dill. Musc. t. 75. f. 2. ul.
1475. *M. conica*, calyce communi quinqueloculari subovato. S. N. 955. Fl. N. 808. Fl. D. 274. Dill. Musc. t. 75. f. 1. ul.
1476. *M. tenella*, calyce communi hemisphaerico quinqueloculari subovato, margine radiato. S. N. 955. Fl. Isl. 14. Dill. Musc. t. 75. f. 4. ul.

BLASIA.

1477. *B. pusilla*. Fl. Sv. 1053. Fl. N. 787. Fl. D. 45. Fl. Isl. 14. Dill. Musc. t. 31. f. 7. hum. a.

RICCIA.

1478. *R. crustallina*, frondibus superficiei pappillofis. Fl. Sv. 1055. Fl. N. 873. Dill. Musc. t. 78. f. 12. pr. hum.
1479. *R. minima*, frondibus glabris bipartitis acutis. Fl. Sv. 1054. Dill. Musc. t. 78. f. 11. pasc. hum.
1480. *R. glauca*, frondibus glabris canaliculatis bilobis obtusis. S. N. 956. Dill. Musc. t. 78. f. 10. a. hum.
1481. *R. fluitans*, frondibus dichotomis lineari-filiformibus. Fl. Sv. 1056. Fl. D. 275. Dill. Musc. t. 74. f. 47. f). foss. caesp.
- † 1482. *R. natans*, frondibus obcordatis ciliatis. S. N. 956. Dill. Musc. t. 78. f. 18. g). stagn.

ANTHOCEROS.

1483. *A. punctatus*, frondibus indivisis sinuatis punctatis. S. N. 956. Fl. N. 872. Fl. D. 396. Dill. Musc. t. 68. f. 1. ul.

1484. *A*

f) Fructificatio globosa sessilis, evalvis, polysperma in latere inferiore frondium. *Ehrh. Beyr. 3. p. 81.*

g) Cel. Kölpin.

1484. *A. laevis*, frondibus indivisis sinuatis laevibus.
S. N. 956. Fl. N. 874. Dill. Musc. t. 68. f. 2. ul.

LICHEN.

* *leprosi tuberculati*.

1485. *L. scriptus*, leprosus albicans, lineolis nigris ramosis characteriformibus. Fl. Sv. 1057. Fl. N. 758.
Dill. Musc. t. 18. f. 1. cort. arb.
1486. *L. geographicus*, leprosus flavescent, lineolis nigris mappam referens. Fl. Sv. 1058. Fl. N. 177.
Fl. D. 168. f. 1. Dill. Musc. t. 18. f. 5. lap.
1487. *L. atrovirens*, leprosus viridis, margine tuberculisque atris. Fl. Sv. 1062. h). *Verrucaria atrovirens* Hoffm. Lich. I. p. 83. t. 17. f. 4. lap.
1488. *L. lacteus*, leprosus albus, tuberculis concoloribus hemisphaericis. M. Pl. 132. Fl. N. 877. lap.
1489. *L. rupicola*, leprosus albidus, tuberculis pallidis albo-marginatis. M. Pl. 132. *Verrucaria contorta* Hoffm. Lich. I. p. 97. t. 22. f. 1-4. ? lap.
1490. *L. pertusus*, leprosus, verrucis subtesselatis laevigatis pertusis poro uno alterove cylindrico. M. Pl. 131. Fl. D. 766. Dill. Musc. t. 18. f. 9. arb. lap.
1491. *L. rugosus*, leprosus albicans, lineolis simplicibus punctisque nigris confertis. Fl. Sv. 1059. Dill. Musc. t. 18. f. 2. arb.
1492. *L. sanguinarius*, leprosus cinereo-virescent tuberculis atris. Fl. Sv. 1060. *Verruc. sanguinaria* Hoffm. Lich. I. p. 58. t. 41. f. 1. arb. alp.
1493. *L. Oderi*, leprosus, crusta albo-glauescente, rimis in maculas subrotundas divisa: tuberculis atris in lineas curvas dispositis. Fl. N. 1025. Fl. D. 472. f. 3. i). rup. mar. 1494.

h) Eundem cum praecedente statuit *Cel. Afzelius* Diff. cit. p. 35.

i) Nomina huic et sequenti a Rev. *Gunnaro* primum imposita, in priore hujus Prodr. editione allata, etiam nunc retinui.

1494. *L. Koenigii*, leprofus ruber, tuberculis nigris. Fl. N. 1020. Fl. D. 470. f. 1. Fl. Isl. 15. *L. Oederi* S. N. 957. *Patellaria filacea* Hoffm. Lich. I. p. 89. t. 19. f. 2. 3. rup.
1495. *L. fusco-ater*, leprofus fuscus, tuberculis atris. Fl. Sv. 1063. lap.
1496. *L. vernalis*, leprofus albidus, tuberculis subrotundis ferrugineis. S. N. 957. Dill. Musc. t. 18. f. 4. arb.
1497. *L. calcareus*, leprofus candidus, tuberculis atris. Fl. Sv. 1061. Fl. N. 942. Dill. Musc. t. 18. f. 8. lap.
1498. *L. cinereus*, leprofus cinereus, tuberculis nigris albo marginatis. M. Pl. 132. lap.
1499. *L. atro-albus*, leprofus niger, tuberculis atroalboque mistis. Fl. Sv. 1064. lap.
1500. *L. niger*, leprofus niger, tuberculis concoloribus. N. Act. Upf. 4. p. 245. k). lap. terr.
1501. *L. confluens*, leprofus albicans, tuberculis globosis magnis atris aetate confluentibus. N. A. Upf. 4. p. 245. Web. Spic. t. 1. *Verruc. confluens* Hoffm. Lich. I. p. 87. t. 19. f. 1. rup.
1502. *L. terminalis*, crustaceus niger, tuberculis concoloribus. Fl. D. 879. f. 1. rup. mar.
1503. *L. ventosus*, leprofus flavus, tuberculis rubris. Fl. Sv. 1065. Fl. N. 939. Fl. D. 472. f. 1. 2. *L. Scopulorum* Fl. D. 712. f. 2. *L. cruentus* Web. Spic. t. 2. *L. flavesceus* Jacqu. Misc. 2. 79. t. 9. f. 1. *Verruc. ventosa* Hoffm. Lich. 2. p. 11. t. 27. f. 1. lap. mont.
1504. *L. Haematomma*, crustaceus albicans, tuberculis marginatis sanguineis. *L. Haematomma* Ehrhart Beytr. 2. p. 157. *Verruca Haematomma* Hoffm. Lich. I. p. 53. t. 11. f. 1. lap.

1505. *L.*

k) An idem cum terminali?

1505. *L. fagineus*, leprosus albus, tuberculis albis farinaceis. Fl. Sv. 1066. Fl. N. 941. l). cort. arb.
1506. *L. carpineus*, leprosus cinereus, tuberculis albidis rugosis. Fl. Sv. 1067. cort. arb.
1507. *L. corallinus*, leprosus ramosus teres fasciculatus fastigiatus confertissimus albus. M. Pl. 131. *L. defraudans* Fl. Isl. p. 17. ? rup.
1508. *L. pseudocorallinus*, leproso-crustaceus albidus, tuberculis subcylindricis inaequalibus concoloribus. m). rup.
1509. *L. Baeromyces*, leprosus albicans, tuberculis stipitatis carneis. Suppl. Pl. 450. *L. Ericetorum stipitatus* Fl. D. 472. f. 4. 1003. f. 2. Dill. Musc. t. 14. f. 1. terr.
1510. *L. Icmadophila*, leprosus cinereus, tuberculis sessilibus planis carneis. Suppl. Pl. 450. *L. Ericetorum sessilis* S. N. *L. parasiticus* Prodr. Fl. Scand. 1333. γ. musc.
1511. *L. candidus*, leprosus albus, sublobatus, tuberculis glauco-coerulescentibus. Web. Spic. 193. Fl. D. 1064. f. 1. rup.
1512. *L. variegatus*, leprosus albo rubroque colore variegatus. Nat. H. S. Skr. 2. 1. p. 17. Hag. Lich. p. 41. t. 1. f. 1. lap.
1513. *L. ocellatus*, leprosus cinereus, tuberculis concoloribus centro nigris. *Verrucaria ocellata* Hoffm. Lich. I. p. 92. t. 20. f. 2. lap.
β. *L. punctatus*. Prodr. Fl. Scand. 1335. Fl. D. 468! f. 2. lap.
1514. *L. marmoreus*, leprosus roseus, tuberculis concoloribus. Jacqu. Coll. 2. p. 178. t. 13. f. 1. *Verruc. purpurascens* Hoffm. Lich. I. p. 74. t. 15. f. 2. lap.

1515.

h) A *carpineo* haud diversum statuit Cel. *Afzelius* l. cit.
m) Cel. *Westring*.

1515. *L. rosaceus*, leprosus cinerascens, tuberculis niveis nigro punctatis. Fl. D. 825. f. 1. cort. arb.
- † 1516. *L. trapeziformis*, leprosus virescens, angulato-rimofus, tuberculis nigris. Fl. Isl. p. 15. Dill. Musc. t. 30. f. 133. terr.
1517. *L. rufescens*, crustaceus subcinereus, tuberculis rufis. Fl. D. 825. f. 2. *Patell. rufescens* Hoffm. Lich. l. p. 80. t. 17. f. 1. Sax.
1518. *L. fulgens*, crustaceus flavescens sublobatus, tuberculis carneis. N. Aët. Upf. 4. p. 246. terr.
1519. *L. rufus*, leprosus virescens, tuberculis substipitatis planiusculis fuscis. N. Aët. Upf. 4. p. 244. *L. Ericetorum fuscus* Prodr. Fl. Scand. 1333. β. Dill. Musc. t. 14. f. 2. terr.
1520. *L. sphaerocephalus*, crustaceus virescens, tuberculo stipitato nigro. N. Aët. Upf. 4. p. 245. *Mucor sphaerocephalus* Fl. Sv. 1286.
β. *Quercinus*. *Mucor Lichenoides*. Fl. Sv. 1287. Dill. Musc. t. 14. f. 3. cort. arb.
1521. *L. globularis*, crustaceus furfuraceus viridis, tuberculis globosis stipitatis fuscis. *L. furfuraceus* N. Aët. Upf. 4. p. 245. *Mucor furfuraceus* Fl. Sv. 1290. terr.
1522. *L. fulvus*, crustaceus furfuraceus pallidus, tuberculis globosis stipitatis fulvis. N. Aët. Upf. 4. p. 246. *L. sulphureus* Aët. Holm. 1769. p. 249. Prodr. Fl. Scand. 1334. *Mucor fulvus* Fl. Sv. 1289. lab. arb.

** *Leprosi scutellati*.

1523. *L. Jolithus*, leprosus ruber, peltis minutissimis planis turbinatis albidis. N. Aët. Upf. 4. p. 244. *Byssus Jolithus* Fl. Sv. 1178. Fl. N. 551. Fl. D. 899. f. 1. lap.
1524. *L. vitellinus*, leprosus vitellinus, scutellis concoloribus. *Patellaria vitellina* Hoffm. Lich. 2. p. 5. t. 26. f. 1. t. 27. f. 2. sep. lign.

1525. *L.*

1525. *L. candelarius*, crustaceus flavus, scutellis luteis. Fl. Sv. 1069. Dill. Musc. t. 1. f. 18. mr. par.
1526. *L. flavescent*, crustaceus flavo-virescens, scutellis rufo-luteis. N. Act. Upf. 4. p. 246. mr. lap. arb.
1527. *L. tartareus*, crustaceus ex albido-virescens, scutellis flavescentibus margine albo. Fl. Sv. 1070. Fl. N. 209. Dill. Musc. t. 18. f. 12. n). mont.
1528. *L. gelidus*, crustaceus albicans, peltis tuberculis rugosis testaceis. M. Pl. 133. Fl. N. 276. lap.
β. L. Heclae. Fl. N. 1021. Fl. D. 470. f. 2. lap.
1529. *L. rimosus*, crustaceus albicans, frustulis subrotundo-angulatis, scutellis glaucis albo-marginatis. Fl. D. 478. f. 3. lap.
1530. *L. frigidus*, crustaceus suffruticulosus albicans, ramulis brevissimis confertis, peltis marginatis testaceis. N. Act. Upf. 4. p. 247. Am. Acad. X. p. 117. t. 5. f. 4. alp.
1531. *L. pallescens*, crustaceus albicans, scutellis pallidis. Fl. Sv. 1071. Fl. N. 556. Dill. Musc. t. 18. f. 17. arb.
1532. *L. lentigerus*, crustaceus albicans sublobatus, scutellis albo-marginatis aetate flavescentibus convexisque. N. Act. Upf. 4. p. 246. Web. Spic. p. 192. t. 3. lap. terr.
1533. *L. subfuscus*, crustaceus albicans, scutellis nigris: junioribus urceolatis cavis. Fl. Sv. 1072. Fl. N. 579. Dill. Musc. t. 18. f. 16. arb.
1534. *L. muscorum*, crustaceus incanus, tuberculis atris connexis. N. Act. Upf. 4. p. 245. Fl. D. 1003. *Patell. muscorum* Hoffm. Lich. I. p. 93. t. 21. f. 1. terr. musc.

S 2

1535.

n) A frigido haud diversum contendit *Cel. Ehrhart* Beytr. 2. p. 46.

1535. *L. Saxorum*, crustaceus cinerascens, scutellis pallide rufis albo-marginatis. Fl. D. 712. f. 1. sax.
1536. *L. Parellus*, crustaceus albus, peltis concavis obtusis pallidis. Mant. Pl. p. 132. Dill. Musc. t. 18. f. 10. *Patell. Perella* Hoffm. Lich. I. p. 60. t. 12. f. 5. mr. lap.
1537. *L. subcarneus*, crustaceus albidus, scutellis confertissimis subcarneis difformibus o). lap.
1538. *L. Upsalienfis*, crustaceus, foliolis subulatis striatis. Fl. Sv. 1073. Fl. N. 562. *Patell. Upsalienfi* Hoffm. Lich. I. p. 95. t. 21. f. 2. p).
1539. *L. Pezizoides*, farinaceus glaucus, scutellis planis confertis fulvis marginatis. N. A. Upf. 4. p. 247. Web. Spic. p. 200. terr. ul.
1540. *L. brunneus*, crustaceus fuscus, scutellis planis rufis. N. Aët. Upf. 4. p. 247. terr.
1541. *L. Ulmi*, crustaceus albicans, scutellis minutis rufis crenatis albo-marginatis. N. A. Upf. 4. p. 247. cort. Ulmi.
1542. *L. ater*, crustaceus albicans rugosus, scutellis atris margine albo subcrenato. Hudf. Fl. Angl. II. p. 530. Dill. Musc. t. 18. f. 15. cort. arb. my.
1543. *L. amylaceus*, crustaceus candidus superficie subcinerascens rimosus, scutellis atris margine albis integris q).
1544. *L. varius*, crustaceus, crusta fugaci, scutellis congestis marginatis cinerascens subfuscisque. *Patellaria varia* Hoffm. Lich. I. p. 102. t. 23. f. 4. lign.

1545. *L.*o) *Cel. Westring.*

p) Foliola adesse negat *Cel. Ehrhart* Beytr. I. p. 137. nec illa in meo specimine video, sed crustam muscorum reliquias obducentem peltasque albas intra marginem farina quasi asperfas.

q) *Cel. Westring.* Crusta candida quasi ex farina compacta in superficie cinerascens rimosa. Forte varietas aliqua sub *Lich. atro* Hudf. a *Dillenio* relata huc pertinet.

1545. *L. impressus*, leprofus albicans, scutellis marginatis concavis atris. N. Aët. Upf. 4. p. 245. lap.
1546. *L. scruposus*, crustaceus cinereus granosus, scutellis immerfis atris: margine crenulato. Schreb. Lipf. 1133. Dickf. Crypt. I. p. 11. *Patell. scruposa* Hoffm. Lich. 1. p. 54. t. 11. f. 2. Dill. Musc. t. 18. f. 15. B. mr.
- † 1547. *L. Vahlîi*, crustaceus pulverulentus lacteus, scutellis coccineis marginatis: margine integro. *L. lacteus* N. Hist. S. Skr. II. 1. p. 69. mont.
1548. *L. aurantiacus*, leprofus albido-cinereus, scutellis aurantiacis. Lightf. Fl. Scot. II. p. 810. r). cort. lign. putr. lap.

*** *Imbricati.*

1549. *L. microphyllus*, imbricatus crustaeformis cinereus, foliolis minutis granulosis, peltis junioribus congestis rufis: adultis discretis nigricantibus s). saxa.
1550. *L. Hypnorum*, imbricatus crustaeformis: foliolis minutis lobatis virescentibus, scutellis ferrugineis margine crenulatis subtus granulatis. N. Hist. S. Skr. II. 1. p. 55. Fl. D. 956. musc. alp.
1551. *L. crassus*, imbricatus, foliolis lobatis crenatis rotundatis virescentibus, scutellis planis fulvis. N. Aët. Upf. 4. p. 247. Fl. D. 1006. Dill. Musc. t. 24. f. 74. lap. terr.

S 3

1552. *L.*

- r) Ab hoc diversissimus *L. aurant.* Latourette vel *Platisma armatum* Hoffm.
- s) *Cel. Westring.* Peltae juniores forma tuberculorum congestorum apparent.

1552. *L. saxicola*, imbricatus, foliolis dense congestis pallide viridulis margine albicantibus sublobatis, scutellis flavis congestis margine subcrenulato. *L. saxicola* Poll. 3. p. 225. t). lap.
1553. *L. luridus*, imbricatus crustaeformis, foliolis minutis sinuatis crassis fusco viridibus subtus albidis, tuberculis nigris. Dickf. Crypt. Fasc. 2. p. 20. N. Hist. S. Skr. 2. 1. p. 60. N. Act. Ups. 4. p. 247. Fl. D. 1064. f. 2. Dill. Musc. t. 30. f. 34. rup.
1554. *L. fillicidiorum*, imbricatus crustaeformis viridicinerascens, scutellis flavis cinereo-marginatis. N. Hist. S. Skr. 2. 1. p. 28. Fl. D. 1063. f. 2. musc. rup. hum.
1555. *L. Jungermanniae*, imbricatus crustaeformis cinereus, scutellis convexis fulvis margine planis. N. Hist. S. Skr. 2. 1. p. 29. Fl. D. 1063. f. 1. jung. cil.
1556. *L. leucophaeus*, imbricatus crustaeformis, foliolis crenato-multifidis glauco-fuscis, scutellis convexis fuscis albedo marginatis. N. Hist. S. Skr. 2. 1. p. 14. Fl. D. 955. f. 2. Dill. Musc. t. 82. f. 2. rup.
1557. *L. pulverulentus*, imbricatus, foliolis lobatis crenatis obtusis viridibus polline conspersis, scutellis nigris margine crenatis. Schreb. Lips. 1123. Web. Spic. 276. Dill. Musc. t. 24. f. 71. *Squamaria pulverulenta* Hoffm. Lich. I. p. 39. t. 8. f. 2. cort. lap.

1558. *L.*

- *) *Cel. Westring.* Dubius sum an hic vel *L. crassus* genuinus sit *saxicola* Pollichii.

155. *L. centrifugus*, imbricatus, foliolis obsolete multifidis laevibus albidis centrifugis, scutellis rufofuscis. Fl. Sv. 1074. Fl. N. 568. Fl. L. t. 11. f. 2. *Squamaria centrifuga* Hoffm. Lich. I. p. 78. t. 16. f. 2. Dill. Musc. t. 24. f. 75. u). lap.
1559. *L. cartilagineus*, imbricatus, foliolis crenatis pallidis cartilagineis, scutellis planis fulvis. Lightf. Fl. Scot. 2. p. 815. Dill. Musc. t. 24. f. 74. rup.
1560. *L. saxatilis*, imbricatus, foliolis finuatis scabris lacunosis, scutellis badiis. Fl. Sv. 1075. Fl. N. 210. Dill. Musc. t. 24. f. 83. lap.
1561. *L. omphalodes*, imbricatus, foliolis multifidis glabris obtusis incanis, punctis vagis eminentibus. Fl. Sv. 1076. Dill. Musc. t. 24. f. 80. lap. pr.
1562. *L. olivaceus*, imbricatus, foliolis lobatis nitidis lividis, scutellis crenatis. Fl. Sv. 1077. Dill. Musc. t. 24. f. 78. lap. arb.
1563. *L. pullus*, imbricatus, foliolis lobatis nitidis, scutellis integerrimis. Schreb. Lipf. p. 131. N. Aët. Upf. 4. p. 249. arb. lap.
1564. *L. diffusus*, imbricatus glauco albescens, fronde lineari-multifida in centro pulverulenta versus margines scutellifera. Web. Spic. 278. w). arb.
1565. *L. Fahlunensis*, imbricatus, foliolis linearibus dichotomis planiusculis acutis nigris, scutellis atris. Fl. Sv. 1078. Fl. D. 958. Jacqu. Misc. 2. p. 90. t. 10. f. 2. *Squamaria Fahlunensis* Hoffm. Lich. 2. p. 45. t. 36. f. 2. Dill. Musc. t. 24. f. 81. rup.
1566. *L. stygius*, imbricatus, foliolis palmatis recurvis atris. Fl. Sv. 1079. Fl. N. 550. *Squamaria stygia* Hoffm. Lich. 2. p. 3. t. 25. f. 2. rup.

S 4

1567. *L.*

u) Diversum ab hoc faciunt *L. conspersum*, ego autem ex exemplari a Cel. Westring. communicato differentiam determinare haud potui.

w) Cel. Westring. at supra lapidea.

1567. *L. crispus*, imbricatus, foliis lobatis truncatis crenatis atro-viridibus, scutellis concoloribus. S. N. 959. Dill. Musc. t. 19, f. 23. lap. mr.
1568. *L. sinuatus*, imbricatus, foliolis subrotundis sinuatis crenatis fusco-viridibus, scutellis concoloribus. Hudf. Fl. Angl. 2. p. 535. Dill. Musc. t. 19, f. 33. rup.
1569. *L. muralis*, imbricatus, flavo virescens, scutellis concoloribus demum lutescentibus margine pallido. Schreb. Lipf. p. 130. *L. Ochroleucus* Jacqu. Coll. 2. 192. t. 13. f. 4. *Pjõra muralis* Hoffm. Lich. I. p. 77. t. 16. f. 1. x).
1570. *L. parietinus*, imbricatus, foliolis crispis fulvis, peltis concoloribus fulvis. Fl. Sv. 1080. Fl. N. 207. Fl. D. 1005. Dill. Musc. t. 24, f. 76. lap. lign.
1571. *L. physodes*, imbricatus, laciniis obtusis subinflatis, Fl. Sv. 1081. Fl. N. 563. Dill. Musc. t. 20. f. 49. arb.
1572. *L. stellaris*, imbricatus, foliolis oblongis laciniatis angustis cinereis, scutellis pullis. Fl. Sv. 1082. Fl. D. 957. Dill. Musc. t. 24. f. 70. arb.

**** *Foliacei.*

1573. *L. Burgeffi*, subfoliaceus crispus, peltis elevatis extus muricato-crispis: fundo depresso plano. S. N. 959. Lightf. Scot. 2. p. 827. t. 26. arb. frut.
1574. *L. ciliaris*, foliaceus, erectiusculus, laciniis linearibus ciliatis, scutellis pedunculatis crenatis. Fl. Sv. 1083. Fl. N. 575. *Lichenoides ciliare* Hoffm. Lich. I. p. 16. t. 3. f. 4. Dill. Musc. t. 20. f. 45. Vaill. Par. t. 20. f. 4. arb.

1575. *L.**) A *L. pallescens* Linn, certe diversus.

1575. *L. hispidus*, foliaceus albus, laciniis subturbina-
tis ciliato-spinulosis, scutellis subseffilibus. *L. hi-*
spidus Schreb. Lips. 1120. *L. tenellus* N. Act. Upf. 4.
p. 248. *Lichen. hispidum* Hoffm. Lich. I. p. 13. t. 3.
f. 3. Dill. Musc. t. 20. f. 46. Vaill. Par. t. 20. f. 5.
arb. lap.
1576. *L. Islandicus*, foliaceus adscendens laciniatus
marginibus elevatis ciliatis. Fl. Sv. 1085. Fl. N.
425. terr.
 α . *rigidus*. Dill. Musc. t. 28. f. 111. Fl. D. 155.
 Lichenoid. islandicum Hoffm. Lich. I. p. 40.
 t. 9. f. 1.
 β . *tenuifolius*. Dill. Musc. t. 28. f. 112.
 γ . *nigricans*. ibid. t. 17. f. 32. y).
- † 1577. *L. Groenlandicus*, foliaceus repens, laciniis ob-
liquis rotundis concavis viridulo-glaucis: inferne
nigris, scutellis marginalibus subpedunculatis,
Fl. D. 466. Fl. N. 1019. terr.
1578. *L. nivalis*, foliaceus adscendens laciniatus cri-
spus glaber lacunofus albus, margine elevato.
Fl. Sv. 1086. Fl. N. 316. Fl. D. 227. Fl. L. t. 111.
f. 1. Dill. Musc. t. 21. f. 56. fab.
1579. *L. pulmonarius*, foliaceus laciniatus obtusus gla-
ber, supra lacunofus, subtus tomentosus, scutellis
marginalibus. Fl. Sv. 1087. Fl. N. 538. Fl. D.
1007. Dill. Musc. t. 29. f. 113. *Pulmonaria reticu-*
lata Hoffm. Lich. I. p. 4. t. 1. f. 2. arb.
1580. *L. furfuraceus*, foliaceus decumbens furfuraceus,
laciniis acutis subtus lacunofis atris. Fl. Sv. 1088.
Fl. N. 775. Dill. Musc. t. 21. f. 52. *Lichenoid. fur-*
furaceum Hoffm. Lich. I. p. 45. t. 9. f. 2. arb.
- S 5
1581. *L.*

y) Hunc mihi e Germania communicavit amicus sub nomine
L. aculeatus Schreb., at ab illo Hoffm. Lich. I. p. 26. t. 5. f. 2.
diversissimus est.

1581. *L. farinaceus*, foliaceus erectus compressus ramifus, verrucis marginalibus farinosis. Fl. Sv. 1089. Fl. N. 691. Dill. Musc. t. 23. f. 63. arb.
1582. *L. calicaris*, foliaceus erectus lineari-ramosus lacunofus convexus mucronatus. Fl. Sv. 1090. Dill. Musc. t. 23. f. 62. arb.
1583. *L. scopulorum*, foliaceus rectus compressus linearis laciniatus nitidus, lateribus verrucosis. Obs. Bot. Fasc. IV. p. 30. *L. calicaris* Fl. D. 959. Dill. Musc. t. 22. f. 60. rup. mar.
1584. *L. fraxineus*, foliaceus erectus oblongus lanceolatus sublaciniatus lacunofus glaber, scutellis subpendunculatis. Fl. Sv. 1091. Fl. N. 577. Dill. Musc. t. 22. f. 59. *Platisma fraxineum* Hoffm. Lich. I. p. 85. t. 18. arb.
1585. *L. Prunastri*, foliaceus erectiusculus lacunofus, subtus tomentosus albus. Fl. Sv. 1092. Fl. N. 553. arb.
- α. incanus*. Dill. Musc. t. 21. f. 55.
- β. albidus*. ibid. f. 54.
1586. *L. Juniperinus*, foliaceus laciniatus crispus fulvus, peltis lividis. Fl. Sv. 1093. Fl. N. 204. Fl. D. 1004. Dill. Musc. t. 24. f. 76. *Squamar. Juniperina* Hoffm. Lich. I. p. 35. t. 7. f. 2. arb.
1587. *L. caperatus*, foliaceus repens lobatus, supra luteo-virescens, infra laevis ater, scutellis verrucosis sessilibus umbilicatis rubris. Scop. Fl. Carn. 1394. S. N. 960. Fl. N. 690. Jacqu. Coll. 4. p. 280. t. 20. f. 1. Dill. Musc. t. 25. f. 97. *Platisma caperatum* Hoffm. Lich. II. p. 50. t. 37. f. 1. t. 38. f. 1. t. 42. f. 1. arb. lap.
1588. *L. glaucus*, foliaceus depressus lobatus glaber margine crispo farinaceo. Fl. Sv. 1094. Fl. N. 761. Fl. D. 598. Dill. Musc. t. 25. f. 96. arb.

1589. *L. nigrescens*, foliaceus gelatinosus subrotundus lobatus rugosus atrovirens, scutellis rufis. S. N. 961. N. Aët. Upf. 4. p. 249. *L. Lactuca* Web. Spic. p. 253. *L. Vespertilio* Lightf. Fl. Scot. 2. p. 840. *L. lacer* Prodr. Fl. Scand. 1369. Fl. N. 1022. Fl. D. 470. f. 3. Dill. Musc. t. 19. f. 20. arb. lap.

1590. *L. granulatus*, foliaceus gelatinosus, laciniis subrotundis crenatis scabriusculis atroviridibus, scutellis fulvis. N. A. Upf. 4. p. 249. Hudf. Fl. Angl. 2. p. 536. Dill. Musc. t. 19. f. 24. lap. mr. ar.

β. *Tremella Lichenoides* Fl. Sv. 1138. Dill. Musc. t. 19. f. 31. *L. tremelloides* Hudf. Fl. Angl. 2. p. 537. S. N. 961. *Collema plicatum* Hoffm. Lich. II. p. 41. t. 35. f. 2.

1591. *L. rupestris*, foliaceus membranaceo-gelatinosus lobatus, laciniis obtusis, scutellis minutis sparsis fufis. N. Aët. Upf. 4. p. 249. S. N. 960. Dill. Musc. t. 19. f. 22. rup. hum.

1592. *L. tenax*, foliaceus gelatinosus plumbeus, laciniis lobatis sinuatis, scutellis sparsis concavis rufis. N. Aët. Upf. 4. p. 249. terr. musc.

1593. *L. palmatus*, foliaceus gelatinosus, laciniis imbricatis subpalmatis crenatis glaucis, scutellis rufis. N. Aët. Upf. 4. p. 249. Dill. Musc. t. 19. f. 30. Vaill. Par. t. 21. f. 15. eric. mont.

1594. *L. marginalis*, foliaceus gelatinosus, laciniis imbricatis multifidis dentato-crenatis atroviridibus, scutellis marginalibus planiusculis atro-fuscis. N. Aët. Upf. 4. p. 250. Dill. Musc. t. 19. f. 25. 2). rup.

1595. *L. fascicularis*, foliaceus gelatinosus, tuberculis turbinatis fasciculatis fronde majoribus. S. N. 961. Fl. D. 462. f. 2. Dill. Musc. t. 19. f. 27. terr.

† 1596.

2) *L. palmatus* et *marginalis* forte melius *L. imbricatis* adderentur, at non vidi ipse.

- † 1596. *L. granosus*, gelatinosus sinuato - plicatus, lobis erectiusculis rotundatis, tuberculis sphaeroidibus gelatinosis aggregatis. Jacqu. Coll. 3. p. 131. t. 10. f. 2. Fl. D. 462. f. 1. Dill. Musc. t. 19. f. 24. ul. musc.
- † 1597. *L. gyrosus*, gelatinosus, foliorum verticalium marginibus subaequalibus, superficie densi caespitis eleganter gyrosa, peltis marginalibus spadicis. Fl. N. 974. Fl. D. 463. f. 2. terr.

***** *Coriacei.*

1598. *L. herbaceus*, coriaceus repens obtuse lobatus crenatus glaber herbaceus, scutellis sparsis rufis. N. Aët. Upf. 4. p. 250. Dill. Musc. t. 25. f. 98. *Pulmonar. herbacea* Hoffm. Lich. I. p. 51. t. 10. f. 2, arb. alp.
1599. *L. scrobiculatus*, coriaceus repens lobatus obtusus superne scrobiculatus verrucosus, inferne villosus albidus. N. Aët. Upf. 4. p. 250. *L. verrucosus* Hudf. Fl. Angl. 2. p. 545. Dill. Musc. t. 29. f. 114. Jacqu. Coll. 4. p. 278. t. 18. f. 2. *Pulmonar. verrucosa* Hoffm. Lich. I. p. 1. t. 1. f. 1. arb. rad.
1600. *L. aquaticus*, coriaceus repens lobatus obtusus, peltis hemisphaericis maximis. Fl. Sv. 1095. Fl. N. 762. Dill. Musc. t. 20. f. 44. pal.
1601. *L. fluviatilis*, coriaceus repens lobatus obtusus, lobis margine integerrimis, superne viridis, inferne ochroleucus lacunofus. N. Aët. Upf. 4. p. 250. Web. Spic. p. 265. t. 4. Dill. Musc. t. 30. f. 28. lap. rivul.
1602. *L. resupinatus*, coriaceus repens lobatus, peltis marginalibus posticis. Fl. Sv. 1096. Fl. N. 763. Fl. D. 764. Dill. Musc. t. 28. f. 105. Jacqu. Coll. 4. p. 257. t. 12. f. 1. terr.

1603. *L. venosus*, coriaceus repens ovatus planus, sub-
tus venosus villosus, peltis marginalibus horizon-
talibus. Fl. Sv. 1097. Fl. N. 764. Dill. Musc.
t. 28. f. 109. *Peltigera venosa* Hoffm. Lich. I. p. 31.
t. 6. f. 2. terr.
1604. *L. aphthosus*, coriaceus repens lobatus obtusus,
verrucis sparsis, pelta marginali adscendente. Fl.
Sv. 1098. Fl. N. 208. Fl. D. 767. f. 1. Dill. Musc.
t. 28. f. 106. Jacqu. Coll. 4. p. 266. t. 17. *Pelti-
gera aphthosa* Hoffm. Lich. I. p. 28. t. 6. f. 1.
terr. lap. sylv.
1605. *L. arcticus*, coriaceus repens lobatus obtusus pla-
nus laevis, subtus avenius villosus. Fl. Sv. 1099.
Fl. N. 765. *L. antarcticus* S. N. 960. Jacqu. Misc. 2.
p. 370. t. 10. f. 1. a). terr.
1606. *L. caninus*, coriaceus repens lobatus obtusus pla-
nus subtus venosus villosus, pelta marginali adscen-
dente. Fl. Sv. 1100. Fl. N. 557. Fl. D. 767. f. 2.
Dill. Musc. t. 27. f. 102. Jacqu. Coll. 4. p. 260.
t. 14. f. 1. sylv. lap.
1607. *L. sylvaticus*, coriaceus repens laciniatus lacuno-
sus, peltis marginalibus adscendentibus. S. N. 961.
Dill. Musc. t. 27. f. 101. Jacqu. Coll. 4. p. 258.
t. 12. f. 2. *Peltig. sylvatica* Hoffm. Lich. I. p. 21.
t. 4. f. 2. lap. arb.
1608. *L. polydactylus*, coriaceus repens lobatus, lobis
digitatis, peltis marginalibus adscendentibus. Jacqu.
Coll. 4. p. 262. t. 14. f. 2. Dill. Musc. t. 28. f. 107.
Peltig. polydactylon Hoffm. Lich. I. p. 19. t. 4. f. 1.
sylv. lap.

1609. *L.*

a) *L. arcticus* noster et *antarcticus* Jacqu. absolute idem, etiam
peltarum magnitudine.

1609. *L. horizontalis*, coriaceus repens planus subtus
 avenius, peltis marginalibus horizontalibus. S. N.
 961. Fl. N. 1058. Fl. D. 533 et 765. Dill. Musc.
 t. 28. f. 104. Jacqu. Coll. 4. p. 265. t. 16. sylv.
 1610. *L. faccatus* coriaceus repens subrotundus, peltis
 depressis subtus faccatis. S. N. 962. Fl. N. 1057.
 Fl. D. 532. f. 3. Dill. Musc. t. 30. f. 121. terr.
 1611. *L. croceus*, coriaceus repens subrotundus planus,
 subtus venosus villosus croceus, peltis sparsis ad-
 natis. Fl. Sv. 1101. Fl. N. 735. Fl. D. 263. Fl.
 L. t. 11. f. 3. Dill. Musc. t. 30. f. 120. Jacqu.
 Coll. 4. p. 255. t. 11. f. 23. *Peltig. crocea* Hoffm.
 Lich. 2. p. 60. t. 41. f. 2-4. t. 42. f. 4. 5. alp.

***** *Umbilicati.*

1612. *L. griseus*, umbilicatus subrotundus, sublobatus,
 laevis, supra griseo-glaucus subtus nigrescens
 punctis elevatis papillofus. *L. griseus* N. Act. Svec.
 1793. p. 52. b). lap.
 1613. *L. miniatus*, umbilicatus gibbus punctatus, subtus
 fulvus. S. N. 962. Fl. N. 1056. Fl. D. 532. f. 1. Jacqu.
 Misc. 2. p. 90. t. 10. f. 3. Dill. Musc. t. 30. f. 127. lap.
 1614. *L. velleus*, umbilicatus cinereus subtus hirsutissi-
 mus, peltis immerfis planis. Fl. Sv. 1103. Fl. N.
 555. Web. Spic. 287. c). lap.
 β. *L. glaucus*, umbilicatus margine lobatus, sub-
 inde polyphyllus, rigidus, supra glaucus sub-
 tus hirsutus niger. N. Act. Holm. 1793.
 p. 59. d). lap.

1615. *L.*

b) *Cel. Westring.*

c) A Dill. t. 82. f. 5. diversus videtur, nec cum Halleriano
 1997 convenit. Peltae in adulto tantum adfunt. *Umbil.*
vellea Hoffm. Lich. 2. p. 9. t. 26. f. 3. a nostro differt, quod
 nimis lobatus sit peltasque convexas habeat. *Umbil. cirrofa*
 ibid. 1. p. 9. t. 2. f. 3. 4. propius nostro accedit et certe ejus
 est varietas.

d) *Cel. Westring.* diversum credidit, mihi varietas *vellei* videtur.

1615. *L. pustulatus*, umbilicatus subtus lacunofus, furfure nigro adsperfus. Fl. Sv. 1107. Fl. N. 552. Fl. D. 597. f. 2. Dill. Musc. t. 30. f. 131. *Umbilicaria pustulata* Hoffm. Lich. II. p. 13. t. 28. f. 1. 2. t. 29. f. 3. lap.

† 1616. *L. exasperatus*, umbilicatus laevis, tuberculis contortis nigris. *L. exasperatus* Pr. Fl. Scand. 1383. Fl. N. 1024. Fl. D. 471. f. 3. *Umbilic. exasperata* Hoffm. Lich. I. p. 7. t. 2. f. 1. 2. rup.

1617. *L. erosus*, umbilicatus diaphanus, subtus albescens fibrosus, peltis turbinatis atris. N. Aët. Upf. 4. p. 250. Web. Spic. p. 259. Dill. Musc. t. 30. f. 118. e).

1618. *L. Cribellum*, umbilicatus laevis niger sublobatus, margine lacinulato denticulatus, poris irregularibus pertusis reticulatus. *L. reticularis* N. A. Holm. 1793. p. 45. f). lap.

1619. *L.*

e) Haud vidi, nisi sequentem intelligat *Cel. Swartz*, quem ab *eroso* *Web.* plane diversum judico, si descriptioni aliqua fides habenda; neque enim diaphanus nec subtus albescens vel cirrosus. *Dillenii* descriptio ad figuram 118 quoque a *Cribello* longe alienam indicat speciem.

f) Pulcherrimus Lichen difficile describitur. Frons in centro radice unica, suborbicularis, supra nigra, subtus pallidior, in medio integra, dehinc quasi ex foliis varie lacinulatis componitur. Laciniae hae margine denticulatae, et hi denticuli conjungunt lacinias in unam frondem varie perforatam cribri instar; at in margine lacinulae liberae. Pagina superior e centro nonnullas rugas elevatas habet et praeterea peltas multas atras planiusculas rugosas.

1619. *L. proboscideus*, umbilicatus laciniatus laevis, laciniis margine ramoso setaceo-ciliatis, peltis turbinatis subpedunculatis. *L. proboscideus* L. Spec. Pl. 1617. Fl. Sv. 1106. Jacqu. Misc. T. 2. p. 80. t. 9. f. 2. g). *L. crinitus* Lightf. Fl. Scot. p. 660. Dill. Musc. t. 29. f. 116. A. B. Fl. D. 471. f. 1. 2. *L. corneus* Pr. Fl. Scand. 1381. Fl. N. 1023. alp.
1620. *L. deustus*, umbilicatus undique laevis. Fl. Sv. 1105. Fl. N. 736. Dill. Musc. t. 30. f. 117.
1621. *L. polyphyllus*, umbilicatus polyphyllus undique laevis atrovirens crenatus. Fl. Sv. 1104. Fl. N. 767. Dill. Musc. t. 30. f. 129. lap.
 β . *L. polyphyllus*. N. Aët. Holm. 1793. p. 46.
 γ . *L. glaber*. ibid. p. 48.
1622. *L. complicatus*, umbilicatus polyphyllus imbricatus lobatus punctatus fusco-viridis. N. Aët. Upf. 4. p. 251. *L. miniatus* β . Fl. D. 532. f. 2. Fl. Scot. 2. p. 857. *L. polyphyllus* Jacqu. Coll. 4. p. 190. t. 16. f. 1. Dill. Musc. t. 30. f. 128. rup. mar.
1623. *L. polyrhizos*, umbilicatus polyphyllus supra laevis, polyrhizos. Fl. Sv. 1108. Fl. N. 692. Fl. D. 97. f. 1. ? Dill. Musc. t. 30. f. 130. *L. hirsutus* N. Aët. Holm. 1793. p. 47. lap.

† 1624.

g) *Lichenoides* Dill. t. 20. f. 42. sine dubio est *L. perforatus* Jacqu. Coll. I. p. 116. t. 3. (confirmante etiam Nob. Smith in litt.) quem in prima Spec. Pl. edit. *L. cylindricum* vocavit ill. à Linné, dein vero perperam cum *proboscideo* alienissimo conjunxit. Ratio hujus confusionis mihi patuit ex exemplari, in quo laciniarum sinus ita flexi atque simul elevati sunt, ut peltas conicas centro perforatas, oculo minus attente scrutanti, haud male referant. Coeterum variat noster colore cinerascens et atro, folio unico varie laciniato et pluribus, ciliis marginalibus solitariis vel etiam simul hinc inde e pagina erumpentibus, brevioribus vel demum longioribus. In hoc praeterea cilia vel setae variae ramosae, in *L. polyrhizo* breviores sunt, simplices, at apice quasi penicillatae, ut antea observavit *Lightfoot*.

† 1624. *L. reticulatus*, umbilicatus multifidus, laciniis margine conglutinatis. Fl. Isl. p. 16.

***** *Scyphiferi h).*

1625. *L. cocciferus*, scyphifer simplex integerrimus, stipite cylindrico, tuberculis coccineis. Fl. Sv. 1109. Fl. N. 359. terr. lap.

α. simplex, tuberculis sessilibus stipitatisve.

Dill. Musc. t. 14. f. 7. A. F. H. I.

β. simplex, tuberculis totum discum occupantibus. Vaill. Par. t. 21. f. 10.

γ. e margine prolifer tuberculatus. Dill. Musc. t. 14. f. 7. K. L.

δ. e centro prolifer. ibid. f. M.

1626. *L. cornucopioides*, scyphifer simplex folio brevior, tuberculis coccineis. Fl. Sv. 1110. Fl. N. 360. Dill. Musc. t. 14. f. 9. terr.

1627. *L. pyxidatus*, scyphifer simplex crenulatus, tuberculis fuscis. Fl. Sv. 1111. Fl. N. 241. terr.

α. simplex tuberculis nullis. Dill. Musc. t. 14. f. 5. A. B.

β. simplex tuberculis sessilibus stipitatisve. ibid. f. 6. C.

γ. e disco prolifer, tuberculis nullis. ibid. f. 6. D. E. F. G.

δ. e disco prolifer tuberculatus. ibid. f. 6. H.

ε. e margine prolifer, tuberculis nullis. ibid. f. 6. I.

ζ. e margine prolifer tuberculatus. ibid. f. 6. K. L. M.

1628. *L. fimbriatus*, scyphifer simplex denticulatus, stipite cylindrico. Fl. Sv. 1112. Fl. N. 968. terr.

α. simplex. Dill. Musc. t. 14. f. 8. A. B.

β. prolifer. ibid. f. 8. C.

1629.

h) Num Scyphiferorum haec distributio genuina? confirmatione eget.

1629. *L. gracilis*, scyphifer ramosus denticulatus filiformis. Fl. Sv. 1113. 1). terr.
 α. simpliciusculus, tuberculis nullis. Dill. Musc. t. 14. f. 13. C. D.
 β. simpliciusculus scyphoideus tuberculatus, ibid. f. 13. A. B.
 γ. ramosus, tuberculis minimis. ibid. f. 13. E.
1630. *L. digitatus*, scyphifer ramosissimus, ramis cylindricis, calycibus integris nodosis. Fl. Sv. 1114. Fl. N. 769. terr.
 α. tuberculatus. Dill. Musc. t. 15. f. 19. C.
 β. etuberculatus. ibid. t. 15. f. A. B.
1631. *L. cornutus*, scyphifer simpliciusculus subventricosus, calycibus integris. Fl. Sv. 1115. Fl. N. 770. Dill. Musc. t. 15. f. 14. 18. *Cladonia cornuta* Hoffm. Lich. II. p. 1. t. 25. f. 1. terr.
1632. *L. deformis*, scyphifer simpliciusculus subventricosus, calycibus dentatis. Fl. Sv. 1116. Fl. N. 771. Fl. L. t. 11. f. 5. terr.
1633. *L. radiatus*, scyphifer subramosus cylindraceus elongatus, calycibus inaequalibus dentatis radiatisque. Schreb. Lipf. 122. Dill. Musc. t. 15. f. 16. terr.
1634. *L. cristatus*, scyphifer, scypho ramuloso-proli-fero, ramulis fusco-tuberculatis. *L. pyxidatus cristatus* Weis Crypt. p. 94. Dill. Musc. t. 15. f. 17. terr.
1635. *L. foliaceus*, scyphifer, foliis adscendentibus multifido-laciniatis subtus niveis, scyphulis conicis brevissimis. Schreb. Lipf. 122. Dill. Musc. t. 14. f. 12. A. terr.

***** Fru.

1) In *digitatum* abire observare voluit ill. Pallas it. T. 2. p. 44.

***** *Fruticulosi*.

1636. *L. Papillaria*, fruticulosus perforatus albidus, ramis paucis obtusis brevissimis, tuberculis terminalibus carneis. Ehrh. Beytr. 2. p. 106. Dill. Musc. t. 16. f. 28. *L. madreporaeformis* Jacqu. Coll. 3. p. 105. t. 16. f. 2. terr.
1637. *L. rangiferinus*, fruticulosus perforatus ramosissimus, ramulis nutantibus. Fl. Sv. 1117. Fl. N. 269.
 α. *alpestris*, Fl. D. 180. Dill. Musc. t. 16. f. 29. alp.
 β. *sylvaticus*. Fl. D. 539. Dill. Musc. t. 16. f. 30. campi.
- † 1638. *L. pallidus*, fruticulosus ramosissimus pallide sulphureus, ramis implexis apice viridi-nigricantibus, farinoso tuberculatis. *L. Sulphureus* Fl. Isl. p. 16. an *aurantiaco ater* Jacqu. Misc. 2. p. 369. t. 11. f. 2. ? k).
1639. *L. uncialis*, fruticulosus perforatus, ramulis brevissimis acutis. Fl. Sv. 1118. Fl. N. 772. Dill. Musc. t. 16. f. 22. sylv.
1640. *L. subulatus*, fruticulosus tubulosus subdichotomus, ramis simplicibus subulatis, Fl. Sv. 1119. Fl. N. 560. Dill. Musc. t. 16. f. 26. sylv.
1641. *L. Vermicularis*, fruticulosus laevis subramosus, ramis diffusis tuberculis sparsis. S. N. 963. N. Act. Upf. 4. p. 248. *L. subuliformis* Ehrh. Beytr. 3. p. 82. *L. tauricus* Jacqu. Coll. 2. p. 177. t. 12. f. 2. *Cladonia subuliformis* Hoffm. Lieh. 2. p. 15. t. 29. f. 1-3. alp.

T 2

† 1642.

k) Specimen meum ex Islandia sine scutellis et minus divaricatis ramis, videtur atrum esse corticeque flavo pulverulento inductum.

- † 1642. *L. arenarius*, fruticulosus rugosus ramosissimus candicans, ramis diffusis. *l.* ar. vol.
1643. *L. globiferus*, fruticulosus solidus laevis, tuberculis globosis cavis terminalibus. S. N. 962. Fl. D. 960. *Coralloides globiferum* Hoffm. Lich. II. p. 25. t. 31. f. 2. Dill. Musc. t. 17. f. 35. lap. sylv.
1644. *L. paschalis*, fruticulosus solidus tectus foliolis crustaceis. Fl. Sv. 1120. Fl. N. 268. Fl. D. 151. Dill. Musc. t. 17. f. 33. *Coralloides paschale* Hoffm. Lich. I. p. 23. t. 5. f. 1. lap. lign.
1645. *L. fragilis*, fruticulosus solidus, ramulis teretibus nudis obtusis. Fl. Sv. 1121. Fl. N. 663. Dill. Musc. t. 17. f. 34. *Coralloid. fragile* Hoffm. Lich. II. p. 34. t. 33. f. 3. lap.
1646. *L. Botrytis*, fruticulosus subramosus, tuberculis terminalibus sphaericis flavo-brunneis maximis. N. Hist. S. Skr. 2. 1. p. 28. Hag. Lich. t. 9. f. 9. lign.
1647. *L. Muscicola*, fruticulosus ramosissimus, ramis brevissimis subulatis atroviridibus, scutellis concoloribus. N. Act. Ups. 4. p. 248. Dickf. Fasc. II. t. 6. f. 9. musc.
1648. *L. tristis*, fruticulosus solidus, ramulis compressis ramosis, apicibus nigricantibus, scutellis terminalibus. S. N. 963. Web. Spic. p. 209. t. 5. *L. rigidus* S. N. 959. Jacqu. Coll. 2. p. 187. t. 13. f. 5. ? *L. fucoides* Jacqu. Coll. 3. p. 143. t. 12. f. 3. *Cornicularia tristis* Hoffm. Lich. II. p. 36. t. 34. f. 1. *L. Normoericus* Pr. Fl. Scand. 1416. Fl. N. 973. t. 2. f. 9-14. Dill. Musc. t. 17. f. 37. mont. ar. vol.

***** *Fi.*

l. Medius inter fruticulosos et Usneas. Fructificationes nullas vidi. Legit *Cel. Weigel* in collibus arenosis Insulae Fasmund.

***** *Filamentosi.*

1649. *L. plicatus*, filamentosus pendulus, ramis implexis scutellis radiatis. Fl. Sv. 1122. Fl. N. 176. Dill. Musc. t. 11. f. 1. arb.
1650. *L. barbatus*, filamentosus pendulus subarticulatus, ramis patentibus. Fl. Sv. 1123. Fl. N. 175. Dill. Musc. t. 12. f. 6. m). arb.
1651. *L. divaricatus*, filamentosus pendulus angulatus articulatus tomentosus, ramis divaricatis, peltis orbiculatis sessilibus. S. N. 964. Dill. Musc. t. 12. f. 5. arb.
1652. *L. jubatus*, filamentosus pendulus axillis comprefsis. Fl. Sv. 1124. Fl. N. 212. Dill. Musc. t. 12. f. 7. n). arb.
1653. *L. lanatus*, filamentosus ramosissimus decumbens implicatus opacus. Fl. Sv. 1125. Fl. N. 773. Dill. Musc. t. 13. f. 8. lap.
1654. *L. pubescens*, filamentosus ramosissimus decumbens implicatus nitidus. Fl. Sv. 1126. Fl. N. 774. Dill. Musc. t. 13. f. 9. lap.
1655. *L. chalybeiformis*, filamentosus ramosus divaricatus decumbens implicato-flexuosus. Fl. Sv. 1127. Fl. N. 805. Fl. D. 262. Dill. Musc. t. 13. f. 10. lap. lign.
1656. *L. hirtus*, filamentosus ramosissimus erectus, tuberculis farinaceis sparsis. Fl. Sv. 1128. Fl. N. 572. Dill. Musc. t. 13. f. 12. *Ufnea hirta* Hoffm. Lich. II. p. 17. t. 30. f. 1. o). arb. lign.

T 3

1657.

- m) Ab antecedente haud diversum docet Cl. Ehrhart Beytr. I. p. 145.
- n) Aetate in *chalybeiformem* mutari asserit Cel. Afzelius Diff. cit. p. 35.
- o) *L. hirtum a florido* haud differre credit Cl. Ehrh. Beytr. I. p. 145. contendit Cel. Afzelius, de *plicato* etiam dubius.

294 CRYPTOGRAMIA. ALGAE. Tremella.

1657. *L. floridus*, filamentosus ramosus erectus, scutellis radiatis. Fl. Sv. 1130. Fl. N. 739. Dill. Musc. t. 13. f. 13. *Usnea florida* Hoffm. Lich. II. p. 19. t. 30. f. 2. arb.

1658. *L. vulpinus*, filamentosus ramosissimus erectus fastigiatus inaequali angulosus. Fl. Sv. 1129. Fl. N. 421. Dill. Musc. t. 13. f. 16. lign.

* *dubii generis*

1659. *L. radiformis*, filamentosus teres ramosissimus glaber radiformis. S. N. 964. N. A. Upf. 4. p. 251. Fl. D. 713. p). sub cort. arb.

TREMELLA.

1660. *T. juniperina*, sessilis membranacea auriformis fulva. Fl. Sv. 1135. Fl. N. 569. *Tr. sagarum* Fl. D. 885. f. 3. an? junip.

-1661. *T. mesenterica*, sessilis membranaceo-gelatinosa congesta sinuosa aurea. Act. Holm. 1769. p. 249. Vaill. Bot. t. 14. f. 4. arb. semiputr.

1662. *T. Noëst*, plicata undulata. Fl. Sv. 1136. Fl. N. 753. Dill. Musc. t. 10. f. 14. Fl. D. 885. f. 1. pr.

1663. *T. crispa*, terrestris tenera crispa virescens. Schr. Lips. 1151. Willd. Berol. 1221. Dill. Musc. t. 10. f. 12. *Tr. intestinalis* Fl. D. 885. f. 2. terr.

1664. *T. sagarum* sessilis membranacea difformis venosa subfusca. Act. Holm. 1769. p. 249. Dill. Musc. t. 10. f. 15. *Tr. atra* Fl. D. 884. *Tr. arborea* Hudf. Angl. 2. p. 563. arb.

† 1665. *T. affinis*, sessilis suborbicularis concava flava. Fl. D. 840. f. 1. arb.

1666. *T.*

p) *L. radiformis* Weberi subterraneus an a nostro subcorticali diversus? Nostrum pro radice Sphaeriae suae 2194 habet Haller, quo jure?

1666. *T. orbicularis*, orbiculata concava viridis. A. Holm.
 1769. p. 249. arb.
 1667. *T. rubra*, sublobata, externe glabra sicca, interne
 liquore viscida. Berl. Beschäft. N. F. Fr. III. p. 354.
 t. 7. f. 7. 8. lign.
 1668. *T. agaricoides*, stipitata nigra inquinans, pileo
 subconvexo. A. Holm. 1769. p. 250. *Tr. turbinata*
 Hudf. Angl. 2. p. 563. Hall. St. Helv. 2219. t. 48.
 f. 8. *Limsvampen* Aët. Holm. 1762. p. 105. t. 3.
Peziza polymorpha Fl. D. 464. *Lycoperdon trunca-*
tum S. N. 981. Fag. Querc.
 1669. *T. verrucosa*, tuberculosa solida rugosa. Fl. Sv.
 1139. Fl. N. 1093. Dill. Musc. t. 10. f. 11.
 lap. inund.
 1670. *T. difformis*, subrotunda sinuato-difformis gela-
 tinosa. Fl. Sv. 1140. conferv. marin.
 1671. *T. hemisphaerica*, hemisphaerica sparsa. Fl. Sv.
 1141. Fl. N. 952. Weig. Obs. p. 39. t. 2. f. 3.
 rup. mar.
 1672. *T. viridis*, hemisphaerica viridis lubrica faxis ad-
 nata inundatis. Aët. Holm. 1769. p. 251. Willd.
 Berol. 1222. lap. rivul.
 1673. *T. purpurea*, subglobosa sessilis solitaria glabra.
 Fl. Sv. 1142. Dill. Musc. t. 18. f. 6. lign.
 1674. *T. granulosa*, granulata congesta sessilis. A. Holm.
 1769. p. 250. Mich. Nov. Gen. p. 105. t. 54. f. 5.
 lign.
 α. nigra.
 β. rubra. Sphaeria Mori Weig. Obs. p. 45. t. 2.
 f. 11. Fl. D. 840. f. 2. Hall. St. Helv. t. 47.
 f. 9.
 1675. *T. adnata*, rotunda imbricata livida. Fl. Sv. 1143.
 lap. mar.

FUCUS.

1676. *F. ferratus*, fronde plana dichotoma costata serrato-dentata, fructificationibus terminalibus tuberculosis. Fl. Sv. 1144. Fl. N. 44.
1677. *F. vesiculosus*, fronde plana dichotoma costata integerrima, vesiculis axillaribus geminis; terminalibus tuberculatis. Fl. Sv. 145. Fl. N. 191.
1678. *F. divaricatus*, fronde plana dichotoma integerrima, axillis divaricatis, vesiculis axillaribus geminis. Sp. Pl. 1627. q).
1679. *F. inflatus*, fronde plana dichotoma integerrima punctata ovato-lanceolata inflata; apice diviso, Fl. Sv. 1148. Fl. N. 471.
1680. *F. ceranoides*, fronde plana dichotoma integerrima punctata, fructificationibus tuberculatis bifidis terminalibus. Fl. Sv. 1146. Fl. N. 606.
- β . *lacerus* Spec. Pl. 1627.
1681. *F. spiralis*, fronde plana dichotoma integerrima punctata; inferne lineari-canaliculata, fructificationibus tuberculatis geminis, Fl. Sv. 1147. Fl. N. 602. Fl. D. 286.
1682. *F. canaliculatus*, fronde plana dichotoma integerrima lineari-canaliculata, fructificationibus tuberculatis bipartitis obtusis. S. N. 967. Fl. D. 214.
- β . *F. excisus*. Sp. Pl. 1627. Fl. N. 314.
1683. *F. distichus*, fronde plana dichotoma integerrima lineari, fructificationibus tuberculatis mucronatis. S. N. 967. Fl. N. 987. Fl. D. 351. *F. filiformis* Gmel. Fuc. t. 1. f. 1. Fl. N. 835.
- † 1684. *F. soboliferus*, frondibus membranaceis dichotomis dilatatis integerrimis ferratisque apice bifidis: laciniis divaricatis, inferioribus appendiculatis. Fl. D. 1065.

† 1685. *F.*q) Varietatem tantum *F. vesiculosi* credunt Hudson, Aszelius.

- † 1685. *F. pumilus*, fronde compressa dichotoma, axillis divaricatis, vesiculis subsolitariis axillis ultimis innatis, fructificationibus globosis terminalibus. Fl. D. 1066.
- † 1686. *F. nodosus*, fronde compressa dichotoma, foliis integerrimis distichis, vesiculis innatis solitariis dilatatis. Fl. Sv. 1149. Fl. N. 244. Fl. D. 146. Gmel. Fuc. t. 1. B.
- † 1687. *F. filiquosus*, stirpe filiformi dichotoma, frondibus membranaceis ensiformibus solitariis ferratis: terminalibus petiolo inflatis. Fl. Sv. 1150. Fl. N. 243. Fl. D. 106. Gmel. Fuc. t. 2. B.
- † 1688. *F. corallinus*, fronde tereti, ramis alternis subulatis, foliis filiformibus medio in vesiculas fusiformes distensis. Fl. D. 709.
- † 1689. *F. elongatus*, fronde filiformi compressa dichotoma articulata, geniculis tumidiusculis. S. N. 968. Fl. N. 1095.
1690. *F. loreus*, fronde filiformi compressa dichotoma undique tuberculata, S. N. 968. Fl. N. 988. Fl. D. 710. r).
1691. *F. foeniculaceus*, fronde filiformi ramosissima, vesiculis ovatis terminatis, foliolis multipartitis obtusis apice fructificantibus. S. N. 968. Fl. N. 685. Gmel. Fuc. t. 2. A. f. 2.
- β. *F. barbatus*. Sp. Pl. 1629.
- † 1692. *F. granulatus*, fronde filiformi ramosissima, ramulis acuminatis, vesiculis subrotundis cumulatis, ramis foliisque acutis omnibus innatis. S. N. 968. Fl. N. 1072. Fl. D. 591.
- T 5 1693. *F.*
- r) Icon Fl. D. 420. hujus Fuci initium. Cel. Vahl.

1693. *F. Selaginoides*, fronde filiformi ramosissima; ramis dichotomis, foliis subulatis alternis basi vesiculosis. Mant. Pl. I. p. 134. Fl. N. 746. Gmel. Fuc. t. 2. A. f. 1.
- † 1694. *F. ericoides*, fronde filiformi ramosissima hirta. Sp. Pl. 1631. Fl. N. 851. Gmel. Fuc. t. 11. f. 2.
1695. *F. aculeatus*, fronde filiformi compressa ramosissima, dentibus marginalibus subulatis alternis erectis. S. N. 969. Fl. D. 355. *F. virgatus* Fl. N. 93. Aët. Nidr. IV. t. 7.
β. muscoides. Sp. Pl. 1630. Fl. N. 1071. Gmel. Fuc. t. 12.
1696. *F. Lycopodioides*, fronde filiformi tereti subramosa obtecta undique setis. S. N. 969. Fl. N. 838. *Conserua squarrosa* Fl. D. 357. Fl. N. 838.
- † 1697. *F. hirsutus*, fronde filiformi tereti dichotoma obtecta undique hirsutie. S. N. 969. Fl. N. 408.
- † 1698. *F. discors*, fronde tereti inermi aculeatissima, foliis distichis subpinnatis lineari-lanceolatis serratis. S. N. 969.
1699. *F. Filum*, fronde filiformi subfragili opaca. Fl. Sv. 1153. Fl. N. 347. Fl. D. 821.
- † 1700. *F. lanosus*, frondibus capillaceis dichotomis ramosissimis scabris. S. N. 969. Fl. N. 687.
1701. *F. fastigiatus*, fronde filiformi dichotoma ramosissima fastigiata obtusa. Fl. Sv. 1152. Fl. N. 102. Fl. D. 393. Gmel. Fuc. t. 6. f. 1.
1702. *F. furcellatus*, fronde filiformi dichotoma ramosissima acuminata. S. N. 969. Fl. D. 419. Fl. N. 677. *F. caprinus* Fl. N. 311. A. Nidr. IV. t. 5. f. 4. *F. lumbricalis* Gmel. Fuc. t. 6. f. 2.
- † 1703. *F.*

- † 1703. *F. palmatus*, fronde palmata plana dichotoma, laciniis linearibus bifidis. Sp. Pl. 1630. s).
- † 1704. *F. miniatus*, frondibus membranaceis dichotomis enerviis digitato proliferis. Fl. D. 769.
1705. *F. ovinus*, caule brevissimo, foliis inaequalibus profunde fissis, laciniis inferne angustatis apice hinc fissis illinc dentatis. Fl. N. 310. A. Nidr. IV. t. 9.
- † 1706. *F. bullatus*, fronde palmata, foliis disco inflatis, apice divisis. Fl. D. 770. t).
1707. *F. lanceolatus*, fronde membranacea foliis linearilanceolatis enerviis, margine apiceque proliferis.
 β *F. epiphyllus*. Fl. D. 708.
1708. *F. digitatus*, fronde palmata, foliolis ensiformibus, stirpe tereti. M. Plant. I. p. 134. Fl. D. 392.
F. palmatus Gmel. Fuc. t. 30. *F. hyperboreus* Fl. N. 61. t. 3.
 β *F. bifurcatus*. Fl. N. 312. A. Nidr. IV. t. 6.
- † 1709. *F. esculentus*, fronde simplici indivisa ensiformi, stirpe pinnata tetragona folium percurrente. M. Plant. I. p. 135. Fl. D. 417. *F. pinnatus* Fl. N. 313. A. Nidr. IV. t. 8. t. 1. *F. fimbriatus* Gmel. Fuc. t. 29, f. 1.
1710. *F. saccharinus*, fronde subsimplici ensiformi, stirpe tereti brevissima. Fl. Sv. 1151. Fl. N. 116. Fl. D. 416. Gmel. Fuc. t. 27. 28. *Ulva longissima* Fl. N. 1003. t. 7. f. 2.

1711. *F.*

s) *F. palmatus*, *ovinus*, *miniatus*, *rubens*, *lanceolatus* definitione certiore egent, ut quid revera sit species vel varietas innotescat, id quod ex Herbario Linneano haud difficile erit. In edit. XIII. Syst. Nat. conjungitur *F. ovinus* cum *palmato*, an recte? *F. lanceolatus* figura bene convenit cum *F. crenato* Gmel, at folia in *lanceolato* omnino enervia.

z) Eundem cum *F. ovino* suspicor.

1711. *F. sanguineus*, frondibus membranaceis ovato-oblongis integerrimis petiolatis nervosis, caule tereti ramofo. M. Plant. I. p. 136. Fl. N. 752. Fl. D. 349. Gmel. Fuc. t. 24. f. 2.
1712. *F. roseus*, frondibus membranaceis subovatis lanceolatisque sublaciniatis serratis nervosis, caule tereti ramofo. *F. roseus* Fl. D. 652.
- † 1713. *F. ciliatus*, frondibus membranaceis lanceolatis proliferis ciliatis. M. Plant. I. p. 136. Fl. N. 782. Fl. D. 353. *F. caulescens* Gmel. Fuc. t. 20. f. 2.
1714. *F. crispus*, frondibus submembranaceis dichotomis: laciniis dilatatis. M. Plant. I. p. 134. Fl. N. 748. *F. ligulatus* Gmel. Fuc. t. 21. f. 3. *Ulva caprina* Fl. N. 1001. t. 6. f. 4. *Ulva delicatula* ibid. 1035. t. 8.
1715. *F. crispatus*, frondibus membranaceis sublineari-bus ramosissimis crispis coloratis. S. N. 970. Fl. D. 826.
1716. *F. alatus*, frondibus membranaceis subdichotomis costatis, laciniis alternis decurrentibus bifidis. M. Plant. I. p. 135. Fl. N. 750. Fl. D. 352. Gmel. Fuc. t. 25.
1717. *F. dentatus*, frondibus membranaceis enerviis alternatim pinnatifidis, sinibus obtusis, laciniis apice erosis. M. Plant. I. p. 135. Fl. N. 749. *F. pinnatifidus* Fl. N. 832. Fl. D. 354.
1718. *F. rubens*, frondibus membranaceis oblongis undulatis sinuatis, stirpe ramofo rereti. Spec. Pl. 1630. Fl. N. 629. Fl. D. 827. *F. palmetta* Gmel. Fuc. t. 22. f. 3. t. 23.
1719. *F. vittatus*, frondibus membranaceis divisiformibus dentato-crispis. S. N. 970. Fl. N. 454. u).
- † 1720. *F.*

u) An a ciliato diversus? Linné iconem Fl. D. 353. ad utramque attulit.

† 1720. *F. ramentaceus*, frondibus filiformibus simplicibus hinc ramentis foliaceis confertis. S. N. 970. Fl. N. 686. *F. barbatus* Fl. N. 1007. *Ulva sobolifera* Fl. D. 356. 837.

† 1721. *F. plumosus*, frondibus cartilagineis lanceolatis bipinnatis plumosis, caule filiformi compresso ramoso. M. Plant. I. p. 134. Fl. N. 747. Fl. D. 350. *F. pedunculatus* Fl. N. 972. t. 2. f. 8. *F. ptilotus* ibid. 1045. t. 2. f. 15.

† 1722. *F. abrotanifolius*, fronde filiformi compressa bipinnata, apicibus vesiculosis dilatatis terminatis fructificationibus hinc tuberculatis. S. N. 971. Fl. N. 684. *F. capensis* Gmel. Fuc. t. 17. f. 1.

† 1723. *F. cartilagineus*, fronde cartilaginea compressa supradecomposita: laciniis linearibus. S. N. 971. Fl. N. 864. t. 3. f. 5. *F. vindicatus* Fl. N. 975. *F. versicolor* Gmel. Fuc. t. 17. f. 2.

† 1724. *F. gigartinus*, fronde cartilaginea filiformi compressa dichotoma, fructificationibus globosis pedunculatis terminalibus subjacente arista. S. N. 971. Fl. N. 847. Fl. D. 394. ? w).

† 1725. *F. spermophorus*, fronde membranacea dichotoma compressa capillacea, fructificationibus pedunculatis lateralibus, foliis linearibus multifidis. S. N. 970. Fl. N. 414.

† 1726. *F. confervoides*, frondibus linearibus membranaceis compressis ramosis, fructificationibus sparsis sessilibus subrotundis. S. N. 971. Fl. N. 754. *F. verrucosus* Gmel. Fuc. t. 14. f. 1.

† 1727. *F. Scorpioides*, caule tereti ramoso, ramis alternis ramosissimis apice inflexis. Fl. D. 887. *F. longissimus* Gmel. Fuc. t. 13.

1728. *F.*

w) Desunt enim in islandico aristae.

1728. *F. albus*, cartilagineo - corneus teres ramis vagis multifidis, apicibus acutis. Fl. N. 757. Fl. D. 408.
F. plicatus Gmel. Fuc. t. 14. f. 2.

* *Fuci nondum satis determinati.*

- † 1729. *F. flagelliformis*, fronde filiformi subdichotoma ramosa, flagellis teretibus longis. Fl. D. 650.
 † 1730. *F. viridis*, stirpe tereti simplici, ramis oppositis ramosissimis capillaceis. Fl. D. 886.
 † 1731. *F. laciniatus*, frondibus ramosis membranaceis, ramis dilatatis. Fl. N. 1097. Gmel. Fuc. t. 22. f. 2.
 † 1732. *F. Oederi*, ramosissimus, ramis vagis approximatis teretibus, vesicis solidis sparsis caulinis. Fl. N. 809. Fl. D. 276.
 † 1733. *F. norvegicus*, fronde plana sublineari dichotoma, apicibus obtusis hinc incurvis, divaricata purpurea. Fl. D. 971. t. 3. f. 4.
 † 1734. *F. fascia*, fronde plana simplici libera aequali, basi angustiore. Fl. D. 768.
 † 1735. *F. clavatus*, dichotomus teres extremitatibus clavatis apice excavatis. Fl. Isl. 19.
 † 1736. *F. coronopifolius*, frondibus cartilagineis subdichotomis canaliculatis subtus et margine obsitis verrucis seu ramulorum rudimentis confertis. Fl. Isl. 19.

ULVA.

- † 1737. *U. umbilicalis*, plana orbicularis monophylla peltata coriacea. Sp. Pl. 1633. Fl. N. 959. Dill. Musc. t. 8: f. 3. mar.
 † 1738. *U. plicata*, orbicularis plicata sessilis polyphylla peltata membranacea. *U. plicata* Fl. D. 829. Aët. Hafn. X. p. 257. t. g. f. 9. mar.
 1739. *U. intestinalis*, tubulosa simplex. Fl. Sv. 1154. Fl. N. 955. Dill. Musc. t. 9. f. 7. mar. aqu. dulc.
 β. *prolifera* e lateribus. Fl. D. 763. f. 1.

1740. *U.*

1740. *U. spongiformis*, tubulosa ramosa teres, ramis simplicissimis. Fl. D. 763. f. 2. mar.
1741. *U. compressa*, tubulosa compressa ramosa. Fl. Sv. 1155. Fl. N. 956. Dill. Musc. t. 9. f. 8. mar.
- † 1742. *U. incrassata*, tubulosa? ramosa, ramis simplicissimis teretibus lubricis apicem versus crassioribus. Fl. D. 653. mar.
- † 1743. *U. clavata*, filiformis ramosa viridis, ramis ramulisque apice clavatis. Fl. D. 949. mar.
1744. *U. latissima*, oblonga plana undulata membranacea viridis. Fl. Sv. 1156. Fl. N. 115. mar.
 β. *U. maxima*. Fl. N. 1002. t. 7. f. 5. x).
1745. *U. Laetuca*, palmata prolifera membranacea, ramentis inferne angustatis. Fl. Sv. 1157. Fl. N. 958. Dill. Musc. t. 8. f. 1. mar.
1746. *U. lanceolata*, lanceolata plana. S. N. 972. Dill. Musc. t. 9. f. 5. mar.
1747. *U. Linza*, fronde oblonga bullata. Fl. Sv. 1158. Fl. N. 957. Fl. D. 889. Dill. Musc. t. 9. f. 6. mar.
- † 1748. *U. flaves cens*, gelatinosa subtereti uscula subramosa flaves cens, ramis brevissimis obtusis. Hudf. Fl. Angl. p. 570. mar.
1749. *U. pruniformis*, subglobosa solitaria intus succulenta. Fl. Sv. 1159. Fl. N. 737. t. 2. f. 6. 7. Weig. Obs. p. 40. t. 2. f. 4. lac.
1750. *U. granulata*, globosa farcta viridis. S. N. 972. M. Plant. 136. Fl. N. 878. *Conserua Pisum* Fl. D. 660. fluv. rip.
1751. *U. radicata*, sphaerica aggregata, radicibus ramosis, viridis. A. Holm. 1769. p. 251. *U. granulata* Sp. Pl. 1663. Fl. Sv. 1160. Dill. Musc. t. 10. f. 17. Fl. D. 705. *Tremella granulata* Hudf. Fl. Angl. 2. p. 566. inund.

CON.

*) Dubium etiamnum an Ulvae vel potius Fuci species.

CONFERVA.

1752. *C. rivularis*, filamentis simplicissimis aequalibus longissimis. Fl. Sv. 1161. Fl. N. 608. Fl. D. 881. Dill. Musc. t. 2. f. 1. riv.
1753. *C. fontinalis*, filamentis simplicissimis aequalibus digito brevioribus. Fl. Sv. 1163. Fl. N. 1048. Fl. D. 651. f. 3. Dill. Musc. t. 2. f. 3. font.
1754. *C. bullosa*, filamentis aequalibus ramosis bullas aëreas includentibus. Fl. Sv. 1166. Fl. N. 844. Dill. Musc. t. 3. f. 2. aqu.
1755. *C. canalicularis*, filamentis aequalibus versus basin ramosioribus. Fl. Sv. 1162. Fl. N. 842. Dill. Musc. t. 4. f. 15. canal.
- † 1756. *C. distorta*, filamentis aequalibus ramosis, ramis ramulisque distortis. Fl. D. 820. lac.
- † 1757. *C. hirta*, filamentis aequalibus viridibus hirsutis, ramis ramulisque simplicibus flexuosis. Fl. D. 947.
1758. *C. amphibia*, filamentis aequalibus ramosis exsiccatione abeuntibus in aculeos. Fl. Sv. 1164. Fl. N. 1085. Dill. Musc. t. 4. f. 17. hum.
1759. *C. littoralis*, filamentis aequalibus ramosissimis elongatis scabriusculis. Fl. Sv. 1170. Fl. N. 845. Dill. Musc. t. 3. f. 13. mar.
1760. *C. dichotoma*, filamentis aequalibus dichotomis. Sp. Pl. 1635. Fl. N. 839. Fl. D. 358. Dill. Musc. t. 3. f. 9. foss.
- † 1761. *C. nidifica*, filamentis simplicibus apice fasciculato-proliferis graniferisque. Fl. D. 761. mar.
1762. *C. cancellata*, filis ramosis, filamentis alternis brevibus multipartitis digitatis. Fl. Sv. 1171. Dill. Musc. t. 4. f. 22. mar.
1763. *C. elongata*, filiformis longissima fusca, ramis alternis simplicibus brevibus pallidis. Fl. Sv. 1172. Fl. N. 927. mar.
1764. *C. omissa*, capillacea ramosissima fusca. Fl. Sv. 1173. Fl. N. 928. mar.

1765. *C.*

1765. *C. reticulata*, filis reticulato-coadunatis. Fl. Sv. 1167. Fl. N. 735. Dill. Musc. t. 4. f. 14. fluv.

** *Filamentis articulatis.*

1766. *C. fluviatilis*, filis simplicissimis fetiformibus rectis geniculis crassioribus angulatis. Fl. Sv. 1168. Fl. N. 843. Dill. Musc. t. 7. f. 47. fluv.

- † 1767. *C. Linum*, filis simplicissimis geniculatis longissimis, geniculis cylindricis. Fl. D. 771. f. 2.

- † 1768. *C. nitida*, filis simplicibus longis articulatis. Fl. D. 819. aqu.

- † 1769. *C. jugalis*, filamentis geniculatis simplicibus saepe conjugatis intus globiferis. Fl. D. 883. lac.

- † 1770. *C. Pennatula*, filamentis geniculatis ramosis viridibus, articulis cylindricis tuberculatis pilosis, pilis sparsis fasciculatis. Fl. D. 945.

- † 1771. *C. fracta*, filamentis geniculatis ramosissimis viridibus, ramis dichotomis divaricatis, articulis cylindricis. Fl. D. 946.

- † 1772. *C. hirta*, filamentis aequalibus viridibus hirsutis, ramis ramulisque simplicibus flexuosis. Fl. D. 947.

- † 1773. *C. flexuosa*, filamentis geniculatis flexuosis, ramulis intimis pinnatis: externis longissimis simplicibus. Fl. D. 882. mar. litt.

- † 1774. *C. uncialis*, viridissima sericea, filamentis geniculatis ramosis: pinnis secundis. Fl. D. 771. f. 1. rup. mar.

1775. *C. gelatinosa*, filis ramosis moniliformibus, articulis globosis gelatinosis. Fl. Sv. 1165. Fl. N. 1086. Dill. Musc. t. 7. f. 42. 43. *Chara batrachosperma* Weis Crypt. Gott. p. 33. f. 1. 2. font. riv.

1776. *C. polymorpha*, filamentis geniculatis, ramis fasciculatis. Fl. Sv. 1174. Fl. N. 756.
 α. *fascicularis*. Dill. Musc. t. 6. f. 35.
 β. *barbata*. Fl. Isl. p. 20. mar.
 γ. *corallina*. Fl. Isl. p. 20. mar.
 δ. *congesta*. Fl. D. 395. mar.
 ε. *caudata*. Fl. D. 651. f. 1. A. Holm. 1769. p. 251. Dill. Musc. t. 5. f. 33. riv. fluv.
1777. *C. seposita*, capillacea ramosissima pallida, geniculis purpureis. Fl. Sv. 1175. Fl. N. 929. Dill. Musc. t. 7. f. 41. mar.
1778. *C. vagabunda*, filis geniculatis flexuosis, ramis ramulisque brevioribus. Fl. Sv. 1177. Fl. N. 854. Dill. Musc. t. 5. f. 32. mar.
1779. *C. glomerata*, filis geniculatis, ramulis brevioribus multifidis. Sp. Pl. 1637. Fl. N. 1091. Fl. D. 651. f. 2. Dill. Musc. t. 5. f. 31. font. riv.
1780. *C. rupestris*, filis geniculatis ramosissimis viridibus. Fl. Sv. 1176. Fl. N. 853. t. 3. f. 3. Fl. D. 948. Dill. Musc. t. 5. f. 29. mar.
1781. *C. floccosa*, filamentis ramosis geniculatis pinnatis, pinnis oppositis, geniculis quadrangulis. Fl. D. 828. f. 1. mar.
- † 1782. *C. hypnoides*, filamentis pinnatis subramosis procumbentibus, pinnis geniculatis tremulis. Fl. D. 828. f. 2. lign. putr. terr. ul.
- † 1783. *C. diaphana*, filamentis geniculatis ramosissimis, ramis dichotomis apice forcipatis, septis ruberrimis, articulis diaphanis. Lightf. Fl. Scot. II. p. 996. Fl. D. 951. rup. mar.
1784. *C. Aegagropila*, filis geniculatis ramosissimis e centro confertissimis globum constituentibus. Fl. Sv. 1169. Fl. N. 1087. lac.
- † 1785. *C. stellaris*, filis e basi orbiculari parallelis. Fl. D. 660. f. 1. hum.

CORALLINA y).

1786. *C. officinalis*, trichotoma, articulis cylindricis terminali-ovato. Fn. Svec. 2234. Ell. Cor. t. 24. f. A. mar.
1787. *C. squamata*, bipinnata, articulis obovatis compressis. Fn. Sv. 2235. Ell. Cor. t. 24. f. C. mar.
1788. *C. corniculata*, dichotoma, articulis bicornibus: corniculis articulatis. Fn. Svec. 2236. Ell. Cor. t. 24. f. D. mar.
1789. *C. rubens*, dichotoma capillaris, articulis cylindricis brevissimis: dichotomiae subclavatis. Fn. Svec. 2237. Ell. Cor. t. 24. f. E. mar.
1790. *C. cristata*, dichotoma capillaris: articulis omnibus clavatis. Fn. Svec. 2238. Ell. Cor. t. 24. f. F. mar.

BYSSUS.

1791. *B. septica*, capillacea molliissima parallela fragilissima pallida. S. N. 973.
1792. *B. aquatica*, hyalina filamentis simplicibus apice clavatis. Fl. D. 896. inf. putr.
1793. *B. Flos Aquae*, filamentis plumosis natantibus. Fl. Sv. 1182. Fl. N. 908. aqu.
1794. *B. cancellata*, filis exacte undique cancellata. S. N. 974. aqu.
1795. *B. phosphorea*, lanuginosa violacea lignis adhaesens. Fl. Sv. 1186. Fl. N. 912. Dill. Musc. t. 1. f. 6.
1796. *B. velutina*, capillacea viridis, filamentis ramosis. Fl. Sv. 1183. Fl. N. 910. Dill. Musc. t. 1. f. 14. terr.

U 2

1797. *B.*

y) *Corallinae*, ab ill. a Linné ad *Zoophyta* relatae, ex observationibus Nob. Cavolini ad vegetabile regnum potius pertinent et *Confervis articulatis* nobis proximae videntur. Conf. CAVOLINI Mem. III. p. 254 et sequ.

1797. *B. bombycina*, lanuginosa altissima, filamentis longioribus implexis, lignis adnascens. A. Holm. 1769. p. 251.

1798. *B. aurea*, capillacea pulverulenta, fructificationibus sparsis, filamentis simplicibus ramosisque. S. N. 974. Fl. D. 718. f. 1. Dill. Musc. t. 1. f. 16. terr.

1799. *B. Cryptarum*, capillacea perennis cinerea tenax rupis innata. Fl. Sv. 1181. Fl. N. 1090.

1800. *B. antiquitatis*, pulverulenta atra. Fl. Sv. 1180. Fl. N. 1089. Dill. Musc. t. 1. f. 18. mr.

1801. *B. saxatilis*, pulverulenta cinerea rupes operiens. Fl. Sv. 1179. Fl. N. 909.

1802. *B. candelaris*, pulverulenta flava lignis adnascens. Fl. Sv. 1185. Fl. N. 911. Fl. D. 899. f. 2. Dill. Musc. t. 1. f. 4.

1803. *B. botryoides*, pulverulenta viridis terrae nudae. Fl. Sv. 1187. Fl. N. 913. Fl. D. 899. f. 3. Dill. Musc. t. 1. f. 5.

1804. *B. incana*, pulverulenta incana farinae instar strata terrae nudae. Fl. Sv. 1188. Fl. N. 1088. Dill. Musc. t. 1. f. 3.

1805. *B. lactea*, pulverulento²-crustacea albissima. Fl. Sv. 1189. Fl. N. 914. Fl. D. 840. f. 4. Dill. Musc. t. 1. f. 2. Cort. musc.

† 1806. *B. rosea*, (Lichen) pulverulenta conglomerata rosea. Schreb. Lipf. p. 140. 2).

† 1807. *B. fruticulosa*, candidissima, filamentis erectis bifurcatis. Fl. D. 718. f. 2. a). lign. putr.

4. FUNGI

2) Cel. Kölpin.

a) Numne potius *Mucoris*, species?

4 F U N G I.

AGARICUS.

* *Pileo albo.*

1808. *A. umbelliferus*, stipitatus, pileo plicato membranaceo, lamellis basi latioribus. Fl. Sv. 1192. Fl. N. 894. sylv.
1809. *A. androsaceus*, stipitatus albus, pileo plicato membranaceo, stipite nigro. Fl. Sv. 1193. Fl. N. 895. acer.
1810. *A. piperatus*, stipitatus, pileo planiusculo lactescente: margine deflexo, lamellis incarnato-pallidis. Fl. Sv. 1195. Fl. N. 896. sylv.
1811. *A. exstimatorius*, stipitatus, pileo campaniformi albido lacero, lamellis niveis, stipite subbulboso subulato nudo. Fl. Sv. 1196. Fl. N. 897. sylv.
β. cinctus, stipitatus pileo campanulato albo radio cinereo, lamellis albis ad medium ramosis. Act. Holm. 1769. p. 252. pr.
1812. *A. pratorum*, stipitatus, pileo convexo albido viscido, stipite longo annulato. Fl. Sv. 1198. Fl. D. 600. pr. sylv.
1813. *A. campestris*, stipitatus, pileo convexo squamato albido, lamellis rufis. Fl. Sv. 1203. Fl. N. 537. Fl. D. 714. pr. sep.
- † 1814. *A. Kremlinga*, stipitatus, pilei convexi viridescentis margine striato, stipite annulato basi tuberoso. Fl. N. 448. sylv.
- † 1815. *A. procerus*, stipitatus, pileo convexo squamato lamellis ex albido cinerascens; stipite annulato bulbosus. Fl. D. 772. b).

U 3

† 1816.

b) Anne mera *A. campestris* varietas?

919 CRYPTOGAMIA. FUNGI. Agaricus.

- †1816. *A. praealtus*, stipitatus, lamellis albis, pileo subviscido, stipiteque longo concolore. Fl. D. 830. f. 1. sylv.
- †1817. *A. praecox*, stipitatus, lamellis albis, pileo conico subfarinoso, stipite gracili longo. Fl. D. 830. f. 2. sylv.
- †1818. *A. gramineus*, stipitatus, lamellis rubris, pileo stipiteque brevi carnosus. Fl. D. 832. f. 1. gramin.
- †1819. *A. cernuus*, stipitatus, pileo hemisphaerico al-bido, lamellis cinereo-nigricantibus, stipite fistu-loso nudo curvato. Fl. D. 1008. f. 1. pr. ster.
- †1820. *A. niveus*, stipitatus, albidus, pileo conve-xiusculo umbilicato membranaceo, lamellis re-motis decurrentibus, stipite fistuloso nudo. Fl. D. 1015. sylv. acer.
1821. *A. nitidus*, stipitatus, pileo candicante, lamellis paucis, stipite fusco splendente. Fl. N. 996. Fl. D. 773. Vaill. Par. t. 11. f. 21-23. hort.
1822. *A. stellatus*, stipitatus, pileo planiusculo suban-gulato pellucido, lamellis paucioribus rectis. Act. Holm. 1769. p. 252. arb. annosae.

** *Pileo flavo.*

1823. *A. cinnamomeus*, stipitatus, pileo sordide flavo, lamellis luteo-rufis. Fl. Sv. 1205. Fl. N. 368. sylv.
- †1824. *A. Buillardii*, stipitatus, pileo cinnamomeo centro depresso margine crenato. Fl. D. 1069. f. 1.
- †1825. *A. rubescens*, stipitatus, pileo cinnamomeo centro depresso margine aequali, stipite solido, lactescens. Fl. D. 1069. f. 2.
1826. *A. dentatus*, stipitatus, pileo convexo, lamellis basi mucrone dentatis. Fl. Sv. 1206. pr.

1827. *A.*

1827. *A. Cantarellus*, stipitatus, lamellis ramosis decurrentibus. Fl. Sv. 1207. Fl. N. 806. Fl. D. 264. pr. sylv.
1828. *A. Georgii*, stipitatus, pileo flavo convexo, lamellis albis. Fl. Sv. 1210. Fl. N. 517. sylv.
1829. *A. deliciosus*, stipitatus, pileo testaceo succo lutescente. Fl. Sv. 1211. Fl. N. 999. pasco. sylv.
 β. *A. fulvus*, stipitatus, pileo hemisphaerico stipiteque cylindrico laevi fulvis, lamellis albis alternis dimidiatis. Act. Holm. 1769. p. 252.
1830. *A. clavus*, stipitatus, pileo luteo convexo striato, lamellis stipiteque albis. Fl. Sv. 1212. Fl. N. 364. nem.
- †1831. *A. caesareus*, stipitatus, volva maxima, pilei aurei hemisphaerici margine striato, stipite crasso annulato. Fl. N. 1000. acer.
1832. *A. citrinus*, stipitatus citrinus, pileo sinuato, lamellis alternis minoribus. Fl. N. 994. t. 7. f. 7.
1833. *A. aurantius*, stipitatus glutinosus, pileo convexo umbilicato. Fl. D. 599. pr.
1834. *A. squarrosus*, stipitatus, pileo hemisphaerico stipiteque squamosis testaceis: squamis reflexis, lamellis lutescentibus. Weig. Obs. p. 40. Fl. D. 491. sylv.
- †1835. *A. squamatus*, stipitatus, pileo flavo disco aurantio stipiteque squamosis, lamellis murinis.
A. scaber Fl. D. 831. f. 1. lign. putr.
- †1836. *A. conicus*, stipitatus, pileo membranaceo stipiteque cavo aurantiis, lamellis sulphureis. Fl. D. 633. sylv.
- †1837. *A. auratus*, stipitatus, pileo stipiteque flavis, lamellis fordide virentibus. Fl. D. 890. lign. putr.
- †1838. *A. mutabilis*, stipitatus, pileo planiusculo lutescente, lamellis ferrugineis, stipite solitario annulato fistuloso fibroso. Fl. D. 1008. pr.

- †1839. *A. glutinosus*, stipitatus, pileo subcampanulato striato lamellisq. flavis, stipiteq. fistuloso nudo concolore. Fl. D. 1009. f. 2. pr. sylv.
- †1840. *A. plicatus*, stipitatus, pileo planiusculo margine plicato luteo-rufo, lamellis pallidioribus, stipite longo rufo. Fl. D. 1009. f. 3. sylv.
- †1841. *A. cyathiformis*, stipitatus, pileo infundibuliformi luteo subfloccoso striato, lamellis pallidis decurrentibus, stipite solitario brevi nudo. Fl. D. 1011. sylv. acer.
- †1842. *A. membranaceus*, stipitatus, pileo infundibuliformi testaceo laevi membranaceo, lamellis decurrentibus pallidis, stipite fistuloso nudo superne attenuato. Fl. D. 1012. sylv. acer.
- †1843. *A. melleus*, stipitatus, pileo convexo lutescente fusco irrorato, lamellis pallidis, stipitibus aggregatis annulatis fæctis. Fl. D. 1013. sylv.
- †1844. *A. muricatus*, stipitatus, pileo convexiusculo flavo, lamellis albidis, stipitibus aggregatis fistulosis muricatis annulatis. Fl. D. 1015. f. 2. pr.
- †1845. *A. ochraceus*, sessilis, imbricatus, pileo obovato margine involuto. N. H. S. Skr. 2. 1. p. 27, Jacqu. Fl. Austr. t. 104. lign. putr.
- †1846. *A. nitens*, stipitatus, pileo subhemisphaerico margine subquamoso, vertice fastigiato flavo, lamellis cinereo-nigricantibus, stipite annulato. Fl. D. 1067. f. 1. ster. pecor.
- †1847. *A. cervinus*, stipitatus, pileo conico stipiteq. solido fusco-punctato, lamellis carneis. Fl. D. 1067. f. 2. lign. putr.
- †1848. *A. torminosus*, stipitatus, pileo piloso fulvente: centro depresso, marginibus revolutis, stipite inani. Fl. D. 1068. sylv.
- †1849. *A. unicolor*, stipitatus, pileo hemisphaerico fastigiato spadiceo, stipite annulato lamellisq. concoloribus. Fl. D. 1071. f. 1. pr.

†1850. *A.*

† 1850. *A. nivalis*, stipitatus, pileo planiusculo flavo, stipite fistuloso nudo lamellisque concoloribus. Fl. D. 1071. f. 2. alp.

† 1851. *A. resupinatus*, pileo resupinato velato, vertice elongato stipitiformi. Fl. D. 1072. lign. putr.

*** *Pileo nigricante.*

1852. *A. violaceus*, stipitatus, pilei rimosi margine violaceo tomentoso, stipite coerulescente lana ferruginea. Fl. Sv. 1226. Fl. N. 901. sylv.

† 1853. *A. echinatus*, stipitatus, atropurpureus squamatus, squamis confertis recurvis hinc aculeiformibus. Fl. N. 993. t. 7. f. 6. 9. 10.

† 1854. *A. purpureus*, stipitatus, totus violaceus, pileo subturbinato, lamellis ramosis decurrentibus. Fl. D. 655. f. 3. pr.

† 1855. *A. scaber*, stipitatus, pileo hemisphaerico fusco squamoso, lamellis stipiteque albis. Fl. D. 832. f. 3. sylv.

† 1856. *A. caesus*, stipitatus, pileo planiusculo fusco, lamellis griseo-fuscis, stipite inani nudo basi albido. Fl. D. 1010. pr.

† 1857. *A. cyamipes*, stipitatus, pileo convexo nigro, stipite inani basi crassiore lamellisque dilutioribus. Fl. D. 1071. f. 3. pr. alp.

† 1858. *A. bicolor*, stipitatus, pileo campanulato striato margine lacero lamellisque nigris, stipite fistuloso albo. Fl. D. 1070. pr. alp.

**** *Pileo rubro.*

1859. *A. lactifluus*, stipitatus, pileo plano carneo, lamellis rufis, stipite longo carneo. Fl. Sv. 1228. sylv.

1860. *A. viscidus*, stipitatus, pileo purpurascente fusco viscido, lamellis fusco purpurascensibus subdecurrentibus, stipite flavo. Fl. Sv. 1229. Fl. N. 903. sylv.

1861. *A. Kermesinus*, stipitatus, pileo lamellis stipiteque eoccineis. Fl. D. 715. sylv.
1862. *A. integer*, stipitatus, pilei rubri lamellis omnibus magnitudine aequalibus stipiteque albis. Fl. Sv. 1230. Fl. N. 902. Fl. D. 1009. f. 1. sylv.
1863. *A. barbatus*, stipitatus, pilei convexi pallide rubri radio barbato, lamellis simplicibus stipiteque albissimis. A. Holm. 1769. p. 252. sylv.
1864. *A. Apicii*, stipitatus, pileolo umbilicato rufescente, lamellis candidis, stipite fistuloso longo rufo. Fl. N. 804. pr. ster.
1865. *A. muscarius*, stipitatus, pileo rubro albo-squamoso, lamellis albis dimidiatis solitariis, stipite volvato apice dilatato, basi ovata. Fl. Sv. 1135. Fl. N. 642. sylv.

***** *Pileo cinereo.*

1866. *A. campanulatus*, stipitatus, pileo campanulato striato pellucido, lamellis adscendentibus, stipite nudo. Fl. Sv. 1213. Fl. N. 330. pr.
1867. *A. fimetarius*, stipitatus, pileo campanulato lacero, lamellis nigris lateraliter flexuosis, stipite fistuloso. Fl. Sv. 1215. Fl. N. 377. *A. comatus*. Fl. D. 834. fimet.
1868. *A. clypeatus*, stipitatus, pileo hemisphaerico viscido acuminato, lamellis albis, stipite longo cylindrico albo. Fl. Sv. 1216. Fl. N. 404. sylv.
1869. *A. mammosus*, stipitatus, pileo convexo acuminato, lamellis convexis griseis crenatis, stipite nudo. Fl. Sv. 1218. Fl. N. 898. sylv.
1870. *A. equestris*, stipitatus, pilei pallidi disco stellatim luteo, lamellis sulphureis. Fl. Sv. 1219. Fl. N. 900. pasc. sylv.
1871. *A. separatus*, stipitatus, pileo laevi livido, lamellis nigricantibus separatis, stipite bulboso volvato. Fl. Sv. 1220. Fl. N. 899. pasc.

1872. *A.*

1872. *A. quinquepartitus*, stipitatus, pileo subflavescente partito, lamellis albidioribus interne dentato connexis. Fl. Sv. 1222. pr.
1873. *A. fimbriatus*, stipitatus, pileo cinereo fusco margine reflexo in strias et lacinias erumpente, lamellis stipiteque albis. Fl. D. 891. lign. putr.
1874. *A. plumbeus*, stipitatus, pileo convexo cinereo, verrucis lamellisque albis, stipite basi crassiusculo, Fl. Sv. 1236. Fl. D. 1014. c).
- † 1875. *A. stellaris*, stipitatus, pileo orbiculari striato, striis binis basi contiguis, lamellis albidis, stipite laevi fistuloso. Fl. N. 955. t. 7. f. 8. fimet.
- † 1876. *A. mutabilis*, stipitatus, pileo juniore castaneo adultiore cinerascente, lamellis purpureis, stipite cavo. Fl. D. 774. sylv.
- † 1877. *A. crassipes*, stipitatus, pileo murino granulis albis, lamellis stipiteque basi incrassato albis, Fl. D. 831. sylv.
- † 1878. *A. crenulatus*, stipitatus, membranaceus coeruleo-lutescens sulcatus, centro papillari, stipite exili, Fl. D. 832. f. 2. fimet.

***** *Acaules dimidiati.*

- † 1879. *A. nigricans*, acaulis, subreniformis sordide violaceus, lamellis latis albis., Fl. D. 892, lign. putr.
- † 1880. *A. striatus*, acaulis, utrinque sordide rufus, supra striatus margineque subviolaceus. Fl. D. 1073. f. 1. lign. putr.
- † 1881. *A. pubescens*, acaulis, pubescens niveus, lamellis simplicibus. Fl. D. 1073. f. 2. rami putr.
- † 1882. *A. Quercinus*, acaulis, lamellis labyrinthiformibus. Fl. Sv. 1241. Fl. N. 907. arb.

1883. *A.*

c) An ab *A. crassipede* diversus?

Reliquos Agaricos stipitatos, nondum rite determinatos, sicco pede transeo.

1883. *A. Betulinus*, acaulis coriaceus villosus margine obtuso, lamellis ramosis anastomosantibus. Fl. Sv. 1244. Fl. D. 775. f. 1. arb.
1884. *A. Alneus*, acaulis, lamellis bifidis pulverulentis. Fl. Sv. 1242. Fl. N. 906. Weig. Obs. p. 41. t. 2. f. 6. arb.

BOLETUS.

- † 1885. *B. inuervus*, stipitatus, stipite livido suberoso, pileo ovato fusco extus papilloso. Fl. N. 540. arb. putr.
1886. *B. perennis*, stipitatus perennis, pileo utrinque planiusculo. Fl. Sv. 1245. Fl. N. 905. Fl. D. 1075. f. 2. arb. putr.
1887. *B. Polyporus*, stipitatus perennis, pileo subhemisphaerico subtus planiusculo fulvo. Act. Holm. 1769. p. 253. pr.
- † 1888. *B. lateralis*, stipitatus perennis, pileo dimidiato. Fl. D. 1075. f. 1. arb. putr.
1889. *B. bovinus*, stipitatus, pileo glabro pulvinato marginato, poris compositis acutis: porulis angulatis brevioribus. Fl. Sv. 1241. Fl. N. 133. Fl. D. 1074. pr. sylv.
1890. *B. rubiginosus*, stipitatus, pileo campanulato rubiginoso margine inflexo subtus albo, poris tenuissimis simplicibus papillofis, stipite cylindrico superne attenuato albo. A. Holm. 1769. p. 253. sylv.
1891. *B. luteus*, stipitatus, pileo pulvinato subviscido, poris rotundatis convexis flavissimis, stipite albo. Fl. Sv. 1247. Fl. N. 904. sylv.
1892. *B. viscidus*, stipitatus, pileo pulvinato viscido, poris teretibus convexis immersis distinctis, stipite lacero. Fl. Sv. 1248. Fl. N. 1053. sylv.

1893. *B. convexus*, stipitatus, pileo utrinque convexo margine inflexo, supra sordide fulvo, subtus griseo, stipite lacero fusco. N. Act. Holm. 1769. p. 253. sylv.
1894. *B. granulatus*, stipitatus, pileo viscido pulvinato, poris teretibus subangulatisque truncatis; angulo granulato. Fl. Sv. 1249. Fl. N. 104. sylv.
- † 1895. *B. gregarius*, stipitatus, gregarius, pileo convexiusculo irregulari submembranaceo, subtus flavescente, poris oblongis inaequalibus. Fl. D. 1018. pr.
- † 1896. *B. terreus*, stipitatus, pileo convexo supra murino, subtus stipiteque sordide albo, poris simplicibus. Fl. D. 833. f. 2.
1897. *B. subtomentosus*, stipitatus, pileo flavo subtomentoso, poris subangulatis difformibus fulvis planis, stipite flavo. Fl. Sv. 1251. sylv.
1898. *B. subsquamosus*, stipitatus, pileo albido, poris difformibus oblongo flexuosis niveis. Fl. Sv. 1250. Fl. N. 1055. sylv.
- † 1899. *B. resupinus*, stipitatus, stipite verticali, supra castaneo-fusco-undatus, subtus albidus, margine costato sinuatoque. Fl. D. 894. arb.
- † 1900. *B. frondosus*, subacaulis, frondosus caespitosus fuscus, frondibus imbricatis planiusculis flexuosis, poris albis. Dickf. Crypt. f. 1. p. 18. Fl. D. 952. arb.
- † 1901. *B. caudicinus*, acaulis, caespitosus carnosus lobatus, margine incrassato lobato, poris tenuissimis flavis. Fl. D. 1019. arb.
1902. *B. Buglossum*, acaulis, utrinque granulato-papillosum, papillis poruliferis, superne convexiusculus carneus, inferne planus subalbidus, substantia fibroso-gelatinosa. N. Act. Holm. 1769. p. 253. Querc.

1903. *B. fomentarius*, acaulis, pulvinatus inaequalis obtusus, poris teretibus aequalibus glaucis. Fl. Sv. 1252. Fl. N. 327. arb.
1904. *B. suberosus*, acaulis, pulvinatus albus laevis, poris acutis difformibus. Fl. Sv. 1253. Fl. N. 326. arb.
1905. *B. ignarius*, acaulis, pulvinatus laevis, poris tenuissimis. Fl. Sv. 1256. Fl. N. 334. Fl. D. 953. arb.
1906. *B. versicolor*, acaulis, fasciis discoloribus, poris albis. Fl. Sv. 1254. Fl. N. 1051. arb.
1907. *B. suaveolens*, acaulis, superne laevis salicinus odoratus. Fl. Sv. 1255. Fl. N. 1054. Salic.
- † 1908. *B. Favus*, acaulis, subpulvinatus scaber setis erectis ramosis, poris angulatis. S. N. 977. Fl. D. 893. ? arb.
- † 1909. *B. cellulofus*, crustaceus albus, poris cellulosis patulis. Fl. D. 716. f. 1. lign. putr.
- † 1910. *B. dubius*, crustaceus pallidus, poris sinuatis irregularibus patulis. *Merulius crispatus* Fl. D. 716. f. 2. d). lign. putr.

HYDNUM.

1911. *H. imbricatum*, stipitatum, pileo convexo imbricato. Fl. Sv. 1257. Fl. N. 797. Fl. D. 176. sylv.
- † 1912. *H. cyathiforme*, stipitatum, pileo infundibuliformi fasciis discoloribus, aculeis albidis. Fl. D. 1020. f. 2. sylv. acer.
1913. *H. repandum*, stipitatum, pileo convexo laevi flexuoso. Fl. Sv. 1258. Fl. N. 1041. Fl. D. 310. sylv.
1914. *H. tomentosum*, stipitatum, pileo plano-infundibuliformi. Fl. Sv. 1259. Fl. N. 1040. sylv.

1915. *H.*a) *B. cellulofus* et *dubius* an hujus generis?

1915. *H. Erinaceus*, stipitatum, pileo lobato-ramoso. Fl. D. 450. sylv.
1916. *H. Auriscalpium*, stipitatum, pileo dimidiato coriaceo. Fl. Sv. 1260. Fl. N. 1039. Fl. D. 1020. f. 1. sylv. acer.
1917. *H. gelatinosum*, stipitatum gelatinosum, pileo dimidiato supra glaberrimo. S. N. 978. *H. crystallinum* Fl. D. 717. sylv. acer.
1918. *H. parasiticum*, acaule arcuato-rugosum tomentosum. S. N. 978. Fl. D. 465. 534. f. 3. arb.
1919. *H. Thaelaephora*, acaule complicatum lacerum scabrum, supra fasciculatim pilosum subtus papillosum. *Thaelaephora Mesenteriformis* Wild. Fl. Berol. 1152. f. 15. Nat. Hist. Selfk. Skr. 2. 1. p. 25. terr. hum.

PHALLUS.

1920. *P. esculentus*, pileo ovato celluloso, stipite nudo rugoso. Fl. Sv. 1262. Fl. N. 547. Fl. D. 53. sylv.
1921. *P. impudicus*, volvatus, stipitatus, pileo celluloso. Fl. Sv. 1261. Act. Holm. 1742. t. 2. Fl. N. 346. Fl. D. 175. sylv.

CLATHRUS.

- † 1922. *C. cancellatus*, acaulis subrotundus. Sp. Pl. 1648. Fl. N. 979. Gled. Fung. t. IV. sylv.
1923. *C. nudus*, stipitatus, capitulo oblongo axi longitudinali adnato. Fl. Sv. 1263. Fl. N. 802. Fl. D. 216. lign. putr.
1924. *C. denudatus*, stipitatus, capitulo oblongo volvato. S. N. 979. Fl. N. 977. lign. putr.

1925. *C.*

320 CRYPTOGAMIA. FUNGI. Helvella.

1925. *C. recutitus*, stipitatus, capitulo globofo, glande ovali. Fl. Sv. 1264. Fl. N. 978. lign. putr.

1926. *C. ramosus*, stipitatus, capitulo difformi, stipite ramoso. Aët. Holm. 1769. p. 253. lign. putr.

α. aureus.

β. rufus.

HELVELLA.

1927. *H. Mitra*, pileo deflexo multilobato adnato, stipite fistuloso laevi. Afz. N. A. Holm. 1783. p. 303. Fl. Sv. 1265. *β.* Fl. D. 116. Fl. N. 177. sylv.

α. tomentosa.

β. nuda.

1928. *H. lacunosa*, pileo deflexo multilobato adnato, stipite fistuloso lacunis sulcisque irregularibus. Afz. N. Aët. Holm. 1783. p. 304. Fl. Sv. 1265. *α.* Gled. Fung. t. 2. f. aa. sylv.

α. lurida.

β. pallida.

1929. *H. sulcata*, pileo deflexo bilobato adnato, stipite solido sulcis irregularibus. Afz. N. A. Holm. 1783. p. 305. t. 10. f. 1. sylv.

α. fusca.

β. cinerea.

1930. *H. atra*, pileo deflexo bilobato libero subtus nudo, stipite solido. Afz. N. Aët. Holm. 1783. p. 307. Prodr. Fl. Scand. 1596. Fl. N. 1059. Fl. D. 534. f. 1. Fl. Isl. p. 20. sylv.

1931. *H. Pezizoides*, pileo deflexo bilobato libero subtus scabro, stipite fistuloso. Afz. N. Aët. Holm. 1783. p. 308. t. 10. f. 2. sylv.

† 1932. *H. aeruginosa*, minima, pileo difformi viridissima. Fl. D. 534. f. 2. arb. putr.

† 1933.

- † 1933. *H. triceps*, pileo trilobato libero, stipite fistuloso. *Phallus triceps* Fl. D. 835. fylv.
- † 1934. *H. lubrica*, pileo convexo margine nodoso-lobato, stipite fistuloso. *Phallus lubricus* Fl. D. 719. fylv.
1935. *H. revoluta*, pileo plano depresso margine revolutato integerrimo, stipite fistuloso compressiusculo. Afz. N. A. Holm. 1783. p. 309. Vaill. Par. t. 13. f. 7. 8. 9. fylv.
1936. *H. horizontalis*, pileo plano infundibuliformi margine horizontali crenulato, stipite solido tereti. Afz. N. A&. Holm. 1783. p. 311. t. 10. f. 3. fylv.
1937. *H. spathulata*, pileo verticali, basi adnata utrinque deorsum porrecto, stipite inserto. Afz. N. A. Holm. 1783. p. 312. *Clavaria spathulata* Prodr. Fl. Scand. 1603. Fl. D. 658. pasc. apr.
- † 1938. *H. verticalis*, stipitata; pileo verticali subrotundo pagina averſa plicato. N. H. Selfk. Skr. 2. 1. p. 26. ar. hum.

** *Forte ad alia genera amandanda.*

- † 1939. *H. membranacea*, acaulis membranacea difformis subfusca, margine varie lobato, subtus venosa: venis ramosis anastomosantibus. Dickf. crypt. F. I. p. 21. N. H. Selfk. Skr. 2. 1. p. 61. Fl. D. 1017. f. 1. N. A&. Hafn. 1. p. 286. f. 7. pal. musc.
1940. *H. Pineti*, acaulis. S. N. 979. Fl. D. 950. ? Pini putr.

CLAVARIA.

- † 1941. *C. pilipes*, simplicissima, gracilis, clavata solitaria, stipite inferne piloso. Fl. D. 1076. f. 1. sylv. ab.
1942. *C. simplex*, simplicissima caespitosa tubulosa subaequalis alba. N. Act. Holm. 1769. p. 254. pasc. apr.
- † 1943. *C. vermicularis*, simplicissima caespitosa farcta subclavata alba. Fl. D. 775. f. 2. sylv.
1944. *C. pistillaris*, simplicissima clavaeformis inanis. Fl. Sv. 1266. Fl. N. 890. Vaill. Par. t. 7. f. 5. sylv.
- † 1945. *C. media*, simplicissima clavata obtusa lutea. Fl. D. 837. f. 1. sylv.
1946. *C. ophioglossoides*, clavata integerrima compressa obtusa. S. N. 980. Fl. N. 1034. Fl. D. 1076. f. 2. Vaill. Par. t. 7. f. 3. sylv.
- † 1947. *C. inaequalis*, caespitosa compressa lutea apice acuminata. Fl. D. 836. f. 1. pr.
- † 1948. *C. purpurea*, caespitosa rufo-fusca subcompressa, apice acuminata. Fl. D. 837. f. 2. sylv.
- † 1949. *C. ascendens*, caespitosa, simplicissima, farcta, acuminata, alba. lign. putr.
- † 1950. *C. polymorpha*, caespitosa, simplex, rufa, apicibus sublobatis. Fl. D. 775. f. 1. sylv.
1951. *C. militaris*, clavata integerrima, capite squamoso polymorpha. S. N. 980. Fl. D. 657. f. 1. Vaill. Par. t. 7. f. 4. sylv.
1952. *C. Hypoxylon*, ramoso-cornuta compressa. Fl. Sv. 1267. Fl. N. 886. Weig. Obf. t. 3. f. 5. arb. putr.
- † 1953. *C. cornuta*, ramoso-cornuta, ramulis subverticillatis. Fl. D. 405. lign. putr.

† 1954.

1954. *C. Carchariae*, composita caulescens atra aspera, medulla fibrosa. Weig. Obf. p. 43. t. 3. f. 6. *C. digitata* e) Prodr. Fl. Scand. 1606. Fl. D. 900. lign. putr.
1955. *C. pileata*, stipite tereti laevi, pileo ovali adnato atro punctis elevatis scabro. Fl. D. 540. f). lign. putr.
1956. *C. coralloides*, ramis confertis ramosissimis inaequalibus. Fl. Sv. 1268. Fl. N. 887. Vaill. Par. t. 8. f. 4. sylv.
1957. *C. fastigiata*, ramis confertis ramosissimis fastigiatis obtusis luteis. Fl. Sv. 1269. Fl. N. 132. Ragi Syn. 3. t. 24. f. 5. sylv.
1958. *C. muscosa*, ramis ramosis acuminatis inaequalibus luteis. Fl. Sv. 1270. Fl. N. 888. Fl. D. 775. f. 3. 836. f. 2. sylv.

PEZIZA.

1959. *P. lentifera*, campanulata lentifera. Fl. Sv. 1271. Fl. N. 515. terr.
 α. *laevis*. Fl. D. 105.
 β. *hirsuta*. Fl. D. 469. f. 1.
 γ. *sericea*. Fl. D. 780. f. 1.
- † 1960. *P. granulifera*, globosa, tomentosa, cinerea, granulifera. *P. scutellaris* Fl. D. 780. f. 2. lign. putr.

X 2

1961. *P.*

e) *Cl. digitata* Linnei alius sine dubio fungus lignosus.

f) Haec et antecedens ad Sphaerias Halleri et Weigelii pertinent, haud bene Clavariis quidem adnumerantur, at difficilium cum Lichene pertuso et variis Lycoperdis junguntur.

1961. *P. punctata*, turbinata truncata, disco punctato. Fl. Sv. 1275. Fl. N. 811. Fl. D. 288. fim.
1962. *P. cornucopioides*, infundibuliformis, disco patente sinuato, punctato. S. N. 979. Fl. N. 743. Vaill. Par. t. 13. f. 2. 3. sylv.
1963. *P. Acetabulum*, cyathiformis, extus angulata venis ramosis. S. N. 979. Vaill. Par. t. 13. f. 1.
- † 1964. *P. pyxidata*, stipitata flavescens, disco cavo. Fl. D. 1016. f. 1. alp. musc.
- † 1965. *P. virgultorum*, stipitata flavescens, disco turbinato. Fl. D. 1016. f. 2. rami putr.
- † 1966. *P. denticulata*, stipitata flavescens extus glabra, disco planiusculo margine denticulato. Fl. D. 1016. f. 3. rami putr.
- † 1967. *P. parvula*, stipitata nivea extus hirsutula, disco cavo margine ciliato. Fl. D. 1016. f. 4. rami putr.
- † 1968. *P. Cibarium*, stipitata, cyathiformis, spadicea, margine integerrimo. Fl. D. 1078. f. 1. pal. alp.
- † 1969. *P. Phiala*, stipitata cyathiformis fordide lutea, margine integerrimo. Fl. D. 1078. f. 2. rami putr.
- † 1970. *P. turbinata*, carnosa sessilis turbinata subdiaphana albido-purpurascens, margine denticulato. Fl. D. 1017. f. 2. rami putr.
- † 1971. *P. carnosa*, carnosa sessilis, turbinata purpurea, margine undulato. Fl. D. 1017. f. 1. rami putr.
- † 1972. *P. cyathoides*, cyathiformis, margine obtuso erecto. Fl. Sv. 1274. Fl. N. 328. Raji Syn. 3. t. 24. f. 4. terr.
- † 1973. *P. scabra*, subfulva intus laevis nitida, extus granulosa. Fl. D. 655. f. 2. fimet.

1974. *P. cupularis*, globofo-campanulata, margine crenato. Fl. Sv. 1273. Fl. N. 760. Fl. D. 469. f. 3. sylv. exult.

† 1975. *P. urceolata*, sessilis urceolata cinerea, extus pilosiuscula. Fl. D. 1017. f. 3. lign. putr.

1976. *P. scutellata*, plana, margine convexo piloso. Fl. Sv. 1272. Fl. N. 329. Vaill. Bot. t. 13. f. 13. 14. lign. putr.

β. *albicans*. Fl. D. 656. f. 1. terr.

γ. *laevis*, non ciliata coccinea. Fl. D. 656. f. 2. sylv.

† 1977. *P. pusilla*, concava subcoerulescens subtus albidula, margine crenulato. *Elvela pusilla* Fl. D. 779. f. 1. lign. putr.

† 1978. *P. minuta*, concava flava, subtus lanuginosa alba. *Elvela minuta* Fl. D. 779. f. 2. lign. putr.

† 1979. *P. equina*, concava flava, extus margineque setosa. *Elvela equina* Fl. D. 779. f. 3. fim. equ.

1980. *P. cochleata*, turbinata cochleata. Sp. Pl. 1651. α. *fulva*. Fl. D. 657. f. 2. Vaill. Bot. t. 11. f. 8.

β. *ochracea*. Fl. D. 1077. f. 2. terr.

1981. *P. Auricula*, concava rugosa auriformis. S. N. 979. *Tremella Auricula* Fl. Sv. 1137. arb.

† 1982. *P. zonalis*, pileo patente membranaceo fibroso striato zonis picto. Fl. Isl. p. 20. *P. radiata* Fl. N. 1050. Fl. D. 469. f. 2.

LYCOPERDON.

* *Subterranea absque radice*.

1983. *L. Tuber*, globosum solidum muricatum, radice destitutum. Fl. Sv. 1221. Fl. N. 501. sylv.

326 CRYPTOGAMIA. FUNGI. Lycoperdon.

1984. *L. Semen*, globosum congestum rugosum. *L. olivaceum* Poll. III. p. 314. Hall. St. Helv. 2178. *Huitkals-Swampen* N. Ael. Holm. 1765. p. 208. fol. Brass.

** *Pulverulenta radicata.*

1985. *L. Bovista*, subrotundum lacerato-dehiscens. Fl. Sv. 1280. Fl. N. 143. pr. pasc.
 1986. *L. stellatum*, volva multifida patente perenni, capitula glabro: ore acuminato plicato. Fl. Sv. 1277. Mich. Gen. t. 100. f. 4. 5.
 1987. *L. multifidum*, volva multifida patente, evanida, radiis ovato-lanceolatis. Schreb. Lips. p. 113. *L. stellatum* Fl. N. 840. Fl. D. 360. pasc. arg.
 1988. *L. Carpopolus*, volva multifida, fructu globoso ex feminibus combinatis. S. N. 981. Fl. N. 998. Fl. D. 895. lign. putr.
 1989. *L. radiatum*, disco hemisphaerico, radio colorato. Sp. Pl. 1654. g). lign. putr.
 1990. *L. pedunculatum*, stipite longo, capitulo globoso glabro, ore cylindrico integerrimo. Fl. Sv. 1276. ar.

*** *Parasitica.*

1991. *L. cancellatum*, parasiticum foliorum, cancellatum filis parallelis. S. N. 981. Fl. D. 704. fol. Pyri.
 1992. *L. corniferum*, tuberculo flavo, tubulis curvatis integris. Fl. D. 833. fol. Sorbi.
 † 1993. *L. penicillatum*, tuberculo flavo, tubulis rectis in filamenta dehiscentibus. Fl. D. 839. fol. Pyri.

1994. *L.*

g) Cave ne cum Lichene Ulmi confundes, ut varii fecerunt.

1994. *L. aggregatum*, parasiticum ovato-oblongum congestum fuscum, ore irregulari, lana fulva elastice exsiliante, stipite nullo. A. Holm. 1769. p. 254. Obs. Bot. Fasc. I. p. 33. Hall. St. Helv. t. 48. f. 7. Fl. D. 655. f. 1. ? lign. putr.
- † 1995. *L. mammosum*, sessile subglobosum, cortice viridescente, farina atra. Fl. D. 1079. lign. putr.
1996. *L. sessile*, parasiticum subrotundum aggregatum, cortice farinaque flavo brunnea, ore lacero, stipite nullo. Aët. Holm. 1769. p. 254. lign. putr.
1997. *L. gregarium*, parasiticum congestum sphaericum cortice fusco nitente, farina aurea, demum pedunculatum. Obs. Bot. Fasc. I. p. 33. lign. putr.
1998. *L. stipitatum*, parasiticum, stipite flavo, capitulo globofo brunneo, ore irregulari. Aët. Holm. 1769. p. 254. Fl. D. 467. f. 3. rad. annos.
 β. Fl. D. 467. f. 1.
 γ. Fl. D. 467. f. 2.
1999. *L. variolosum*, parasiticum sessile subrotundum, cortice exteriore fulvo deciduo, farina atra compacta. S. N. 981. Dill. Musc. t. 18. f. 1. *Tremella carbonacea* Aët. Holm. 1769. p. 250. *Sphaeria Lycoperdoides* Weig. Obs. p. 47. t. 3. f. 2. 3. h). lign. putr.
2000. *L. pistiforme*, globosum scabrum ore perforato. S. N. 982.

X 4

2001. *L.*

h) Ex specimine quod mecum benevole communicavit amicis, Weigel.

2001. *L. Epidendrum*, cortice farinaque purpurea. Fl. Sv. 1279. Fl. N. 1042. Fl. D. 720. lign. putr.
2002. *L. epiphyllum*, aggregatum parasiticum, ore multifido lacero, pulvere fulvo. Fl. Sv. 1278. Fl. N. 1043. *L. poculiforme* Jacqu. Coll. I. p. 123. t. 4. f. 2. fol.
 β . *L. poculiforme*. Jacqu. Coll. I. p. 122. t. 4. f. 1. i).

MUCOR.

2003. *M. urceolatus*, stipitatus fugax, stipite superne ventricoso pellucido rorido, capitulo subrotundo elastico nigro. Dickf. Crypt. F. 1. p. 25. t. 3. f. 6. Fl. D. 1080. fim. equ.
2004. *M. Embolus*, seta nigra, villo fusco. Fl. Sv. 1288. Fl. N. 882. Hall. St. Helv. t. 48. f. 1. lign. putr.
2005. *M. obvelatus*, volva communi lacera, capitulo oblongo deciduo, calyce concavo pedicellato remanente. *Embolus obvelatus* Fl. D. 536. lign. putr.
- † 2006. *M. Scoparius*, fetis fasciculatim contortis, capitulis umbellatis. *Byffus Scoparia* Fl. D. 897. f. 1. putr.
- † 2007. *M. conoideus*, capitulo conico acuminato, stipite setaceo. *Byffus conoidea* Fl. D. 897. f. 2. infus.
2008. *M. Mucedo*, stipitatus, capsula globosa. Fl. Sv. 1282. putr.

2009. *M.*i) In foliis et fructu *Grossulariae* hortuli mei copiose.

2009. *M. glaucus*, stipitatus, capitulo subrotundo aggregato. Sp. Plant. 1656. Fl. D. 777. f. 2. k). fruct. putr.
2010. *M. Druparum*, stipitatus, capitulis oblongis alternis. Fl. D. 777. f. 1. Drup. putr.
2011. *M. crustaceus*, stipitatus, spicis digitatis. Fl. Sv. 1283. Fl. N. 889. l). esculenta.
2012. *M. viridescens*, viridis granulofus. Mant. Pl. 2. p. 313. lign. putr.
2013. *M. Erysiphe*, albus, capitulis fuscis sessilibus. Fl. Sv. 1291. Fl. N. 879. fol.
2014. *M. septicus*, mucosus flavus. Fl. Sv. 1282. 1285. Fl. D. 778. lign. putr.
- † 2015. *M. tubulosus*, hemisphaericus ex tubulis congestis.
- a. *Tubulifera Cremor*. Fl. D. 659. f. 1. lign. putr.
- β. *Tubulifera Ceratum*. Fl. D. 659. f. 2. fol. putr.

LACHNUM.

2016. *L. Agaricinum*, stipitatum, pileo subconcavo subtus stipiteque villosis. N. Aët. Holm. 1769. p. 255. *Peziza nivea* Dickf. Crypt. F. 1. p. 21. fylv.

X 5

* *Fungi*

k) Fl. D. 840. f. 3. aliena videtur.

l) *M. Scoparius* ab hoc diversus videtur.

* *Fungi dubii generis*

- †2017. *Byssus fruticulosa*, candidissima, filamentis erectis bifurcatis. Fl. D. 718. f. 2. lign. putr.
2018. *Acrospermum unguinosum*, perpendiculare, digitato-caespitosum, unguinosum, apice tumido. Fl. D. 1076. f. 3. lign. putr.
2019. *Acrospermum cylindricum*, teres, erectum, solitarium, rubellum apice pallidum. Fl. D. 1076. f. 4. lign. putr.
-

M U S C O R U M
G E N E R A E T S P E C I E S
SECUNDUM METHODUM
G E L. D. J. H E D W I G.

* Peristomate nullo.

PHASCUM. *Capsula* cum rudimento operculi nunquam
adscendente, sed inaperta de setula de-
cidente.

Ph. subulatum H. L.

Ph. cuspidatum.

Phascum acaulon α. L.

Ph. muticum H. Schr.

** Peristomate nudo.

SPHAGNUM. *Mas. Flos* clavaeformis. Folliculi intra
ejus foliorum alas, filamento succulento
cincti.

S. palustre.

HEDWIGIA. *M. Flos* gemmiformis monoicus.

H. ciliata H.

Bryum apocarpum coccineum L.

H. aquatica H.

Hypnum aquaticum L.

Gymno-

CLAVARIA.

† 1941. *C. pilipes*, simplicissima, gracilis, clavata solitaria, stipite inferne piloso. Fl. D. 1076. f. 1. sylv. ab.

1942. *C. simplex*, simplicissima caespitosa tubulosa subaequalis alba. N. Act. Holm. 1769. p. 254. pasc. apr.

† 1943. *C. vermicularis*, simplicissima caespitosa farcta subclavata alba. Fl. D. 775. f. 2. sylv.

1944. *C. pistillaris*, simplicissima clavaeformis inanis. Fl. Sv. 1266. Fl. N. 890. Vaill. Par. t. 7. f. 5. sylv.

† 1945. *C. media*, simplicissima clavata obtusa lutea. Fl. D. 837. f. 1. sylv.

1946. *C. ophioglossoides*, clavata integerrima compressa obtusa. S. N. 980. Fl. N. 1034. Fl. D. 1076. f. 2. Vaill. Par. t. 7. f. 3. sylv.

† 1947. *C. inaequalis*, caespitosa compressa lutea apice acuminata. Fl. D. 836. f. 1. pr.

† 1948. *C. purpurea*, caespitosa rufo-fusca subcompressa, apice acuminata. Fl. D. 837. f. 2. sylv.

† 1949. *C. ascendens*, caespitosa, simplicissima, farcta, acuminata, alba. lign. putr.

† 1950. *C. polymorpha*, caespitosa, simplex, rufa, apicibus sublobatis. Fl. D. 775. f. 1. sylv.

1951. *C. militaris*, clavata integerrima, capite squamoso polymorpha. S. N. 980. Fl. D. 657. f. 1. Vaill. Par. t. 7. f. 4. sylv.

1952. *C. Hypoxylon*, ramoso-cornuta compressa. Fl. Sv. 1267. Fl. N. 886. Weig. Obf. t. 3. f. 5. arb. putr.

† 1953. *C. cornuta*, ramoso-cornuta, ramulis subverticillatis. Fl. D. 405. lign. putr.

† 1954.

1954. *C. Carchariae*, composita caulescens atra aspera, medulla fibrosa. Weig. Obs. p. 43. t. 3. f. 6.
C. digitata e) Prodr. Fl. Scand. 1606. Fl. D. 900. lign. putr.
1955. *C. pileata*, stipite tereti laevi, pileo ovali adnato atro punctis elevatis scabro. Fl. D. 540. f). lign. putr.
1956. *C. coralloides*, ramis confertis ramosissimis inaequalibus. Fl. Sv. 1268. Fl. N. 887. Vaill. Par. t. 8. f. 4. sylv.
1957. *C. fastigiata*, ramis confertis ramosissimis fastigiatis obtusis luteis. Fl. Sv. 1269. Fl. N. 132. Ragi Syn. 3. t. 24. f. 5. sylv.
1958. *C. mustosa*, ramis ramosis acuminatis inaequalibus luteis. Fl. Sv. 1270. Fl. N. 888. Fl. D. 775. f. 3. 836. f. 2. sylv.

PEZIZA.

1959. *P. lentifera*, campanulata lentifera. Fl. Sv. 1271. Fl. N. 515. terr.
 α. *laevis*. Fl. D. 105.
 β. *hirsuta*. Fl. D. 469. f. 1.
 γ. *sericea*. Fl. D. 780. f. 1.
- †1960. *P. granulifera*, globosa, tomentosa, cinerea, granulifera. *P. scutellaris* Fl. D. 780. f. 2. lign. putr.

X 2

1961. *P.*

e) *Cl. digitata* Linnei alius sine dubio fungus lignosus.

f) Haec et antecedens ad Sphaerias Halleri et Weigeli pertinent, haud bene Clavariis quidem adnumerantur, at facilius cum Lichene pertuso et variis Lycoperdis junguntur.

1961. *P. punctata*, turbinata truncata, disco punctato. Fl. Sv. 1275. Fl. N. 811. Fl. D. 288. firm.
1962. *P. cornucopioides*, infundibuliformis, disco patente sinuato, punctato. S. N. 979. Fl. N. 743. Vaill. Par. t. 13. f. 2. 3. fylv.
1963. *P. Acetabulum*, cyathiformis, extus angulata venis ramosis. S. N. 979. Vaill. Par. t. 13. f. 1.
- † 1964. *P. pyxidata*, stipitata flavescens, disco cavo. Fl. D. 1016. f. 1. alp. musc.
- † 1965. *P. virgultorum*, stipitata flavescens, disco turbinato. Fl. D. 1016. f. 2. rami putr.
- † 1966. *P. denticulata*, stipitata flavescens extus glabra, disco planiusculo margine denticulato. Fl. D. 1016. f. 3. rami putr.
- † 1967. *P. parvula*, stipitata nivea extus hirsutula, disco cavo margine ciliato. Fl. D. 1016. f. 4. rami putr.
- † 1968. *P. Cibarium*, stipitata, cyathiformis, spadicea, margine integerrimo. Fl. D. 1078. f. 1. pal. alp.
- † 1969. *P. Phiala*, stipitata cyathiformis fordide lutea, margine integerrimo. Fl. D. 1078. f. 2. rami putr.
- † 1970. *P. turbinata*, carnosae sessilis turbinata subdiaphana albido-purpurascens, margine denticulato. Fl. D. 1017. f. 2. rami putr.
- † 1971. *P. carnosae*, carnosae sessilis, turbinata purpurea, margine undulato. Fl. D. 1017. f. 1. rami putr.
- † 1972. *P. cyathoides*, cyathiformis, margine obtuso erecto. Fl. Sv. 1274. Fl. N. 328. Raji Syn. 3. t. 24. f. 4. terr.
- † 1973. *P. scabra*, subfulva intus laevis nitida, extus granulosa. Fl. D. 655. f. 2. fimet.

1974. *P. cupularis*, globofo-campanulata, margine crenato. Fl. Sv. 1273. Fl. N. 760. Fl. D. 469. f. 3. sylv. exult.
- † 1975. *P. urceolata*, sessilis urceolata cinerea, extus pilosiuscula. Fl. D. 1017. f. 3. lign. putr.
1976. *P. scutellata*, plana, margine convexo piloso. Fl. Sv. 1272. Fl. N. 329. Vaill. Bot. t. 13. f. 13. 14. lign. putr.
- β. *albicans*. Fl. D. 656. f. 1. terr.
- γ. *laevis*, non ciliata coccinea. Fl. D. 656. f. 2. sylv.
- † 1977. *P. pufilla*, concava subcoerulefcent subtus albidā, margine crenulato. *Elvela pufilla* Fl. D. 779. f. 1. lign. putr.
- † 1978. *P. minuta*, concava flava, subtus lanuginosa alba. *Elvela minuta* Fl. D. 779. f. 2. lign. putr.
- † 1979. *P. equina*, concava flava, extus margineque setosa. *Elvela equina* Fl. D. 779. f. 3. fim. equ.
1980. *P. cochleata*, turbinata cochleata. Sp. Pl. 1651.
- α. *fulva*. Fl. D. 657. f. 2. Vaill. Bot. t. 11. f. 8.
- β. *ochracea*. Fl. D. 1077. f. 2. terr.
1981. *P. Auricula*, concava rugosa auriformis. S. N. 979. *Tremella Auricula* Fl. Sv. 1137. arb.
- † 1982. *P. zonalis*, pileo patente membranaceo fibroso striato zonis picto. Fl. Isl. p. 20. *P. radiata* Fl. N. 1050. Fl. D. 469. f. 2.

LYCOPERDON.

* *Subterranea absque radice.*

1983. *L. Tuber*, globosum solidum muricatum, radice destitutum. Fl. Sv. 1221. Fl. N. 501. sylv.

326 CRYPTOGAMIA. FUNGI. Lycoperdon.

1984. *L. Semen*, globosum congestum rugosum. *L. olivaceum* Poll. III. p. 314. Hall. St. Helv. 2178. *Huitkals-Swampen* N. Aët. Holm. 1765. p. 208. fol. Brass.

** *Pulverulenta radicata*.

1985. *L. Bovista*, subrotundum lacerato-dehiscens. Fl. Sv. 1280. Fl. N. 143. pr. pasc.
 1986. *L. stellatum*, volva multifida patente perenni, capitula glabro: ore acuminata plicata. Fl. Sv. 1277. Mich. Gen. t. 100. f. 4. 5.
 1987. *L. multifidum*, volva multifida patente, evanida, radiis ovato-lanceolatis. Schreb. Lipf. p. 113. *L. stellatum* Fl. N. 840. Fl. D. 360. pasc. arg.
 1988. *L. Carpopolus*, volva multifida, fructu globoso ex feminibus combinatis. S. N. 981. Fl. N. 998. Fl. D. 895. lign. putr.
 1989. *L. radiatum*, disco hemisphaerico, radio colorato. Sp. Pl. 1654. g). lign. putr.
 1990. *L. pedunculatum*, stipite longo, capitulo globoso glabro, ore cylindrico integerrimo. Fl. Sv. 1276. ar.

*** *Parasitica*.

1991. *L. cancellatum*, parasiticum foliorum, cancellatum filis parallelis. S. N. 981. Fl. D. 704. fol. Pyri.
 1992. *L. corniferum*, tuberculo flavo, tubulis curvatis integris. Fl. D. 833. fol. Sorbi.
 † 1993. *L. penicillatum*, tuberculo flavo, tubulis rectis in filamenta dehiscentibus. Fl. D. 839. fol. Pyri.

1994. *L.*

g) Cave ne cum Lichene Ulmi confundes, ut varii fecerunt.

1994. *L. aggregatum*, parasiticum ovato-oblongum congestum fuscum, ore irregulari, lana fulva elastice exsiliante, stipite nullo. A. Holm. 1769. p. 254. Obs. Bot. Fasc. I. p. 33. Hall. St. Helv. t. 48. f. 7. Fl. D. 655. f. 1. ? lign. putr.

† 1995. *L. mammosum*, sessile subglobosum, cortice viridescente, farina atra. Fl. D. 1079. lign. putr.

1996. *L. sessile*, parasiticum subrotundum aggregatum, cortice farinaeque flavo brunnea, ore lacero, stipite nullo. Aët. Holm. 1769. p. 254. lign. putr.

1997. *L. gregarium*, parasiticum congestum sphaericum cortice fusco nitente, farina aurea, demum pedunculatum. Obs. Bot. Fasc. I. p. 33. lign. putr.

1998. *L. stipitatum*, parasiticum, stipite flavo, capitulo globoso brunneo, ore irregulari. Aët. Holm. 1769. p. 254. Fl. D. 467. f. 3. rad. annos.
β. Fl. D. 467. f. 1.
γ. Fl. D. 467. f. 2.

1999. *L. variolosum*, parasiticum sessile subrotundum, cortice exteriori fulvo deciduo, farina atra compacta. S. N. 981. Dill. Musc. t. 18. f. 1. *Tremella carbonacea* Aët. Holm. 1769. p. 250. *Sphaeria Lycoperdoides* Weig. Obs. p. 47. t. 3. f. 2. 3. h). lign. putr.

2000. *L. pistiforme*, globosum scabrum ore perforato. S. N. 982.

X 4

2001. *L.*

h) Ex specimine quod mecum benevole communicavit amicis. Weigel.

328 CRYPTOGRAMIA. FUNGI. Mucor

2001. *L. Epidendrum*, cortice farinaque purpurea. Fl. Sv. 1279. Fl. N. 1042. Fl. D. 720. lign. putr.
2002. *L. epiphyllum*, aggregatum parasiticum, ore multifido lacero, pulvere fulvo. Fl. Sv. 1278. Fl. N. 1043. *L. poculiforme* Jacqu. Coll. I. p. 123. t. 4. f. 2. fol.
 β. *L. poculiforme*. Jacqu. Coll. I. p. 122. t. 4. f. 1. i).

MUCOR.

2003. *M. urceolatus*, stipitatus fugax, stipite superne ventricoso pellucido rorido, capitulo subrotundo elastico nigro. Dickf. Crypt. F. 1. p. 25. t. 3. f. 6. Fl. D. 1080. fim. equ.
2004. *M. Embolus*, seta nigra, villo fusco. Fl. Sv. 1288. Fl. N. 882. Hall. St. Helv. t. 48. f. 1. lign. putr.
2005. *M. obvelatus*, volva communi lacera, capitulo oblongo deciduo, calyce concavo pedicellato remanente. *Embolus obvelatus* Fl. D. 536. lign. putr.
- † 2006. *M. Scoparius*, fetis fasciculatim contortis, capitulis umbellatis. *Byffus Scoparia* Fl. D. 897. f. 1. putr.
- † 2007. *M. conoideus*, capitulo conico acuminato, stipite setaceo. *Byffus conoidea* Fl. D. 897. f. 2. infus.
2008. *M. Mucedo*, stipitatus, capsula globosa. Fl. Sv. 1282. putr.
2009. *M.*

i) In foliis et fructu *Grossulariae* hortuli mei copiose.

2009. *M. glaucus*, stipitatus, capitulo subrotundo aggregato. Sp. Plant. 1656. Fl. D. 777. f. 2. k). fruct. putr.

2010. *M. Druparum*, stipitatus, capitulis oblongis alternis. Fl. D. 777. f. 1. Drup. putr.

2011. *M. crustaceus*, stipitatus, spicis digitatis. Fl. Sv. 1283. Fl. N. 889. l). esculenta.

2012. *M. viridescens*, viridis granulofus. Mant. Pl. 2. p. 313. lign. putr.

2013. *M. Erysiphe*, albus, capitulis fuscis sessilibus. Fl. Sv. 1291. Fl. N. 879. fol.

2014. *M. septicus*, mucofus flavus. Fl. Sv. 1282. 1285. Fl. D. 778. lign. putr.

† 2015. *M. tubulofus*, hemisphaericus ex tubulis congestis.

a. *Tubulifera Cremor*. Fl. D. 659. f. 1. lign. putr.

β. *Tubulifera Ceratum*. Fl. D. 659. f. 2. fol. putr.

LACHNUM.

2016. *L. Agaricinum*, stipitatum, pileo subconcavo subtus stipiteque villofis. N. A&t. Holm. 1769. p. 255. *Peziza nivea* Dickf. Crypt. F. 1. p. 21. fylv.

X 5

* *Fungi*

k) Fl. D. 840. f. 3. aliena videtur.

l) *M. Scoparius* ab hoc diverfus videtur.

* *Fungi dubii generis*

- †2017. *Byssus fruticulosa*, candidissima, filamentis erectis bifurcatis. Fl. D. 718. f. 2. lign. putr.
2018. *Acrospermum unguinosum*, perpendiculare, digitato-caespitosum, unguinosum, apice tumido. Fl. D. 1076. f. 3. lign. putr.
2019. *Acrospermum cylindricum*, teres, erectum, solitarium, rubellum apice pallidum. Fl. D. 1076. f. 4. lign. putr.
-

M U S C O R U M
G E N E R A E T S P E C I E S
SECUNDUM METHODUM
G E L. D. J. H E D W I G.

* Peristomate nullo.

PHASCUM. *Capsula* cum rudimento operculi nunquam
adscendente, sed inaperta de fetula de-
cidente.

Ph. subulatum H. L.

Ph. cuspidatum.

Phascum acaulon α. L.

Ph. muticum H. Schr.

** Peristomate nudo.

SPHAGNUM. *Mas. Flos* clavaeformis. Folliculi intra
ejus foliorum alas, filamento succulento
cincti.

S. palustre.

HEDWIGIA. *M. Flos* gemmiformis monoicus.

H. ciliata H.

Bryum apocarpum coccineum L.

H. aquatica H.

Hypnum aquaticum L.

Gymno-

332 GENERA ET SPECIES MUSCORUM

GYMNOSTOMUM. *M. Flos* disciformis dioicus.

G. pyriforme H.

Bryum pyriforme L.

G. truncatulum H.

Bryum truncatulum L.

G. aestivum H.

Bryum aestivum L.

*** Peristomate figurato, ordine simplici.

TETRAPHIS. *M. Flos* capituliformis pulverulentus.
Peristoma dentibus quatuor pyramidalibus.

T. pellucida H.

Mnium pellucidum L.

LEERSIA. *M. Flos* gemmaeformis in foliorum alis.
Perist. dentibus 16 angustis erectilibus integris.

L. vulgaris H.

Bryum exstinctorium α . L.

L. ciliata H.

Bryum exstinctorium β . L.

GRIMMIA. *M. Flos* gemmaeformis in foliorum alis, disciformis in summitate. *Perist.* dentibus 16 latiusculis reflexis integris.

Gr. apocarpa H.

Bryum apocarpum aureum L.

WEISSIA. *M. Flos* capituliformis terminalis. *Perist.* dentibus 16 integris.

W. acuta H.

Bryum capillaceum Dickf.

W. controversa H.

Bryum viridulum L.

W. pusilla H.

Bryum paludosum L.

POLY-

POLYTRICHUM. *M. Flos* disciformis. *Perist.* dentibus 32 integris membrana apicibus adglutinata.

P. commune H.

P. commune α . L.

P. juniperinum H.

P. commune β . L.

P. piliferum H.

P. commune γ . L.

P. nanum H.

Mnium polytrichoides α . L.

P. aloides H.

Mnium polytrichoides β . L.

P. undulatum H.

Bryum undulatum L.

TRICHOSTOMUM. *M. Flos* gemmaeformis. *Peristoma* dentibus 16 bifidis longis erectis.

T. lanuginosum H.

Bryum hypnoides hirsutum L.

T. canescens H.

Bryum hypnoides pilosum et barbatum L.

FISSIDENS. *M. Flos* gemmaeformis. *Peristoma* dentibus 16 brevioribus inflexis.

F. bryoides H.

Hypnum bryoides L.

F. taxifolius H.

Hypnum taxifolium L.

F. adianthoides H.

Hypnum adianthoides L.

F. pulvinatum H.

Bryum pulvinatum L.

334 GENERA ET SPECIES MUSCORUM

DICRANUM. *M. Flos capituliformis terminalis. Perist. dentibus 16 brevibus, bifidis, inflexis.*

D. purpureum H.

Bryum purpureum L.

D. scoparium H.

Bryum scoparium L.

D. heteromallum H.

Bryum heteromallum L.

D. glaucum H.

Bryum glaucum L.

D. pellucidum H.

Bryum pellucidum L.

D. flexuosum H.

Bryum flexuosum L.

D. aciculare H.

Bryum aciculare L.

D. Celsii H.

Bryum Celsii L.

D. virens H.

Bryum virens Sw.

D. varium H.

Bryum simplex L.

TORTULA. *M. Flos gemmaeformis. Perist. ciliis spiraliter convolutis.*

T. tortifolia H.

Bryum tortuosum L.

BARBULA. *M. Flos capituliformis. Perist. ciliis spiraliter convolutis.*

B. ruralis H.

Bryum rurale L.

B. muralis H.

Bryum murale L.

B. unguiculata H.

Bryum unguiculatum L.

**** Peri-

**** Peristomate figurato, ordine duplici.

NECKERA. *M. Flos* gemmaeformis. *Peristoma* dentibus 16, ciliisque 16 apice liberis alternantibus.

N. crispá H.

Hypnum crispum L.

N. dendroides H.

Hypnum dendroides L.

N. curtipendula H.

Hypnum curtipendulum L.

N. viticulosa H.

Hypnum viticulosum L.

LESKEA. *M. Flos* gemmaeformis. *Perist. extern.* dentibus 16 acutis, *intern.* ciliis uniformibus basi membranacea connexis.

L. trichomanoides H.

Hypnum trichomanoides L.

L. complanata H.

Hypnum complanatum L.

L. sericea H.

Hypnum sericeum curvatum L.

L. palustris H.

Hypnum palustre L.

HYPNUM. *M. Flos* gemmaeformis. *Perist. extern.* dentibus 16 acutis, *intern.* ciliis e membrana difformibus.

H. denticulatum H. L.

H. sylvaticum H. L.

H. lucens H. L.

H. rugosum H. L.

H. triquetrum H. L.

H. rutabulum H. L.

H. filicinum H. L.

H. pro-

- H. proliferum* H. L.
H. delicatulum H. L.
H. parietinum H. L.
H. praelongum H. L.
H. Crista castrensis H. L.
H. abietinum H. L.
H. plumosum H. L.
H. cupressiforme H. L.
H. aduncum H. L.
H. Scorpioides H. L.
H. squarrosum H. L.
H. loreum H. L.
H. alopecurum H. L.
H. purum H. L.
H. illecebrum H. L.
H. riparium H. L.
H. cuspidatum H.
 Hypnum cuspidatum α . L.
H. compressum H.
 Hypnum cuspidatum β . L.
H. lutescens H. Schreb.
 Hypnum fericeum praelongum.
H. velutinum H. L.
H. serpens H. L.

BRYUM. *M. Flos capituliformis. Perist. extern. dentibus 16 acutis, intern. e membrana carinata ciliis difformibus.*

- B. androgynum* H.
 Mnium androgynum L.
B. argenteum H. L.
B. squarrosum H. L.
B. binum H. L.
B. caespiticium H. L.
B. carneum H. L.

MNIUM.

MNIUM. *M. Flos* disciformis. *Perist. extern.* dentibus 16 acutis, *intern.* e membrana ciliis difformibus.

Mn. fontanum H. L.

Mn. palustre H. L.

Mn. annotinum H. L.

Mn. hornum H. L.

Mn. crudum H. L.

Mn. serpyllifolium H.

Mnium serpyllifolium α . L.

Mn. cuspidatum H.

Mnium serpyllifolium β . L.

Mn. roseum H.

Mnium serpyllifolium γ . L.

Mn. ligulatum H.

Mnium serpyllifolium δ . L.

Mn. triquetrum H. L.

KOELREUTERA. *M. Flos* disciformis. *Perist. extern.* dentibus 16 apice cohaerentibus obliquis, *intern.* cilia totidem membranacea plana.

K. hygrometrica H.

Mnium hygrometricum L.

WEBERA. *Perist. extern.* dentibus 16 acuminatis, *intern.* membranaceo-carinatum cum et absque ciliis. *Flores* hermaphroditi terminales.

W. pyriformis H.

Bryum pyriforme L.

BARTHRAmia. *M. Flores* foeminei approximati. *Perist. extern.* dentibus 16 cuneiformibus, *intern.* membranaceum plicato-carinatum apice lacinulatum.

B. pomiformis H.

Bryum pomiforme L.

B. Halleriana H.

Bryum laterale Sw.

338 GENERA ET SPECIES MUSCORUM etc.

ORTHOTRICHUM. *M. Flos gemmaceus. Perist. extern.*
dentibus 16 latiusculis, *intern. dentibus 16*
erofis, *Calyptra conica pilis erectis.*

O. anomalum H.

Bryum striatum minus α . L.

O. striatum H.

Bryum striatum majus β . L.

O. crispum H.

Bryum striatum δ . L.

BUXBAUMIA. *M. Flos discoideus. Perist. exter. denti-*
bus 16 truncatis, *intern. membranaceo-*
plicatum.

B. aphylla H. L.

B. foliosa H. L.

FONTINALIS. *M. Flos gemmiformis. Perist. extern.*
dentibus 16, *intern. reticulatum.*

F. antipyretica H. L.

F. falcata H.

Fontinalis secunda Sw.?

MEESIA. *M. Flos disciformis. Perist. extern. denti-*
bus 16 curtis, *intern. reticulatum.*

M. uliginosa H.

Bryum trichodes L.

CONSPECTUS
 PLANTARUM
 PRODROMI
 FLORAE SCANDINAVIAE
 ORDINE LINNEANO.

I. MONANDRIA.

I. MONOGYNIA.

<i>Salicornia</i>	Pag. 1. 2
herbacea	—
fruticosa	—
<i>Hippuris</i>	1. 2
vulgaris	—
maritima	—

2. DIGYNIA.

<i>Callitriche</i>	1. 2
verna	—
autumnalis	—

II. DIANDRIA.

I. MONOGYNIA.

<i>Ligustrum</i>	3. 5
vulgare	—
<i>Circaea</i>	3. 5
lutetiana	—
alpina	—
<i>Veronica</i>	3. 5
* <i>Spicatae.</i>	
maritima	—
longifolia	—
spicata	—
hybrida	—
officinalis	—

** *Corymbofo-race-
mosae.*

fruticulosa	Pag. 6
alpina	—
serpyllifolia	—
Beccabunga	—
Anagallis	—
scutellata	—
Teucrium	—
montana	—
Chamaedrys	—

*** *Pedunculis unifloris.*

agrestis	—
arvensis	—
hederaefolia	7
triphyllus	—
verna	—
acinifolia	—
peregrina	—
marilandica	—
<i>Gratiola</i>	
officinalis	3. 7
<i>Pinguicula</i>	
vulgaris	3. 7
alpina	—
villosa	—

Y 2

Utri.

<i>Utricularia</i>	Pag. 3.	7	fetaceus	Pag. 16
vulgaris	—	—	caricis	—
minor	—	—	maritimus	—
<i>Verbeña</i>	4.	8	sylvaticus	—
officinalis	—	—	<i>Eriophorum</i>	10. 16
<i>Lycopus</i>	4.	8	vaginatum	—
europaeus	—	—	polystachyon	17
<i>Salvia</i>	4.	8	alpinum	—
pratensis	—	—	<i>Nardus</i>	10. 17
2. DIGYNIA.			stricta	—
<i>Anthoxanthum</i>	4.	8		
odoratum	—	—	2. DIGYNIA.	
III. TRIANDRIA.			<i>Asprella</i>	10. 17
1. MONOGYNIA.			Oryzoides	—
<i>Valeriana</i>	9.	14	<i>Phalaris</i>	10. 17
officinalis	—	—	phleoides	—
dioica	—	—	arundinacea	—
locusta	—	—	<i>Panicum</i>	10. 17
α. olitoria			verticillatum	—
β. radiata?			glaucum	—
<i>Iris</i>	9.	14	viride	18
germanica	—	—	Crus Galli	—
Pseudacorus	—	—	sanguinale	—
fibirica	—	—	<i>Phleum</i>	10. 18
spuria	—	—	pratense	—
<i>Schoenus</i>	9.	14	nodosum	—
Mariscus	—	—	alpinum	—
nigricans	—	—	arenarium	—
ferrugineus	15	—	<i>Alopecurus</i>	10. 19
fuscus	—	—	pratensis	—
albus	—	—	agrestis	—
<i>Cyperus</i>	9.	15	geniculatus	—
flavescens	—	—	panicus	—
fuscus	—	—	<i>Milium</i>	11. 19
<i>Scirpus</i>	9.	15	effusum	—
palustris	—	—	<i>Agrostis</i>	11. 19
Baeothryon	—	—	* aristatae.	—
caespitosus	—	—	spica venti	—
acicularis	—	—	arundinacea	—
fluitans	16	—	rubra	—
lacustris	—	—	canina	—
Holoschaenus	—	—	β. minor	—

** muticae.			Briza	Pag. II. 23
stolonifera	Pag. 20		media	—
α . erecta			Cynofurus	II. 23
β . decumbens			cristatus	—
γ . pumila			coeruleus	—
alba	—		Festuca	12. 24
flava	—		ovina	—
capillaris	—		α . mutica	—
Aira	II. 20		β . vivipara	—
* muticae.			bromoides	—
aquatica	—		rubra	—
** aristatae.			duriuscula	—
subspicata	—		dumetorum	—
caespitosa	21		elatior	—
flexuosa	—		decumbens	—
montana	—		adscendens	—
alpina	—		fluitans	—
canescens	—		Bromus	12. 25
praecox	—		secalinus	—
cariophyllaea	—		α . vitiosus	—
Melica	II. 21		β . hordeaceus	—
ciliata	—		mollis	—
nutans	—		β . nanus	—
uniflora	—		inermis	—
coerulea	22		asper	—
Poa	II. 22		littoreus	—
aquatica	—		sterilis	26
alpina	—		arvensis	—
β . vivipara			tectorum	—
trivialis	—		giganteus	—
angustifolia	—		triflorus	—
pratensis	—		pinnatus	—
annua	—		cristatus	—
salina	—		Stipa	12. 26
palustris	—		pennata	—
compressa	—		Avena	12. 27
nemoralis	23		elatior	—
bulbosa	—		strigosa	—
arenaria	—		fatua	—
cristata	—		flavescens	—
glauca	—		pubescens	—
Dactylis	II. 23		pratensis	—
glomerata	—		Y 3	Arundo

<i>Arundo</i>	Pag. 12. 27	<i>Asperula</i>	Pag. 30. 33
<i>phragmites</i>	—	<i>odorata</i>	—
<i>epigejos</i>	—	<i>tinctoria</i>	34
<i>neglecta</i>	—	<i>Galium</i>	30. 34
<i>Calamagrostis</i>	—	<i>palustre</i>	—
<i>arenaria</i>	28	<i>trifidum</i>	—
<i>Lolium</i>	12. 28	<i>uliginosum</i>	—
<i>perenne</i>	—	<i>saxatile</i>	—
<i>β. - - -</i>	—	<i>pomeranicum</i>	—
<i>γ. - - -</i>	—	<i>verum</i>	—
<i>temulentum</i>	—	<i>Mollugo</i>	35
<i>β. mutica</i>	—	<i>β. angustifolia</i>	—
<i>Rottböllia</i>	28	<i>sylvaticum</i>	—
<i>incurvata</i>	—	<i>boreale</i>	—
<i>Elymus</i>	12. 28	<i>Aparine</i>	—
<i>arenarius</i>	—	<i>Plantago</i>	30. 35
<i>caninus</i>	—	<i>major</i>	—
<i>Hordeum</i>	12. 28	<i>media</i>	—
<i>murinum</i>	—	<i>lanceolata</i>	—
<i>secalinum</i>	—	<i>β. altissima</i>	—
<i>Triticum</i>	12. 29	<i>γ. capitata</i>	—
<i>repens</i>	—	<i>maritima</i>	36
<i>juncum</i>	—	<i>β. squamata</i>	—
3. TRIGYNIA.		<i>γ. serpentina</i>	—
<i>Monia</i>	13. 29	<i>ferraria</i>	—
<i>fontana</i>	—	<i>coronopifolia</i>	—
<i>Holosteum</i>	13. 29	<i>Plyllium</i>	—
<i>umbellatum</i>	—	<i>Centunculus</i>	31. 36
<i>Koenigia</i>	13. 29	<i>minimus</i>	—
<i>islandica</i>	—	<i>Sanguisorba</i>	31. 36
IV. TETRANDRIA.		<i>officinalis</i>	—
I. MONOGYNIA.		<i>Cornus</i>	31. 36
<i>Globularia</i>	30. 33	<i>sanguinea</i>	—
<i>vulgaris</i>	—	<i>suecica</i>	—
<i>Dipsacus</i>	30. 33	<i>Isardia</i>	31. 37
<i>sylvestris</i>	—	<i>palustris</i>	—
<i>pilosus</i>	—	<i>Trapa</i>	31. 37
<i>Scabiosa</i>	30. 33	<i>natans</i>	—
<i>succisa</i>	—	<i>Alchemilla</i>	31. 37
<i>arvensis</i>	—	<i>vulgaris</i>	—
<i>columbaria</i>	—	<i>alpina</i>	—
<i>Sherardia</i>	30. 33	<i>Aphanes</i>	—
<i>arvensis</i>	—		

2. DIGYNIA.

<i>Cuscuta</i>	Pag. 32. 37
<i>europaea</i>	—
<i>Epithymum</i>	—

3. TRIGYNIA.

<i>Ilex</i>	32. 37
<i>Aquifolium</i>	—
<i>Potamogeton</i>	32. 38
<i>natans</i>	—
<i>perfoliatum</i>	—
<i>lucens</i>	—
<i>crispum</i>	—
<i>compressum</i>	—
<i>pectinatum</i>	—
<i>gramineum</i>	—
<i>marinum</i>	—
<i>pusillum</i>	—
<i>Ruppia</i>	32. 38
<i>maritima</i>	—
<i>Sagina</i>	32. 39
<i>procumbens</i>	—
<i>erecta</i>	—
<i>Tillaea</i>	32. 39
<i>aquatica</i>	—

V. PENTANDRIA.

I. MONOGYNIA.

<i>Myosotis</i>	40. 53
<i>scorpioides</i>	—
<i>α. arvensis</i>	—
<i>β. palustris</i>	—
<i>lappula</i>	—
<i>Lithospermum</i>	40. 53
<i>officinale</i>	—
<i>arvense</i>	—
<i>Anchusa</i>	40. 53
<i>officinalis</i>	—
<i>Cynoglossum</i>	40. 53
<i>officinale</i>	—
<i>Pulmonaria</i>	40. 53
<i>officinalis</i>	—
<i>α. - - - -</i>	—
<i>β. - - - -</i>	—

<i>angustifolia</i>	Pag. 54
<i>maritima</i>	—
<i>Symphytum</i>	40. 54
<i>officinale</i>	—
<i>Asperugo</i>	41. 54
<i>procumbens</i>	—
<i>Lycopsis</i>	41. 54
<i>arvensis</i>	—
<i>Echium</i>	41. 54
<i>vulgare</i>	—
<i>Diapensia</i>	41. 54
<i>lapponica</i>	—
<i>Androsace</i>	41. 54
<i>septentrionalis</i>	—
<i>α. acaulis</i>	—
<i>Primula</i>	41. 55
<i>veris</i>	—
<i>α. officinalis</i>	—
<i>β. elatior</i>	—
<i>γ. acaulis</i>	—
<i>farinosa</i>	—
<i>norvegica</i>	—
<i>Menganchus</i>	41. 55
<i>Nymphoides</i>	—
<i>trifoliata</i>	—
<i>Horstonia</i>	42. 55
<i>palustris</i>	—
<i>Lythymachia</i>	42. 55
<i>vulgaris</i>	—
<i>thyrsiflora</i>	56
<i>punctata</i>	—
<i>nemorum</i>	—
<i>Nummularia</i>	—
<i>Anagallis</i>	42. 56
<i>arvensis</i>	—
<i>Azalea</i>	42. 56
<i>lapponica</i>	—
<i>procumbens</i>	—
<i>Convolvulus</i>	42. 56
<i>arvensis</i>	—
<i>sepium</i>	—
<i>Polemonium</i>	42. 57
<i>coeruleum</i>	—
Y 4	Campa-

<i>Campanula</i>	Pag. 42.	57	<i>Ribes</i>	Pag 44.	60
uniflora	—	—	rubrum	—	—
rotundifolia	—	—	alpinum	—	—
β. parviflora	—	—	petraeum	—	—
patula	—	—	nigrum	—	—
Rapunculus	—	—	Groffularia	—	—
perficifolia	—	—	Uva crispa	—	—
latifolia	—	—	<i>Hedera</i>	44.	60
β. — — —	—	—	Helix	—	—
rapunculoides	—	—	<i>Illecebrum</i>	44.	60
Trachaelium	—	—	verticillatum	—	—
glomerata	—	—	<i>Glaux</i>	45.	60
Cervicaria	58	—	maritima	—	—
hederacea	—	—	<i>Thefium</i>	45.	61
<i>Phyteuma</i>	43.	58	alpinum	—	—
ipicata	—	—	linophyllum	—	—
<i>Samolus</i>	43.	58	2. DIGYNIA.		
Valerandi	—	—	<i>Asclepias</i>	45.	61
<i>Lonicera</i>	43.	58	Vincedoxicum	—	—
Periclymenum	—	—	<i>Herniaria</i>	45.	61
Xylosteum	—	—	glabra	—	—
coerulea	—	—	<i>Chenopodium</i>	45.	61
<i>Verbascum</i>	43.	58	Bonus - Henricus	—	—
Thapsus	—	—	urbicum	—	—
Lychnitis	—	—	rubrum	—	—
nigrum	—	—	murale	—	—
<i>Datura</i>	43.	59	viride	—	62
Stramonium	—	—	β. album	—	—
<i>Hyoſcyamus</i>	43.	59	hybridum	—	—
niger	—	—	ambrosioides	—	—
<i>Atropa</i>	43.	59	glaucum	—	—
Belladonna	—	—	Vulvaria	—	—
<i>Physalis</i>	44.	59	polyſpermum	—	—
Alkekengi	—	—	maritimum	—	—
<i>Solanum</i>	44.	59	<i>Salsola</i>	45.	62
nigrum	—	—	Kali	—	—
Dulcamara	—	—	α. decumbens	—	—
<i>Rhamnus</i>	44.	59	β. erecta hisp.	—	—
catharticus	—	—	γ. erecta glaberr.	—	—
Frangula	—	—	hirsuta	—	63
<i>Evonymus</i>	44.	59	<i>Ulmus</i>	46.	63
europaeus	—	—	campeſtris	—	—

Sivernia

PLANTARUM.

345

	Pag. 46. 63	<i>Heracleum</i>	Pag. 48. 66
<i>Sivertia</i>			
fulcata	—	Sphondilium	—
perennis	—	angustifolium	—
<i>Gentiana</i>	46. 63	<i>Ligusticum</i>	48. 66
purpurea	—	Levisticum	—
<i>Pneumonanthe</i>	—	scoticum	—
nivalis	—	<i>Angelica</i>	48. 67
pulchella	64	Archangelica	—
<i>Centaurium</i>	—	sylvestris	—
involucrata	—	<i>Sium</i>	48. 67
Amarella	—	latifolium	—
campestris	—	angustifolium	—
tenella	—	nodiflorum	—
ciliata	—	Falcaria	—
β. detonsa		<i>Sison</i>	49. 67
γ. ferrata		inundatum	—
filiformis	—	<i>Oenanthe</i>	49. 67
<i>Eryngium</i>	46. 65	fistulosa	—
maritimum	—	crocata	—
campestre	—	pimpinelloides	—
<i>Hydrocotyle</i>	46. 65	<i>Phellandrium</i>	49. 68
vulgaris	—	aquaticum	—
<i>Sanicula</i>	46. 65	<i>Aethusa</i>	49. 68
europaea	—	Cynapium	—
<i>Bupleurum</i>	46. 65	<i>Cicuta</i>	49. 68
falcatum	—	virosa	—
tenuissimum	—	<i>Scandix</i>	49. 68
<i>Tordylium</i>	47. 65	odorata	—
Anthriscus	—	Pecten	—
<i>Daucus</i>	47. 65	Cerfolium	—
Carota	—	Anthriscus	—
<i>Bunium</i>	47. 65	infesta	—
Bulbocastanum	—	<i>Chaerophyllum</i>	50. 68
<i>Conium</i>	47. 66	sylvestre	—
maculatum	—	temulum	—
<i>Selinum</i>	47. 66	bulbosum	—
palustre	—	<i>Imperatoria</i>	50. 68
Carvifolia	—	Ostruthium	—
<i>Athamanta</i>	48. 66	<i>Seseli</i>	49. 69
Libanotis	—	montanum	—
Oreoselinum	—	<i>Pastinaca</i>	50. 69
<i>Laserpitium</i>	48. 66	fativa	—
latifolium	—		
prutenicum	—		

<i>Ceram</i>	Pag. 50. 69	<i>Sibbaldia</i>	Pag. 52. 72
Carvi	—	procumbens	—
<i>Pimpinella</i>	50. 69	6. POLYGYNIA.	
<i>Saxifraga</i>	—	<i>Myosurus</i>	52. 72
β. magna	—	minimus	—
γ.	—	VI. HEXANDRIA.	
α.	—	I. MONOGYNIA.	
diffecta	—	<i>Allium</i>	73. 76
<i>Apium</i>	50. 69	<i>Scorodoprasum</i>	—
graveolens	—	arenarium	—
<i>Aegopodium</i>	50. 70	carinatum	—
<i>Podagraria</i>	—	oleraceum	—
3. TRIGYNIA.			
<i>Viburnum</i>	51. 70	ursinum	—
<i>Opulus</i>	—	<i>Schoenoprasum</i>	—
<i>Sambucus</i>	51. 70	<i>Lilium</i>	73. 76
nigra	—	bulbiferum	—
<i>Ebulus</i>	—	Martagon	—
<i>Tamarix</i>	51. 70	<i>Fritillaria</i>	73. 76
germanica	—	Meleagris	—
<i>Corrigiola</i>	51. 70	<i>Tulipa</i>	73. 77
littoralis	—	sylvestris	—
<i>Alfina</i>	51. 70	<i>Ornithogalum</i>	73. 77
media	—	luteum	—
α. pentandra	—	minimum	—
β. decandra	—	umbellatum	—
<i>viscosa</i>	—	nutans	—
4. TETRAGYNIA.			
<i>Parnassia</i>	51. 71	<i>Scilla</i>	73. 77
palustris	—	italica	—
5. PENTAGYNIA.			
<i>Statice</i>	51. 71	bifolia	—
<i>Armeria</i>	—	<i>Anthericum</i>	74. 77
<i>Limonium</i>	—	ramosum	—
<i>Linum</i>	52. 71	<i>Liliago</i>	—
catharticum	—	officinarum	78
<i>Radiola</i>	—	calyculatum	—
<i>Drosera</i>	52. 71	<i>Asparagus</i>	74. 78
rotundifolia	—	officinalis	—
longifolia	—	<i>Convallaria</i>	74. 78
<i>Crassula</i>	52. 71	majalis	—
rubens	—	verticillata	—
		<i>Polygonatum</i>	—
		multiflorum	—
		bifolia	—
		<i>Hyacin-</i>	—

<i>Hyacinthus</i>	Pag. 73. 79	<i>Berberis</i>	Pag. 74. 82
botryoides	—	vulgaris	—
<i>Acorus</i>	74. 79	<i>Peplis</i>	74. 82
Calamus	—	Portula	—
<i>Juncus</i>	74. 79	3. TRIGYNIA.	
* <i>culmis nudis</i> ,	—	<i>Rumex</i>	75. 83
auctus	—	crispus	—
conglomeratus	—	Nemolapathum	—
effusus	—	maritimus	—
α. - - - -	—	acutus	—
β. - - - -	—	obtusifolius	—
γ. - - - -	—	aquaticus	—
inflexus	—	digynus	—
filiformis	—	Acetosa	—
trifidus	80	α. vulgaris	—
squarrosus	—	β. alpina	—
** <i>culmis capitatis</i> ,	—	Acetofella	—
capitatus	—	<i>Scheuchzeria</i>	75. 84
α. - - - -	—	palustris	—
β. - - - -	—	<i>Triglochin</i>	75. 84
setifolius	—	palustre	—
articulatus	—	maritimum	—
α. aquaticus	—	4. POLYGYNIA.	
β. - - - -	—	<i>Alisma</i>	75. 84
bulbosus	—	Plantago aquatica	—
bufonius	81	β. - - - -	—
β. viviparus	—	natans	—
γ. pumilus	—	ranunculoides	—
<i>Tenageja</i>	—	VII. HEPTANDRIA.	
lupinus	—	I. MONOGYNIA.	
stygius	—	<i>Trientalis</i>	85. 86
biglumis	—	europaea	—
triglumis	—	VIII. OCTANDRIA.	
vernalis	—	I. MONOGYNIA.	
parviflorus	—	<i>Oenothera</i>	87. 89
maximus	82	biennis	—
multiflorus	—	<i>Epilobium</i>	87. 89
campestris	—	angustifolium	—
β. lapponicus	—	Halleri	—
γ. tenuis	—	frigidum	—
δ. strictus	—	hirsutum	—
spicatus	—	parvi-	
subverticillatus	—		

<i>parviflorum</i>	Pag. 89	3. TETRAGYNIA.	
<i>obscurum</i>	—	<i>Paris</i>	Pag. 88. 92
<i>montanum</i>	90	<i>quatrifolia</i>	—
<i>roseum</i>	—	<i>Adoxa</i>	88. 93
<i>tetragonum</i>	—	<i>Moschatellina</i>	—
<i>palustre</i>	—	<i>Elatine</i>	88. 93
<i>β. alpinum</i>	—	<i>Hydropiper</i>	—
<i>alpinum</i>	—	<i>Alfinastrum</i>	—
<i>Vaccinium</i>	87. 90		
<i>Myrtillus</i>	—	IX. ENNEANDRIA.	
<i>uliginosum</i>	—	I. TRIGYNIA.	
<i>Vitis idaea</i>	—	<i>Rheum</i>	94. 95
<i>Oxycoccus</i>	—	<i>Rhaponticum</i>	—
<i>Erica</i>	87. 91	<i>Butomus</i>	94. 95
<i>vulgaris</i>	—	<i>umbellatus</i>	—
<i>Tetralix</i>	—		
<i>cinerea</i>	—	X. DECANDRIA.	
<i>Daphne</i>	87. 91	I. MONOGYNIA.	
<i>Mezereum</i>	—	<i>Monotropa</i>	96. 100
3. TRIGYNIA.		<i>Hypopithys</i>	—
<i>Polygonum</i>	87. 91	<i>Ledum</i>	96. 100
<i>Bistorta</i>	—	<i>palustre</i>	—
<i>viviparum</i>	—	<i>groenlandicum</i>	—
<i>amphibium</i>	—	<i>Andromeda</i>	96. 100
<i>α. terrestre</i>	—	<i>tetragona</i>	—
<i>β. aquaticum</i>	—	<i>hypnoides</i>	—
<i>Hydropiper</i>	92	<i>coerulea</i>	—
<i>Perficaria</i>	—	<i>polifolia</i>	—
<i>α. latifolia</i>	—	<i>calyculata</i>	101
<i>β. angustifolia</i>	—	<i>Arbutus</i>	96. 101
<i>γ. maculata</i>	—	<i>alpina</i>	—
<i>δ. pusilla</i>	—	<i>Uva Ursi</i>	—
<i>ε. tomentosa</i>	—	<i>Pyrola</i>	96. 101
<i>aviculare</i>	—	<i>rotundifolia</i>	—
<i>α. latifolium</i>	—	<i>minor</i>	—
<i>β. angustifolium</i>	—	<i>secunda</i>	—
<i>γ. brevifolium</i>	—	<i>umbellata</i>	—
<i>δ. purpurascens</i>	—	<i>uniflora</i>	—
<i>ε. aphyllum</i>	—	2. DIGYNIA.	
<i>Fagopyrum</i>	—	<i>Chrysosplenium</i>	96. 101
<i>Convolvulus</i>	—	<i>alternifolium</i>	—
<i>dumetorum</i>	—	<i>oppositifolium</i>	—

<i>Saxifraga</i>	Pag. 97. 102	<i>Silene</i>	Pag. 97. 105
<i>Cotyledon</i>	—	<i>nutans</i>	—
<i>stellaris</i>	—	<i>noctiflora</i>	—
<i>β. comosa</i>	—	<i>Armeria</i>	—
<i>nivalis</i>	—	<i>rupestris</i>	106
<i>cuneifolia</i>	—	<i>acaulis</i>	—
<i>oppositifolia</i>	—	<i>β. exscapa</i>	—
<i>β. indurata</i>	—	<i>Stellaria</i>	98. 106
<i>Hirculus</i>	—	<i>nemorum</i>	—
<i>Aizoides</i>	—	<i>dichotoma</i>	—
<i>granulata</i>	103	<i>Holosteia</i>	—
<i>bulbifera</i>	—	<i>graminea</i>	—
<i>cernua</i>	—	<i>α. major</i>	—
<i>β. biflora</i>	—	<i>β. minor</i>	—
<i>rivularis</i>	—	<i>uliginosa</i>	—
<i>tridactylites</i>	—	<i>palustris</i>	—
<i>β. integrifolia</i>	—	<i>cerastoides</i>	107
<i>petraea</i>	—	<i>humifusa</i>	—
<i>caespitosa</i>	—	<i>biflora</i>	—
<i>β. groenlandica</i>	—	<i>groenlandica</i>	—
<i>tricuspidata</i>	104	<i>Arenaria</i>	98. 107
<i>hypnoides</i>	—	<i>peploides</i>	—
<i>Scleranthus</i>	97. 104	<i>trinervia</i>	—
<i>annuus</i>	—	<i>ciliata</i>	—
<i>perennis</i>	—	<i>β. norvegica</i>	—
<i>Gypsophila</i>	97. 104	<i>multicaulis</i>	—
<i>fastigiata</i>	—	<i>serpyllifolia</i>	108
<i>muralis</i>	—	<i>rubra</i>	—
<i>Saponaria</i>	97. 104	<i>α. campestris</i>	—
<i>officinalis</i>	—	<i>β. maritima</i>	—
<i>Dianthus</i>	97. 104	4. PENTAGYNIA.	—
<i>Carthusianorum</i>	—	<i>Sedum</i>	98. 108
<i>Armeria</i>	—	<i>Telephium</i>	—
<i>prolifer</i>	—	<i>β. purpureum</i>	—
<i>deltoides</i>	105	<i>reflexum</i>	—
<i>fuperbus</i>	—	<i>rupestre</i>	—
<i>arenarius</i>	—	<i>Oederi</i>	—
3. TRIGYNIA.	—	<i>album</i>	—
<i>Cucubalus</i>	97. 105	<i>acre</i>	—
<i>Behen</i>	—	<i>fexangulare</i>	109
<i>β. repens</i>	—	<i>annuum</i>	—
<i>viscosus</i>	—	<i>anglicum</i>	—
<i>Otites</i>	—	<i>villosum</i>	—

Oxalis

<i>Oxalis</i>	Pag. 98. 109	<i>Euphorbia</i>	Pag. 112. 113
Acetosella	—	Peplus	—
corniculata	—	exigua	—
<i>Agrostemma</i>	98. 109	Helioscopia	114
Githago	—	Efula	—
<i>Lachnis</i>	98. 109	palustris	—
Flos Cuculi	—	amygdaloides	—
Viscaria	—	4. DODECAGYNIA.	
alpina	—	<i>Sempervivum</i>	112. 114
dioica	110	testorum	—
α . purpurea	—	XII. ICOSANDRIA.	
β . alba	—	I. MONOGYNIA.	
γ . hermaphrodita	—	<i>Prunus</i>	115. 118
apetala	—	Padus	—
<i>Cerastium</i>	99. 110	Cerasus	—
vulgatum	—	avium	—
viscosum	—	domestica	—
semidecandrum	—	infirmitia	—
alpinum	—	spinosa	—
α . hirsutum	—	2. DIGYNIA.	
β . glabrum	—	<i>Crataegus</i>	115. 118
aquaticum	—	Aria	—
latifolium	—	α . ulmifolia	—
<i>Spergula</i>	99. 110	β . sinuata	—
arvensis	—	torminalis	119
nodosa	—	Oxyacantha	—
sagmoides	111	monogyna	—
subulata	—	3. TRIGYNIA.	
XI. DODECANDRIA.		<i>Sorbus</i>	115. 119
I. MONOGYNIA.		aucuparia	—
<i>Asarum</i>	112. 113	hybrida	—
europaeum	—	4. PENTAGYNIA.	
<i>Lythrum</i>	112. 113	<i>Mespilus</i>	115. 119
Salicaria	—	Cotoneaster	—
β . trifolia	—	<i>Pyrus</i>	115. 119
γ . lapponica	—	communis	—
2. DIGYNIA.		malus	—
<i>Agrimonia</i>	112. 113	<i>Spiraea</i>	116. 119
Eupatoria	—	falicifolia	—
3. TRIGYNIA.		Filipendula	120
<i>Reseda</i>	112. 113	Ulmaria	—
Luteola	—	5. POLY-	

5. POLYGYNIA.

<i>Rosa</i>	Pag. 116. 120
* <i>germinibus subglobosis.</i>	—
<i>Eglanteria</i>	—
<i>arvensis</i>	—
<i>cinnamomea</i>	—
<i>fluvialis</i>	—
<i>spinofillima</i>	—
<i>villosa</i>	—
** <i>germinibus oblongis.</i>	—
<i>rubiginosa</i>	121
<i>canina</i>	—
<i>Rubus</i>	116. 121
<i>idaeus</i>	—
<i>caesius</i>	—
<i>fruticosus</i>	—
<i>saxatilis</i>	—
<i>arcticus</i>	—
<i>Chamaemorus</i>	—
<i>Fragaria</i>	116. 121.
<i>vesca</i>	—
<i>Potentilla</i>	116. 122
<i>fruticosa</i>	—
<i>Anserina</i>	—
<i>rupestris</i>	—
<i>supina</i>	—
<i>argentea</i>	—
<i>opaca</i>	—
<i>verna</i>	—
<i>aurea</i>	—
<i>reptans</i>	—
<i>norvegica</i>	—
<i>nivea</i>	123
β.	—
γ.	—
<i>retusa</i>	—
<i>Tormentilla</i>	116. 123
<i>erecta</i>	—
<i>Geum</i>	116. 123
<i>urbanum</i>	—
<i>rivale</i>	—
β.	—

<i>Dryas</i>	Pag. 117. 123
<i>octopetala</i>	—
<i>Comarum</i>	117. 123
<i>palustre</i>	—

XIII. POLYANDRIA.

I. MONOGYNIA.

<i>Achaea</i>	124. 127
<i>spicata</i>	—
<i>Chelidonium</i>	124. 127
<i>majus</i>	—
<i>tenuifolium.</i>	—
<i>Glaucium</i>	—
<i>Papaver</i>	124. 127
<i>Argemone</i>	—
<i>nudicaule</i>	—
<i>Rhoeas</i>	—
<i>dubium</i>	—
<i>Nymphaea</i>	124. 127
<i>lutea</i>	—
<i>alba</i>	—
<i>Tilia</i>	124. 128
<i>europaea</i>	—
α. <i>grandifolia</i>	—
β. <i>parvifolia</i>	—
γ. <i>corallina</i>	—
<i>Cistus</i>	124. 128
<i>oelandicus</i>	—
<i>Fumana</i>	—
<i>Helianthemum</i>	—

2. TRIGYNIA.

<i>Delphinium</i>	125. 128
<i>Consolida</i>	—
<i>Aconitum</i>	125. 128
<i>Lycostonum</i>	—
α. <i>coeruleum</i>	—
<i>Napellus</i>	—

3. PENTAGYNIA.

<i>Aquilegia</i>	125. 129
<i>vulgaris</i>	—

4. POLY-

4. POLYGYNIA.

Anemone Pag. 125. 129*Hepatica* —*patens* —*vernalis* —*Pulsatilla* —*pratensis* —*sylvestris* —*nemorosa* —*ranunculoides* —*Thalictrum* 125. 129*alpinum* —*minus* 130*flavum* —*simplex* —*aquilegifolium* —*Adonis* 125. 130*aestivalis* —*vernalis* —*Ranunculus* 126. 130*Flammula* —*reptans* —*Lingua* —*Ficaria* —*auricomus* 131*sceleratus* —*plataniifolius* —*illyricus* —*glacialis* —*nivalis* —*β. pygmaeus**lapponicus* —*hypertoreus* —*bulbosus* 132*Philonotis* —*polyanthemus* —*acris* —*lanuginosus* —*arvensis* —*hederaceus* —

aquatilis

Pag. 133

*α. anomalis**β. heterophyllus**γ. millefolius**δ. peucedanifolius**Trollius* 126. 133*europaeus* —*Helleborus* 126. 133*trifolius* —*Calccha* 126. 133*palustris* —

XIV. DIDYNAMIA.

I. GYMNOSPERMIA.

Ajuga 134. 139*pyramidalis* —*alpina* —*reptans* —*Teucrium* 134. 139*Chamaepithys* —*Scordium* —*Scorodonia* —*Nepeta* 134. 139*Cataria* —*Mentha* 134. 139*sylvestris* —*viridis* —*hirsuta* 140*aquatica* —*farva* —*gentilis* —*arvensis* —*Glecoma* 134. 140*hederacea* —*Lamium* 134. 140*album* —*purpureum* —*amplexicaule* —

Galeopsis Pag. 134. 141

Ladanum —

α. angustifolia —

β. latifolia —

Tetrahit —

α. vulgaris —

β. cannabina —

Galeobdolon —

Betonica 135. 141

officinalis —

Stachys 135. 141.

sylvatica —

palustris —

alpina —

recta —

arvensis 142

Ballota 135. 142

nigra —

alba —

Marrubium 135. 142

vulgare —

Leonurus 135. 142

Cardiaca —

Clinopodium 135. 142

vulgare —

Origanum 135. 142

vulgare —

Thymus 136. 142

Serpillum —

α. rubrum —

β. album —

γ. lanuginosum —

Acinos 143

Dracocephalum 136. 143

Ruischiana —

thymiflorum —

Scutellaria 136. 143

gallericulata —

hastifolia —

Prunella 136. 143

vulgaris —

β. grandiflora —

I. ANGIOSPERMIA.

Barbisa Pag. 136. 143

alpina —

Rhinanthus 136. 144

Crista Galli —

α. foemina —

β. mas —

Euphrasia 136. 144

officinalis —

Odontites —

Melampyrum 137. 144

cristatum —

arvense —

nemorosum —

pratense —

sylvaticum —

Lathraea 137. 144

Squamaria —

Pedicularis 137. 145

palustris —

sylvatica —

Sceptrum Carolinum —

flammea —

hirsuta —

lapponica —

groenlandica —

foliosa 146

Amirrhinum 137. 146

Elatine —

spurium —

minus —

Linaria —

β. multiplicata —

Orontium —

Scrophularia 137. 146

nodosa —

aquatica —

vernalis —

Digitalis 137. 147

purpurea —

Linnaea 137. 147

borealis —

Z

Limo.

<i>Limosella</i>	Pag. 138. 147	<i>Alyssum</i>	Pag. 149. 154
<i>aquatica</i>	—	<i>incanum</i>	—
<i>Orobanchë</i>	138. 147	<i>calycinum</i>	—
<i>major</i>	—	<i>Lunaria</i>	149. 154
		<i>rediviva</i>	—
		<i>annua</i>	155

XV. TETRADYNAMIA.

I. SILICULOSA.

<i>Myagrum</i>	148. 152
<i>lativum</i>	—
<i>paniculatum</i>	—
<i>Subularia</i>	148. 152
<i>aquatica</i>	—
<i>Draba</i>	148. 152
<i>alpina</i>	—
<i>verna</i>	—
<i>muralis</i>	—
<i>nemoralis</i>	—
<i>hirta</i>	—
<i>β. norvegica</i>	—
<i>incana</i>	153
<i>contorta</i>	—
<i>Lepidium</i>	148. 153
<i>alpinum</i>	—
<i>petraeum</i>	—
<i>latifolium</i>	—
<i>ruderales</i>	—
<i>Thlaspi</i>	148. 153
<i>arvense</i>	—
<i>campestre</i>	—
<i>Bursa Pastoris</i>	—
<i>α. sinuata</i>	—
<i>β. integrifolia</i>	—
<i>Cochlearia</i>	148. 154
<i>officinalis</i>	—
<i>danica</i>	—
<i>anglica</i>	—
<i>groenlandica</i>	—
<i>Coronopus</i>	—
<i>Armoracia</i>	—
<i>Iberis</i>	148. 154
<i>nudicaulis</i>	—
<i>β. integrifolia</i>	—

2. SILIQUOSA.

<i>Dentaria</i>	149. 155
<i>bulbifera</i>	—
<i>Cardamine</i>	149. 155
<i>bellidifolia</i>	—
<i>petraea</i>	—
<i>α. hirsuta</i>	—
<i>β. glabra</i>	—
<i>trifolia</i>	—
<i>Impatiens</i>	—
<i>parviflora</i>	—
<i>hirsuta</i>	—
<i>pratensis</i>	—
<i>amara</i>	156
<i>α. glabra</i>	—
<i>β. hirsuta</i>	—
<i>Sisymbrium</i>	149. 156
<i>Nasturtium</i>	—
<i>sylvestre</i>	—
<i>β. islandicum</i>	—
<i>amphibium</i>	—
<i>α. palustre</i>	—
<i>β. aquaticum</i>	—
<i>γ. terrestre</i>	—
<i>supinum</i>	—
<i>arenosum</i>	—
<i>Sophia</i>	157
<i>Iris</i>	—
<i>Erysimum</i>	149. 157
<i>officinale</i>	—
<i>Barbarea</i>	—
<i>Alliaria</i>	—
<i>Cheiranthoides</i>	—
<i>hieracifolium</i>	—
<i>Cheiranthus</i>	149. 157
<i>Erysimoides</i>	—

PLANTARUM.

355

<i>Hesperis</i>	Pag. 150. 157
inodora	—
<i>Arabis</i>	150. 157
alpina	—
Thaliana	158
β.	—
<i>Turritis</i>	150. 158
glabra	—
hirsuta	—
alpina	—
<i>Brassica</i>	150. 158
campestris	—
Napus	—
<i>Sinapis</i>	150. 158
arvensis	—
nigra	—
<i>Raphanus</i>	150. 158
Raphanistrum	—
<i>Bunias</i>	150. 158
Cakile	—
orientalis	—
<i>Isatis</i>	151. 159
tinctoria	—
<i>Crambe</i>	151. 159
maritima	—
XVI. MONADELPHIA.	
I. DECANDRIA.	
<i>Geranium</i>	160. 161
cicutarium	—
phaeum	—
fuscum	—
sylvaticum	—
palustre	—
pratense	—
robertianum	—
lucidum	—
molle	162
Columbinum	—
dissectum	—
rotundifolium	—
sanguineum	—

2. POLYANDRIA.	
<i>Althaea</i>	Pag. 160. 162
officinalis	—
<i>Malva</i>	160. 162
rotundifolia	—
sylvestris	—
Alcea	163
moschata	—
<i>Lavatera</i>	160. 163
thuringiaca	—

XVII. DIADELPHIA.

I. HEXANDRIA.	
<i>Fumaria</i>	164. 167
bulbosa	—
fabacea	—
cava	—
officinalis	—
capreolata	—

2. OCTANDRIA.	
<i>Polygala</i>	164. 167
amara	—
vulgaris	168
α. coerulea	—
β. carnea	—
γ. alba	—
major	—

3. DECANDRIA.	
<i>Spartium</i>	164. 168
scoparium	—
<i>Genista</i>	164. 168
tinctoria	—
pilosa	—
anglica	—
germanica	169
<i>Ulex</i>	164. 169
europaeus	—
<i>Ononis</i>	165. 169
arvensis	—
spinosa	—

<i>Anshyllis</i>	Pag. 165. 169	<i>Hedysarum</i>	Pag. 166. 172
<i>Vulneraria</i>	—	<i>Onobrychis</i>	—
<i>α. lutea</i>	—	<i>Phaca</i>	166. 172
<i>β. alba</i>	—	<i>alpina</i>	—
<i>γ. rubra</i>	—	<i>Astragalus</i>	166. 172
<i>δ. coccinea</i>	—	<i>pilosus</i>	—
<i>Pisum</i>	165. 169	<i>glycyphyllos</i>	—
<i>arvense</i>	—	<i>arenarius</i>	173
<i>maritimum</i>	—	<i>danicus</i>	—
<i>Orobus</i>	165. 170	<i>alpinus</i>	—
<i>vernus</i>	—	<i>Uralensis</i>	—
<i>tuberosus</i>	—	<i>campestris</i>	—
<i>β. grandis</i>	—	<i>Trifolium</i>	166. 173
<i>niger</i>	—	<i>M. officinalis</i>	—
<i>Lathyrus</i>	165. 170	<i>M. Ornithopodioides</i>	—
<i>tuberosus</i>	—	<i>hybridum</i>	—
<i>pratensis</i>	—	<i>repens</i>	—
<i>sylvestris</i>	—	<i>pratense</i>	—
<i>β. platyphyllos</i>	—	<i>flexuosum</i>	174
<i>latifolius</i>	—	<i>alpestre</i>	—
<i>heterophyllus</i>	—	<i>arvense</i>	—
<i>palustris</i>	—	<i>β. pumilum</i>	—
<i>Vicia</i>	165. 171	<i>striatum</i>	—
<i>pisiformis</i>	—	<i>fragiferum</i>	—
<i>dumetorum</i>	—	<i>montanum</i>	—
<i>sylvatica</i>	—	<i>agrarium</i>	—
<i>castubica</i>	—	<i>spadiceum</i>	—
<i>Cracca</i>	—	<i>procumbens</i>	—
<i>fativa</i>	—	<i>filiforme</i>	—
<i>Lathyroides</i>	—	<i>Lotus</i>	166. 175
<i>sepium</i>	—	<i>maritima</i>	—
<i>Ervum</i>	165. 171	<i>corniculata</i>	—
<i>Lens</i>	—	<i>β. procerior</i>	—
<i>tetraspermum</i>	—	<i>γ. tenuifolia</i>	—
<i>hirsutum</i>	172	<i>Medicago</i>	166. 175
<i>Coronilla</i>	166. 172	<i>falcata</i>	—
<i>Emerus</i>	—	<i>Lupulina</i>	—
<i>varia</i>	—	<i>polymorpha</i>	—
<i>Ornithopus</i>	166. 172	<i>α. minima.</i>	—
<i>perpusillus</i>	—		

XVIII. POLYADELPHIA.

Leontodon

Pag. 179. 187.

I. POLYANDRIA.

*Taraxacum**Hypericum* Pag. 176. 177 β . palustre

quadrangulare —

perforatum —

humifusum —

montanum —

hirsutum —

pulchrum —

Hieracium 179. 187

alpinum —

 α . grandiflorum β . parviflorum*Taraxaci* —*Pilosella* —

dubium —

Auricula 188

cymosum —

praemorsum —

aurantiacum —

murorum —

 β . sylvaticum γ . laciniatum δ . myophorum

paludosum —

Sabaudum —

umbellatum —

XIX. SYNGENESIA.

I. POLYG. AEQUAL.

Tragopogon 178. 185

pratense —

porrifolium —

Scorzonera 178. 185

humilis —

purpurea —

Picris 178. 185*Echioides* —*Hieracioides* —*Apargia* 179. 185

autumnalis —

hispida 186

Sonchus 178. 186

maritimus —

palustris —

arvensis —

oleraceus —

 α . laevis β . asper

lapponicus —

alpinus —

sibiricus —

Lactuca 178. 186

quercina —

virosa —

Chondrilla 178. 187

juncea —

Prenanthes 178. 187

muralis —

Crepis 179. 188

testorum —

biennis 189

Hyoferis 179. 189

minima —

Hypochaeris 179. 189

maculata —

glabra —

radicata —

Lapsana 179. 189

communis —

Cichorium 179. 189

Intybus —

Artium 180. 189*Lappa* — α . vulgaris β . arachnoidea

<i>Serratula</i>	Pag. 180. 189	<i>rupestris</i>	Pag. 192
<i>tinctoria</i>	—	<i>Abinthium</i>	—
<i>alpina</i>	190	<i>vulgaris</i>	193
<i>arvensis</i>	—	<i>α. alba</i>	—
<i>α. glabra</i>	—	<i>β. rubra</i>	—
<i>β. tomentosa</i>	—	<i>Gnaphalium</i>	181. 193
<i>Carduus</i>	180. 190	<i>arenarium</i>	—
<i>nutans</i>	—	<i>dioicum</i>	—
<i>Acanthoides</i>	—	<i>α. mas</i>	—
<i>crispus</i>	—	<i>β. foemina</i>	—
<i>Cnicus</i>	180. 190	<i>alpinum</i>	—
<i>acaulis</i>	—	<i>sylvaticum</i>	—
<i>lanceolatus</i>	—	<i>norvegicum</i>	—
<i>palustris</i>	—	<i>uliginosum</i>	—
<i>heterophyllus</i>	191	<i>Conyza</i>	181. 194
<i>helenioides</i>	—	<i>squarrosa</i>	—
<i>oleraceus</i>	—	<i>Erigeron</i>	181. 194
<i>Onopordum</i>	180. 191	<i>acre</i>	—
<i>Acanthium</i>	—	<i>dröbachiense</i>	—
<i>Carlina</i>	180. 191	<i>uniflorum</i>	—
<i>vulgaris</i>	—	<i>Tussilago</i>	182. 194
<i>Bidens</i>	180. 191	<i>Farfara</i>	—
<i>tripartita</i>	—	<i>frigida</i>	—
<i>β. pumila</i>	—	<i>alba</i>	—
<i>minima</i>	—	<i>spuria</i>	—
<i>cernua</i>	—	<i>Petasites</i>	—
<i>β. radiata</i>	—	<i>Senecio</i>	182. 195
<i>frondosa</i>	192	<i>vulgaris</i>	—
<i>Eupatorium</i>	180. 192	<i>viscosus</i>	—
<i>cannabinum</i>	—	<i>β. denticulatus</i>	—
<i>Chrysocoma</i>	181. 192	<i>sylvaticus</i>	—
<i>Lynosyris</i>	—	<i>Jacobaea</i>	—
		<i>β.</i>	—
		<i>aquaticus</i>	—
		<i>paludosus</i>	—
2. POLYGAMIA SU-		<i>Aster</i>	182. 195
PERFLUA.		<i>Tripolium</i>	—
<i>Thunacetum</i>	181. 192	<i>annuus</i>	—
<i>vulgare</i>	—	<i>Solidago</i>	182. 196
<i>Artemisia</i>	181. 192	<i>Virgaurea</i>	—
<i>campestris</i>	—	<i>Cineraria</i>	182. 196
<i>α. alba</i>	—	<i>palustris</i>	—
<i>β. rubra</i>	—	<i>campestris</i>	—
<i>maritima</i>	—		
			<i>Inula</i>

<i>Inula</i>	Pag. 182. 196	4. POLYGAMIA NECESSAR.	
<i>Helenium</i>	—	<i>Calendula</i>	Pag. 184. 199
<i>britannica</i>	—	<i>arvensis</i>	—
<i>dysenterica</i>	—	<i>Filago</i>	184. 200
<i>Pulicaria</i>	—	<i>germanica</i>	—
<i>falicina</i>	197	<i>montana</i>	—
<i>hirta</i>	—	<i>arvensis</i>	—
<i>Arnica</i>	182. 197		
<i>montana</i>	—	5. MONOGAMIA.	
<i>α. pratensis</i>	—	<i>Fasione</i>	184. 200
<i>β. alpina</i>	—	<i>montana</i>	—
<i>Bellis</i>	182. 197	<i>Lobelia</i>	184. 200
<i>perennis</i>	—	<i>Dortmanna</i>	—
<i>Chrysanthemum</i>	183. 197	<i>Viola</i>	184. 200
<i>segetum</i>	—	<i>hirta</i>	—
<i>Leucanthemum</i>	—	<i>palustris</i>	—
<i>inodorum</i>	—	<i>odorata</i>	—
<i>Matricaria</i>	183. 197	<i>canina</i>	—
<i>Parthenium</i>	—	<i>montana</i>	—
<i>maritima</i>	198	<i>mirabilis</i>	201
<i>suaveolens</i>	—	<i>biflora</i>	—
<i>Chamomilla</i>	—	<i>tricolor</i>	—
<i>Cotula</i>	183. 198	<i>β. arvensis</i>	—
<i>coronopifolia</i>	—	<i>Impatiens</i>	184. 201
<i>Anthemis</i>	183. 198	<i>noli tangere</i>	—
<i>maritima</i>	—		
<i>arvensis</i>	—	XX. GYNANDRIA.	
<i>Cotula</i>	—	I. DIANDRIA.	
<i>tinctoria</i>	—	<i>Orchis</i>	202. 204
<i>Achillea</i>	183. 198	* <i>Testiculatae.</i>	
<i>Ptarmica</i>	—	<i>bifolia</i>	—
<i>Millefolium</i>	—	<i>pyramidalis</i>	—
<i>nobilis</i>	199	<i>Merio</i>	—
		<i>mascula</i>	—
3. POLYG. FRUSTAN.		<i>ustulata</i>	—
<i>Coreopsis</i>	183. 199	<i>militaris</i>	—
<i>Bidens</i>	—	<i>pallens</i>	—
<i>Centaurea</i>	183. 199	** <i>Palmarae.</i>	
<i>phrygia</i>	—	<i>latifolia</i>	205
<i>nigra</i>	—	<i>mixta</i>	—
<i>Cyanus</i>	—	<i>α. incarnata</i>	—
<i>Scabiosa</i>	—	<i>β. sambucina</i>	—
<i>Jacea</i>	—		

**** Spicis plurim. androgyn.**

<i>incurva</i>	Pag. 217
<i>chordorhiza</i>	—
<i>arenaria</i>	—
<i>leporina</i>	—
<i>disticha</i>	—
<i>vulpina</i>	218
<i>β. intermedia</i>	—

<i>echinata</i>	—
<i>loliacea</i>	—
<i>Heleonastes</i>	—
<i>microstachia</i>	—
<i>remota</i>	—
<i>elongata</i>	219
<i>canescens</i>	—
<i>tenella</i>	—
<i>multiculmis</i>	—
<i>paniculata</i>	—
<i>β. minor</i>	—
<i>norvegica</i>	—

***** Spic. sexu distinctis:
foem. sessil.**

<i>flava</i>	220
<i>pedata</i>	—
<i>digitata</i>	—
<i>montana</i>	—
<i>globularis</i>	—
<i>filiformis</i>	—
<i>pilulifera</i>	—
<i>laxatilis</i>	221
<i>humilis</i>	—

****** Spic. sexu distinctis:
foem. pedunc.**

<i>atrata</i>	—
<i>limosa</i>	—
<i>β. pauciflora</i>	—
<i>capillaris</i>	—
<i>pallefcens</i>	—
<i>panicea</i>	—
<i>Pseudocyperus</i>	—
<i>sylvatica</i>	222
<i>distans</i>	—
<i>stolonifera</i>	—
<i>caespitosa</i>	—

******* Spic. masc. pluribus.**

<i>maritima</i>	Pag. 222
<i>flacca</i>	—
<i>lasiocarpa</i>	—
<i>acuta</i>	223
<i>α. nigra</i>	—
<i>β. rufa</i>	—
<i>γ. gracilis</i>	—
<i>crassa</i>	—
<i>vesicaria</i>	—
<i>obrusangula</i>	—
<i>hirta</i>	—
<i>Axyris</i>	211. 223
<i>prostrata</i>	—

4. TETRANDRIA.

<i>Littorella</i>	211. 224
<i>lacustris</i>	—
<i>Betula</i>	211. 224
<i>alba</i>	—
<i>α. vulgaris</i>	—
<i>β. pendula</i>	—
<i>γ. laciniata</i>	—

<i>nana</i>	—
<i>Alnus</i>	—
<i>β. pumila</i>	—
<i>incana</i>	—
<i>Urtica</i>	212. 224
<i>urens</i>	—
<i>dioica</i>	225

5. PENTANDRIA.

<i>Xanthium</i>	212. 225
<i>strumarium</i>	—
<i>Amaranthus</i>	212. 225
<i>Blitum</i>	—

6. POLYANDRIA.

<i>Ceratophyllum</i>	212. 225
<i>demersum</i>	—
<i>submersum</i>	—
<i>Myriophyllum</i>	212. 225
<i>spicatum</i>	—
<i>verticillatum</i>	—

<i>Sagittaria</i>	Pag. 213. 225	<i>myrsinites</i>	Pag. 233
<i>sagittifolia</i>	—	<i>arbuscula</i>	—
<i>Poterium</i>	213. 226	α . - - - -	—
<i>Sanguiforba</i>	—	β . - - - -	—
<i>Quercus</i>	—	γ . - - - -	—
<i>Robur</i>	—	herbacea	—
α . <i>pedunculata</i>	—	** <i>Fol. integerr. glabr.</i>	—
β . <i>fessilis</i>	—	reticulata	—
<i>Fagus</i>	213. 226	<i>myrtilloides</i>	234
<i>lylvatica</i>	—	glauca	—
<i>Carpinus</i>	213. 226	*** <i>Fol. integerr. villos.</i>	—
<i>Betulus</i>	—	<i>aurita</i>	—
<i>Corylus</i>	214. 226	<i>lanata</i>	—
<i>Avellana</i>	—	β . <i>depressa</i>	—
7. MONADELPHIA.	—	<i>Lapponum</i>	—
<i>Pinus</i>	214. 226	<i>arenaria</i>	—
<i>lylvestris</i>	—	<i>incubacea</i>	—
<i>Abies</i>	227	<i>repens</i>	—
β . <i>viminalis</i>	—	<i>fulca</i>	—
8. SYNGENESIA.	—	<i>rosmarinifolia</i>	—
<i>Bryonia</i>	214. 227	**** <i>Fol. subserrat. villos.</i>	—
<i>alba</i>	—	<i>capraea</i>	235
XXII. DIOECIA.	—	α . - - - -	—
I. MONANDRIA.	—	β . - - - -	—
<i>Najas</i>	228. 232	<i>viminalis</i>	—
<i>marina</i>	—	<i>cinerea</i>	—
2. DIANDRIA.	—	<i>alba</i>	—
<i>Vallisneria</i>	228. 232	3. TRIANDRIA.	—
<i>spiralis</i>	—	<i>Empetrum</i>	228. 235
<i>Salix</i>	228. 232	<i>nigrum</i>	—
* <i>Fol. serratis glabris.</i>	—	4. TETRANDRIA.	—
hermaphroditica	—	<i>Viscum</i>	229. 235
triandra	—	<i>album</i>	—
pentandra	—	<i>Hippophaë</i>	229. 235
β . <i>polyandra</i>	—	<i>Rhamnoides</i>	—
<i>phylicifolia</i>	—	<i>Myrica</i>	229. 236
β . - - - -	—	<i>Gale</i>	—
<i>Vitellina</i>	—	5. PENTANDRIA.	—
<i>amygdalina</i>	233	<i>Cannabis</i>	229. 236
<i>hastata</i>	—	<i>fativa</i>	—
<i>fragilis</i>	—	<i>Humulus</i>	229. 236
<i>purpurea</i>	—	<i>Lupulus</i>	—
		6. OCTANDRIA.	—

6. OCTANDRIA.

<i>Populus</i>	Pag. 230. 236
<i>alba</i>	—
<i>tremula</i>	—
<i>nigra</i>	—
<i>Rhodiola</i>	230. 236
<i>rosea</i>	—

7. ENNEANDRIA.

<i>Mercurialis</i>	230. 236
<i>perennis</i>	—
<i>annua</i>	—
<i>Hydrocharis</i>	230. 236
<i>Morsus Ranae</i>	—

8. POLYANDRIA.

<i>Serratoides</i>	231. 237
<i>Aloides</i>	—

9. MONADELPHIA.

<i>Juniperus</i>	231. 237
<i>communis</i>	—
<i>α. minor</i>	—
<i>β. major</i>	—

<i>Taxus</i>	231. 237
<i>baccata</i>	—

XXIII. POLYGAMIA.

I. MONOECIA.

<i>Veratrum</i>	238. 240
<i>album</i>	—
<i>Holcus</i>	238. 240
<i>mollis</i>	—
<i>lanatus</i>	—
<i>odoratus</i>	—
<i>Valantia</i>	238. 240
<i>cruciata</i>	—
<i>Parietaria</i>	238. 240
<i>officinalis</i>	—
<i>Atriplex</i>	238. 240
<i>Portulacoides</i>	—
<i>laciniata</i>	—
<i>hastata</i>	—
<i>patula</i>	241
<i>littoralis</i>	—
<i>marina</i>	—
<i>pedunculata</i>	—

Acer

Pag. 239. 241

<i>Pseudoplatanus</i>	—
<i>Platanoides</i>	—
<i>campestre</i>	—

2. DIOECIA.

<i>Fraxinus</i>	239. 241
<i>excelsior</i>	—

XXIV. CRYPTOGRAMIA.

I. FILICES.

<i>Equisetum</i>	242. 247
<i>sylvaticum</i>	—
<i>arvense</i>	—
<i>Telmateja</i>	—
<i>palustre</i>	—
<i>β. polytachion</i>	—
<i>fluviatile</i>	—
<i>limosum</i>	—
<i>hyemale</i>	—
<i>Ophioglossum</i>	242. 247
<i>vulgatum</i>	—
<i>Osmunda</i>	242. 248
<i>Lunaria</i>	—
<i>β. spica gemina</i>	—
<i>γ. rutacea</i>	—
<i>δ. matricariaefolia</i>	—
<i>regalis</i>	—
<i>Struthiopteris</i>	—
<i>Spicant</i>	—
<i>crispa</i>	—
<i>Acrostichum</i>	242. 248
<i>septentrionale</i>	—
<i>hyperboreum</i>	—
<i>ilvense</i>	—
<i>Pteris</i>	242. 248
<i>aquilina</i>	—
<i>Asplenium</i>	242. 249
<i>Scolopendrium</i>	—
<i>Trichomanes</i>	—
<i>Breynii</i>	—
<i>Ruta muraria</i>	—
<i>Adiantum nigrum</i>	—

Poly.

<i>Polypodium</i>	Pag. 242. 249	<i>Buxbaumia</i>	Pag. 243. 253
vulgare	—	aphylla	—
Lonchitis	—	foliosa	—
fontanum	—	<i>Splachnum</i>	243. 253
montanum	—	rubrum	—
Phegopteris	250	luteum	254
cristatum	—	ampullaceum	—
Filix mas	—	vasculosum	—
spinulosum	—	angustatum	—
molle	—	mnioides	—
Filix foemina	—	sphaericum	—
Thelypteris	—	<i>Polytrichum</i>	243. 254
aculeatum	—	commune	—
fragile	251	α. ferrulatum	—
Dryopteris	—	β. ciliatum	—
<i>Trichomanes</i>	242. 251	piliferum	—
tunbrigense	—	alpinum	—
<i>Ptilularia</i>	242. 251	urnigerum	—
globulifera	—	<i>Mnium</i>	244. 255
<i>Isoetes</i>	243. 251	pellucidum	—
lacustris	—	androgynum	—
2. MUSCI.		fontanum	—
<i>Lycopodium</i>	243. 251	palustre	—
clavatum	—	hygrometricum	—
Selaginoides	—	purpureum	—
inundatum	253	setaceum	—
Selago	—	cirrhatum	—
annotinum	—	annotinum	—
alpinum	—	hornum	—
complanatum	—	capillaceum	—
dubium	—	capillare	256
<i>Sphagnum</i>	243. 252	crudum	—
palustre	—	pyriforme	—
<i>Phascum</i>	243. 252	Polytrichoides	—
aculon	—	serpyllifolium	—
muticum	—	α. punctatum	—
subulatum	253	β. cuspidatum	—
<i>Fontinalis</i>	243. 253	γ. proliferum	—
antipyretica	—	δ. undulatum	—
minor	—	triquetrum	—
squamosa	—	α. - - - -	—
capillacea	—	β. - - - -	—
secunda	—	γ. - - - -	—

Bryum

<i>Bryum</i>	Pag. 244. 257			Pag. 260
apocarpum	—		argenteum	—
α . aureum	—		pulvinatum	—
β . coccineum	—		pendulum	—
striatum	—		bimum	—
α . minus	—		caespitium	—
β . majus	—		virens	—
γ . tenerius	—		carneum	—
δ . - - -	—		simplex	—
pomiforme	—		alpinum	—
pyriforme	—		hyperboreum	—
extinctorium	—		Schisti	261
α . minus	—		Oederi	—
β . majus	—		<i>Hypnum</i>	244. 261
subulatum	—		* <i>Frondib. pinnatis.</i>	—
rurale	—		taxifolium	—
murale	—		denticulatum	—
scoparium	258		Bryoides	—
undulatum	—		Adiantoides	—
glaucum	—		Trichomanoides	—
pellucidum	—		complanatum	—
flexuosum	—		sylvaticum	—
unguiculatum	—		** <i>Surculis vagis.</i>	—
capillaceum	—		lucens	—
aciculare	—		rugosum	262
heteromallum	—		undulatum	—
tortuosum	—		crispum	—
truncatum	259		triquetrum	—
viridulum	—		rutabulum	—
paludosum	—		*** <i>Surculis pinnatis.</i>	—
laterale	—		flicinum	—
hypnoides	—		proliferum	—
α . hirsutum	—		delicatum	—
β . pilosum	—		parietinum	—
γ . virescens	—		praelongum	—
δ . microcarpos	—		Crista castrensis	—
ϵ . obtusum	—		abietinum	263
ζ . barbatum	—		plumosum	—
verticillatum	—		*** <i>Foliis recurv.</i>	—
aestivum	—		cupressiforme	—
Celsii	—		aduncum	—
trichodes	—		Scorpioides	—
isuarrosum	260		viriculofum	—
			aquaticum	—
				isquarro.

<i>Isquarrosom</i>	Pag. 263	** <i>Fr. pinn. pinnul. antr.</i>	
<i>α. majus</i>		<i>sup. testis.</i>	
<i>β. minus</i>		<i>nemorea</i>	Pag. 267
<i>palustre</i>	264	<i>resupinata</i>	—
<i>loreum</i>	—	<i>albicans</i>	—
***** <i>Surc. dendr. v. fascic.</i>		<i>trilobata</i>	—
<i>dendroides</i>	—	<i>reptans</i>	—
<i>alopecurum</i>	—	<i>multiflora</i>	—
***** <i>Surc. teretiusc.</i>		<i>purpurea</i>	—
<i>curtipendulum</i>	—	*** <i>Froncl. imbricat.</i>	
<i>puium</i>	—	<i>complanata</i>	—
<i>illecebrum</i>	—	<i>dilatata</i>	—
<i>riparium</i>	—	<i>tamariscifolia</i>	268
<i>cuspidatum</i>	—	<i>platyphylla</i>	—
<i>α. pungens</i>		<i>ciliaris</i>	—
<i>β. inerme</i>		<i>tomentella</i>	—
***** <i>Surcul. confert.</i>		<i>varia</i>	—
<i>sericeum</i>	265	<i>pulcherrima</i>	—
<i>α. curvatum</i>		*** <i>Surc. undique imbricat.</i>	
<i>β. praelongum</i>		<i>julacea</i>	—
<i>velutinum</i>	—	<i>cinninata</i>	—
<i>serpens</i>	—	<i>rupestris</i>	—
<i>sciuroides</i>	—	<i>trichophylla</i>	269
<i>gracile</i>	—	<i>alpina</i>	—
<i>myosuroides</i>	—	<i>Sertularioides</i>	—
<i>clavellatum</i>	—	**** <i>Acaules fr. simpl.</i>	
<i>Halleri</i>	—	<i>epiphylla</i>	—
<i>nitens</i>	—	<i>pinguis</i>	—
		<i>multifida</i>	—
		<i>furcata</i>	—
		<i>pusilla</i>	—
3. ALGAE.		<i>Targionia</i>	244. 269
<i>Jangermannia</i>	244. 266	<i>hypophylla</i>	—
* <i>Froncl. pinnat. secund.</i>		<i>Marchantia</i>	244. 269
<i>Sphagni</i>	—	<i>polymorpha</i>	—
<i>alplenoides</i>	—	<i>α. stellata</i>	
<i>viticulosa</i>	—	<i>β. umbellata</i>	
<i>polyanthos</i>	—	<i>hemisphaerica</i>	270
<i>lanceolata</i>	—	<i>conica</i>	—
<i>bidentata</i>	—	<i>tenella</i>	—
<i>bicuspidata</i>	—	<i>Blasia</i>	244. 270
<i>fissa</i>	—	<i>pusilla</i>	—
<i>quinquedentata</i>	267		

Riccia

PLANTARUM.

367

<i>Riccia</i>	Pag. 245. 270	trapeziformis	Pag. 274
<i>crystallina</i>	—	<i>rufescens</i>	—
<i>minima</i>	—	<i>fulgens</i>	—
<i>glauca</i>	—	<i>rufus</i>	—
<i>fluitans</i>	—	<i>sphaerocephalus</i>	—
<i>natans</i>	—	<i>β. quercinus</i>	—
<i>Anthoceros</i>	245. 270	<i>globularis</i>	—
<i>punctatus</i>	—	<i>fulvus</i>	—
<i>laevis</i>	—	** <i>Lepr. scutell.</i>	—
<i>Lichen</i>	245. 271	<i>Jolithus</i>	—
* <i>Leprosi tuberc.</i>	—	<i>vitellinus</i>	—
<i>scriptus</i>	—	<i>candelarius</i>	275
<i>geographicus</i>	—	<i>flavescens</i>	—
<i>atrovirens</i>	—	<i>tartareus</i>	—
<i>lacteus</i>	—	<i>gelidus</i>	—
<i>rupicola</i>	—	<i>β. Heclae.</i>	—
<i>pertusus</i>	—	<i>rimosus</i>	—
<i>rugosus</i>	—	<i>frigidus</i>	—
<i>sanguinarius</i>	—	<i>palescens</i>	—
<i>Oederi</i>	—	<i>lentigerus</i>	—
<i>Koenigii</i>	272	<i>subfuscus</i>	—
<i>fulco- ater</i>	—	<i>Muscorum</i>	—
<i>vernalis</i>	—	<i>Saxorum</i>	276
<i>calcareus</i>	—	<i>Parellus</i>	—
<i>cinereus</i>	—	<i>subcarneus</i>	—
<i>atro- albus</i>	—	<i>Upfalienfis</i>	—
<i>niger</i>	—	<i>Pezizoides</i>	—
<i>confluent</i>	—	<i>brunneus</i>	—
<i>terminalis</i>	—	<i>Ulmi</i>	—
<i>ventosus</i>	—	<i>ater</i>	—
<i>Haematomma</i>	—	<i>amylaceus</i>	—
<i>fagineus</i>	273	<i>varius</i>	—
<i>carpineus</i>	—	<i>impressus</i>	277
<i>corallinus</i>	—	<i>scruposus</i>	—
<i>pseudocorallinus</i>	—	<i>Vahlh</i>	—
<i>Baeomyces</i>	—	<i>aurantiacus</i>	—
<i>Icmadophila</i>	—	*** <i>Imbricati.</i>	—
<i>candidus</i>	—	<i>microphyllus</i>	—
<i>variegatus</i>	—	<i>Hypnorum</i>	—
<i>ocellatus</i>	—	<i>crassus</i>	—
<i>β. punctatus</i>	—	<i>saxicola</i>	278
<i>marmoreus</i>	—	<i>luridus</i>	—
<i>roseus</i>	274	<i>stillicidiorum</i>	—

Junger.

	Pag. 278		Pag. 283
<i>Jungermanniae</i>	—	<i>granulatus</i>	—
<i>leucophaeus</i>	—	<i>β.</i>	—
<i>pulverulentus</i>	—	<i>rupestris</i>	—
<i>centrifugus</i>	279	<i>tenax</i>	—
<i>cartilagineus</i>	—	<i>palmatus</i>	—
<i>saxatilis</i>	—	<i>marginalis</i>	—
<i>omphalodes</i>	—	<i>fascicularis</i>	—
<i>olivaceus</i>	—	<i>granosus</i>	284
<i>pullus</i>	—	<i>gyrosus</i>	—
<i>diffusus</i>	—	***** <i>Coriacei.</i>	—
<i>Fahlunensis</i>	—	<i>herbaceus</i>	—
<i>stygius</i>	—	<i>scrobiculatus</i>	—
<i>crispus</i>	280	<i>aquaticus</i>	—
<i>sinuatus</i>	—	<i>fluvialis</i>	—
<i>muralis</i>	—	<i>resupinatus</i>	—
<i>parietinus</i>	—	<i>venosus</i>	285
<i>physodes</i>	—	<i>aphthosus</i>	—
<i>stellaris</i>	—	<i>arcticus</i>	—
*** <i>Foliacei.</i>	—	<i>caninus</i>	—
<i>Burgesii</i>	—	<i>sylvaticus</i>	—
<i>ciliaris</i>	—	<i>polydactylus</i>	—
<i>hispidus</i>	281	<i>horizontalis</i>	286
<i>islandicus</i>	—	<i>faccatus</i>	—
<i>α. rigidus</i>	—	<i>croceus</i>	—
<i>β. tenuifolius</i>	—	***** <i>Umbilicati.</i>	—
<i>γ. nigricans</i>	—	<i>griseus</i>	—
<i>groenlandicus</i>	—	<i>miniatus</i>	—
<i>nivalis</i>	—	<i>velleus</i>	—
<i>pulmonarius</i>	—	<i>β. glaucus</i>	—
<i>furfuraceus</i>	—	<i>pustulatus</i>	287
<i>farinaceus</i>	282	<i>exasperatus</i>	—
<i>calicaris</i>	—	<i>erosus</i>	—
<i>scopulorum</i>	—	<i>Cribellum</i>	—
<i>fraxineus</i>	—	<i>proboscideus</i>	288
<i>Pranaetri</i>	—	<i>deustus</i>	—
<i>α. incanus</i>	—	<i>polyphyllus</i>	—
<i>β. albidus</i>	—	<i>β. polyphyllus</i>	—
<i>juniperinus</i>	—	<i>γ. glaber</i>	—
<i>caperatus</i>	—	<i>complicatus</i>	—
<i>glaucus</i>	—	<i>polyrhizos</i>	—
<i>nigrescens</i>	283	<i>reticulatus</i>	289

***** *Scy.*

PLANTARUM

369

***** Scyphiferi.

cocciferus Pag. 289

α.

β.

γ.

δ.

cornucopioides

pixidatus

α.

β.

γ.

δ.

ε.

ζ.

fimbriatus

α. simplex

β. prolifer

gracilis

α.

β.

γ.

digitatus

α. tuberculatus

β. etuberculatus

cornutus

deformis

radiatus

cristatus

foliaceus

***** Fruticulosi.

Papillaria

rangiferinus

α. alpestris

β. sylvaticus

pallidus

uncialis

fubulatus

vermicularis

arenarius

globiferus

paschalis

fragilis

Botrytis

Mulcicola

tristis

***** Filamentosi.

plicatus

barbatus

divaricatus

jubatus

lanatus

pubescens

chalybeiformis

hirtus

floridus

vulpinus

* Dubii generis.

radiciformis

Tremella

juniperina

mesenterica

Noctoc

crispa

sagarum

affinis

orbicularis

rubra

agaricoides

verrucosa

difformis

hemisphaerica

viridis

purpurea

granulosa

α. nigra

β. rubra

adnata

Fucus

ferratus

vesiculosus

divaricatus

inflatus

ceranoides

β. lacerus

As

Pag. 292

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

<i>spiralis</i>	Pag. 296	<i>vittatus</i>	Pag. 300
<i>canaliculatus</i>	—	<i>ramentaceus</i>	301
<i>β. excisus</i>	—	<i>plumosus</i>	—
<i>distichus</i>	—	<i>abrotanifolius</i>	—
<i>soboliferus</i>	—	<i>cartilagineus</i>	—
<i>pumilus</i>	297	<i>gigartinus</i>	—
<i>nodosus</i>	—	<i>spermophorus</i>	—
<i>filiquosus</i>	—	<i>confervoides</i>	—
<i>corallinus</i>	—	<i>Scorpioides</i>	—
<i>elongatus</i>	—	<i>albus</i>	302
<i>loreus</i>	—	* <i>Fuci wond. facis determ.</i>	—
<i>foeniculaceus</i>	—	<i>flagelliformis</i>	—
<i>β. barbatus</i>	—	<i>viridis</i>	—
<i>granulatus</i>	—	<i>laciniatus</i>	—
<i>Selaginoides</i>	298	<i>Oederi</i>	—
<i>ericoides</i>	—	<i>norvegicus</i>	—
<i>aculeatus</i>	—	<i>Fascia</i>	—
<i>β. muscoides</i>	—	<i>clavatus</i>	—
<i>Lycopodioides</i>	—	<i>coronopifolius</i>	—
<i>hirsutus</i>	—	<i>Ulva</i>	245. 302
<i>discors</i>	—	<i>umbilicalis</i>	—
<i>Filum</i>	—	<i>plicata</i>	—
<i>lanosus</i>	—	<i>intestinalis</i>	—
<i>fastigiatus</i>	—	<i>β. prolifera</i>	—
<i>furcellatus</i>	—	<i>spongiformis</i>	303
<i>palmaris</i>	299	<i>compressa</i>	—
<i>miniatus</i>	—	<i>incrassata</i>	—
<i>ovinus</i>	—	<i>clavata</i>	—
<i>bullatus</i>	—	<i>latissima</i>	—
<i>lanceolatus</i>	—	<i>β. maxima</i>	—
<i>β. epiphyllus</i>	—	<i>Lactuca</i>	—
<i>digitatus</i>	—	<i>lanceolata</i>	—
<i>β. bifurcatus</i>	—	<i>Linza</i>	—
<i>esculentus</i>	—	<i>flavescens</i>	—
<i>saccharinus</i>	—	<i>pruniformis</i>	—
<i>sanguineus</i>	300	<i>granulata</i>	—
<i>roseus</i>	—	<i>radicata</i>	—
<i>ciliatus</i>	—	<i>Conferva</i>	245. 304
<i>crispus</i>	—	<i>rivularis</i>	—
<i>crispatus</i>	—	<i>fontinalis</i>	—
<i>alatus</i>	—	<i>bullosa</i>	—
<i>dentatus</i>	—	<i>canalicularis</i>	—
<i>rubens</i>	—	<i>distorta</i>	—
			hirsuta

PLANTARUM.

371

hirta	Pag. 304	aquatica	Pag. 307
amphibia	—	Elos Aquae	—
littoralis	—	cancellata	—
dichotoma	—	phosphorea	—
nidifica	—	velutina	—
cancellata	—	bombycina	308
elongata	—	aurea	—
omissa	—	Cryptarum	—
reticulata	305	antiquitatis	—
** <i>Filaments. articul.</i>		saxatilis	—
fluviatilis	—	candelaris	—
Linum	—	botryoides	—
nitida	—	incana	—
jugalis	—	lactea	—
Pennatula	—	rosea	—
fracta	—	fruticulosa	—
hirta	—		
flexuosa	—	4. FUNGI.	
uncialis	—	<i>Agericus</i>	245. 309
gelatinosa	—	* <i>Pileo albo.</i>	
polymorpha	306	umbelliferus	—
α . fascicularis		androsaceus	—
β . barbata		piperatus	—
γ . corallina		extinctorius	—
δ . congesta		β . cinctus	—
ϵ . caudata		pratorum	—
seposita	—	campestris	—
vagabunda	—	Kremlinga	—
glomerata	—	procerus	—
rupestris	—	praealtus	310
floccosa	—	praecox	—
hypnoides	—	gramineus	—
diaphana	—	cernuus	—
Aegagropila	—	niveus	—
stellaria	—	nitidus	—
<i>Corallina</i>	245. 307	stellatus	—
officinalis	—	** <i>Pileo flavo.</i>	
squamata	—	cinnamomeus	—
corniculata	—	Buillardii	—
rubens	—	rubescens	—
cristata	—	dentatus	—
<i>Byssus</i>	245. 307	Cantarellus	311
leptica	—	Georgii	—
		A & 2	deli.

	Pag. 311		Pag. 314
deliciosus		clypeatus	
β. fulvus		mammosus	
clavus	—	equestris	—
caesarius	—	separatus	—
citrinus	—	quinquepartitus	315
aurantius	—	fimbriatus	—
squarrosus	—	plumbeus	—
squamatus	—	stellaris	—
conicus	—	mutabilis	—
auratus	—	crassipes	—
mutabilis	—	crenulatus	—
glutinosus	312	***** <i>Acaul. dimidiat.</i>	—
plicatus	—	nigricans	—
cyathiformis	—	striatus	—
membranaceus	—	pubescens	—
melleus	—	Quercinus	—
muricatus	—	Betulinus	316
ochraceus	—	Alneus	—
niger	—	<i>Bolus</i>	245. 316
cervinus	—	inversus	—
torminosus	—	perennis	—
unicolor	—	Polyporus	—
nivalis	313	lateralis	—
resupinatus	—	bovinus	—
*** <i>Pileo nigricanoe.</i>	—	rubiginosus	—
violaceus	—	luteus	—
echinatus	—	viscidus	—
purpureus	—	convexus	317
scaber	—	granulatus	—
caesius	—	gregarius	—
cyanipes	—	terreus	—
bicolor	—	subtomentosus	—
**** <i>Pileo rubro.</i>	—	subsquamosus	—
lactifluus	—	resupinatus	—
viscidus	—	frondosus	—
Kermesinus	314	caudicinus	—
integer	—	Buglossum	—
barbatus	—	fomentarius	318
Apicii	—	uberofus	—
muscarius	—	igniarius	—
***** <i>Pileo cinereo.</i>	—	versicolor	—
campanulatus	—	suaveolens	—
fimetarius	—	Favus	—
			cellu-

<i>cellulosus</i>	Pag. 318	* <i>Forte ad alia g. est.</i>	
<i>dubius</i>	—	<i>membranacea</i>	Pag. 321
<i>Hydnum</i>	246. 318	<i>Pineti</i>	—
<i>imbricatum</i>	—	<i>Clavaria</i>	246. 322
<i>cyathiforme</i>	—	<i>pilipes</i>	—
<i>repandum</i>	—	<i>simplex</i>	—
<i>tomentosum</i>	—	<i>vermicularis</i>	—
<i>Erinaceus</i>	319	<i>pistillaris</i>	—
<i>Auriscalpium</i>	—	<i>media</i>	—
<i>gelatinosum</i>	—	<i>ophioglossoides</i>	—
<i>parasiticum</i>	—	<i>inaequalis</i>	—
<i>Thaelaephora</i>	—	<i>purpurea</i>	—
<i>Phallus</i>	246. 319	<i>ascendens</i>	—
<i>esculentus</i>	—	<i>polymorpha</i>	—
<i>impudicus</i>	—	<i>militaris</i>	—
<i>Clathrus</i>	246. 319	<i>Hypoxylon</i>	—
<i>cancellatus</i>	—	<i>cornuta</i>	—
<i>nudus</i>	—	<i>Garchariae</i>	323
<i>denudatus</i>	—	<i>pileata</i>	—
<i>recutitus</i>	320	<i>coralloides</i>	—
<i>ramosus</i>	—	<i>fastigiata</i>	—
<i>α. aureus</i>	—	<i>muscosa</i>	—
<i>β. rufus</i>	—	<i>Peziza</i>	246. 323
<i>Helvella</i>	246. 320	<i>lentifera</i>	—
<i>Mitra</i>	—	<i>α. laevis</i>	—
<i>α. tomentosa</i>	—	<i>β. hirsuta</i>	—
<i>β. nuda</i>	—	<i>γ. sericea</i>	—
<i>lacunosa</i>	—	<i>granulifera</i>	—
<i>α. lurida</i>	—	<i>punctata</i>	324
<i>β. pallida</i>	—	<i>cornucopioides</i>	—
<i>fulcata</i>	—	<i>Acetabulum</i>	—
<i>α. fusca</i>	—	<i>pyxidata</i>	—
<i>β. cinerea</i>	—	<i>virgultorum</i>	—
<i>atra</i>	—	<i>denticulata</i>	—
<i>Pezizoides</i>	—	<i>parvula</i>	—
<i>aeruginosa</i>	—	<i>Ciborium</i>	—
<i>triceps</i>	321	<i>Phiala</i>	—
<i>lubrica</i>	—	<i>turbinata</i>	—
<i>revoluta</i>	—	<i>carnea</i>	—
<i>horizontalis</i>	—	<i>cyathoides</i>	—
<i>spathulata</i>	—	<i>scabra</i>	—
<i>verticalis</i>	—	<i>cupularis</i>	325
<i>Xerium</i>	—	<i>urceolata</i>	—
		<i>A a 3</i>	scu,

374. CONSPECTUS PLANTARUM.

<i>scutellata</i>	Pag. 325	<i>stipitatum</i>	Pag. 327
<i>β. albicans</i>		<i>β.</i>	
<i>γ. laevis</i>		<i>γ.</i>	
<i>pusilla</i>	—	<i>variolosum</i>	—
<i>minuta</i>	—	<i>pisiforme</i>	—
<i>equina</i>	—	<i>Epidendrum</i>	328
<i>cochleata</i>	—	<i>epiphyllum</i>	—
<i>α. fulva</i>		<i>β. poculiforme</i>	
<i>β. ochracea</i>			
<i>Auricula</i>	—	<i>Mucor</i>	246. 328
<i>zonalis</i>	—	<i>urceolatus</i>	—
		<i>Embolus</i>	—
<i>Lycoperdon</i>	246. 325	<i>obvelatus</i>	—
<i>* Subterr. absq. radice.</i>		<i>Scoparius</i>	—
<i>Tuber</i>	—	<i>conoideus</i>	—
<i>Semen</i>	326	<i>Mucedo</i>	—
<i>** Pulverul. radic.</i>		<i>glauca</i>	329
<i>Bovista</i>	—	<i>Druparum</i>	—
<i>stellatum</i>	—	<i>viridescens</i>	—
<i>multifidum</i>	—	<i>Erysiphe</i>	—
<i>Carpobolus</i>	—	<i>septicus</i>	—
<i>radiatum</i>	—	<i>tubulosus</i>	—
<i>pedunculatum</i>	—	<i>α. Tubulif. Cremor</i>	
<i>*** Parasitica.</i>		<i>β. Tubulif. Ceratum</i>	
<i>cancellatum</i>	—	<i>Lachnum</i>	246. 329
<i>corniferum</i>	—	<i>Agaricinum</i>	—
<i>penicillatum</i>	—		
<i>aggregatum</i>	327	<i>* Fungi dubii generis.</i>	
<i>mammosum</i>	—	<i>Byssus fruticulosa</i>	330
<i>fessile</i>	—	<i>Acrospermum unguinosum</i>	—
<i>gregarium</i>	—	<i>Acrospermum cylindricum</i>	—

INDEX

- Acer 1262—1264.
 Achillaea 1048—1050.
 Aconitum 662—663.
 Acorus 416.
 Acrosperrum 2018, 2019.
 Acrostichum 1279—1281.
 Adæa 648.
 Adonia 678—679.
 Adoxa 486.
 Aegopodium 371.
 Aethusa 354.
 Agaricus 1808—1824.
 Agrimonia 589.
 Agrostemma 571.
 Agrostis 88—95.
 Aira 96—104, 120.
 Ajuga 704—706.
 Alchemilla 204—206.
 Alisma 452—454.
 Allium 389—394.
 Alopecurus 83—86.
 Alfine 377—378.
 Althaea 852.
 Alyssum 798, 799.
 Amaranthus 1190.
 Anagallis 249.
 Anchusa 227.
 Andromeda 494—498.
 Androsace 237.
 Anemone 665—672.
 Angelica 342, 343.
 Anthemis 1044—1047.
 Anthericum 405—408.
 Anthoceros 1483, 1484.
 Anthoxanthum 41.
 Anthyllis 874.
 Antirrhinum 762—766.
 Apargia 941, 942.
 Aphanes 206.
 Apium 370.
 Aquilegia 664.
 Arabis 825, 826, 825.
 Arbutus 499, 500.
 Arctium 975.
 Arenaria 553—558.
 Aristolochia 1112.
 Asnica 1034.
 Artemisia 997—1001.

- Arum 1113.
 Arundo 155—159.
 Asarum 587.
 Asclepias 294.
 Asparagus 409.
 Asperugo 233.
 Asperula 179. 180.
 Asplenium 1283—1287.
 Asprella 71.
 Aster 1023. 1024.
 Astragalus 902—907.
 Athamanta 334. 335.
 Atriplex 1255—1261.
 Atropa 27.
 Avena 149—154.
 Axuris 1181.
 Azalea 250. 251.
 Ballota 741. 742.
 Bartfia 744.
 Bellis 1095.
 Berberis 438.
 Betonica 725.
 Betula 1183.
 Bidens 990—993.
 Blasia 1477.
 Boletus 185—190.
 Brassica 830. 831.
 Briza 124. 142.
 Bromus 136—147.
 Bryonia 1203.
 Bryum 1352—1391.
 Bunias 838. 839.
 Bunium 330. 331.
 Bupleurum 326. 327.
 Butomus 832.
 Buxbaumia 1341. 1342.
 Byllus 1791—1869. 1523.
 Calendula 1057.
 Calla 1114.
 Callitriche 5. 6.
 Caltha 703.
 Campanula 255—265.
 Cannabis 1237.
 Cardamine 803—810.
 Carduus 979—981. 982. 983.
 984. 985. 986.
 Carex 1131—1180.
 Carlina 989.
 Carpinus 1199.
 Carum 367.
 Centaurea 1052—1056.
 Centunculus 198.
 Cerastium 577—582.
 Ceratophyllum 1191. 1192.
 Chara 1118—1121. 1175.
 Chaerophyllum 361—363.
 Cheiranthus 825.
 Chelidonium 649. 650.
 Chenopodium 295—305.
 Chondrilla 932.
 Chrysanthemum 1036—1038.
 Chrysocoma 995.
 Chrysosplenium 506. 507.
 Cichorium 974.
 Cicuta 355.
 Cineraria 1026. 1027.
 Circaea 18. 9.
 Cistus 658—660.
 Cladonia 1634. 1641.
 Clathrus 1922—1926.
 Clavaria 1941—1958. 1937.
 Clinopodium 735.
 Cnicus 982—987.
 Cochlearia 701—706.
 Comarum 642.
 Conferva 1751—1785. 1696.
 1750.
 Conium 331.

Con-

Convallaria 410—414.
Convolvulus 252. 253.
Conyza 1008.
Corallina 1786—1790.
Coralloides 1643. 1644. 1645.
Coreopsis 1051.
Cornicularia 1648.
Cornus 200. 201.
Coronilla 897. 898.
Corrigiola 376.
Corylus 1200.
Corula 1043.
Crambe 838.
Crassula 386.
Crataegus 604—607. 609.
Crepis 967. 968.
Cucubalus 535—537.
Cuscuta 207. 208.
Cynoglossum 228.
Cynofunus 125. 126.
Cyperus 54. 55.
Cypripedium 1110. 1111.

D.

Dactylis 123.
Daphne 47.
Datura 274.
Daucus 329.
Delphinium 661.
Dentaria 802.
Dianthus 529—534.
Diapensia 236.
Digitalis 770.
Dipsacus 173. 174.
Draba 777—783.
Dracocephalum 739. 740.
Drosera 384. 385.
Dryas 646.

E.

Echium 235.
Elatine 487. 488.

Elvela 1977. 1978. 1979.
Elymus 163. 164.
Embolus 2005.
Empetrum 1233.
Epilobium 457—467.
Equisetum 1266—1272.
Erica 472—474.
Erigeron 1009—1011.
Eriophorum 67—69.
Ervum 894—896.
Eryngium 322. 323.
Erysimum 818—822.
Euonymus 282.
Eupatorium 994.
Euphorbia 591—596.
Euphrasia 746. 747.

F.

Fagus 1198.
Festuca 127—135.
Filago 1058—1060.
Fontinalis 1317—1321.
Fragaria 630.
Fraxinus 1265.
Fritillaria 397.
Fucus 1676—1736.
Fumaria 858—862.

G.

Galeopsis 722—724.
Galium 181—190.
Genista 867—870.
Gentiana 311—321.
Geranium 839—851.
Geum 644. 645.
Glaux 291.
Glechoma 718.
Globularia 172.

Gnaphalium 1002—1007.

Gratiola 32.

Gypsophila 526. 527.

H.

Hedera 289.

Hedysarum 900.

Helleborus 702.

Helvella 1927.

Heracleum 338. 339.

Herniaria 294.

Hesperis 824.

Hieracium 955—966.

Hippophaë 1235.

Hippuris 3. 4.

Holcus 1250—1252.

Holosteum 170.

Hordeum 165. 166.

Hottbomia 243.

Humulus 1238.

Hwirkälsfvamp 1984.

Hyacinthus 415.

Hydnum 1911—1919.

Hydrocharis 1245.

Hydrocotyle 324.

Hyofcyamus 275.

Hyoseris 969.

Hypericum 929—934.

Hypnum 1392—1436.

Hypochaeris 970—972.

I.

Jacobaea 1027.

Jasione 1061.

Iberis 797.

Ilex 209.

Illecebrum 290.

Impatiens 1071.

Imperatoria 364.

Inula 1028—1033.

Iris 45—48.

Isatis 837.

Isnardia 202.

Isoetes 1304.

Juncus 417—437.

Jungermannia 1437—1471.

Juniperus 1247.

K.

Koenigia 171.

L.

Lachnum 2016.

Lactuca 950. 951.

Lamium 719—721.

Lapfana 973.

Lalerpitium 336. 337.

Lathraea 753.

Lathyrus 880—885.

Lavatera 857.

Ledum 492. 493.

Lemna 1122—1125.

Leontodon 954. 941. 942.

Leonurus 734.

Lepidium 784—787.

Lichen 1485—1659.

Lichenoides 1574. 1576. 1580.

Ligusticum 340. 341.

Ligustrum 7.

Lilium 395. 396.

Limofella 772.

Limfwamp 1668.

Linnea 771.

Linum 302. 303.

Lithospermum 225. 226.

Litto-

Littorella 1182.
Lobelia 1062.
Lolium 160. 161.
Lonicera 268 — 270.
Lotus 924. 925.
Lunaria 800. 801.
Lychnis 572 — 576.
Lycoperdon 1983 — 2002.
 1668.

Lycopodium 1305 — 1312.
Lycopsis 234.
Lycopus 39.
Lysimachia 244 — 248.
Lythrum 588.

M.

Malaxis 1099.
Malva 853 — 856.
Marchantia 1473 — 1476.
Marrubium 733.
Matricaria 1039 — 1042.
Medicago 926 — 928.
Melampyrum 748 — 752.
Melica 105 — 108.
Mentha 711 — 717.
Menyanthes 241. 242.
Mercurialis 1243. 1244.
Mespilus 610.
Milium 87.
Mnium 1235 — 1351. 1444.
 1452.

Monotropa 491.
Montia 169.
Mucor 2003 — 2015.
Myagrum 774. 775.
Myolotis 223. 224.
Myofurus 388.
Myrica 1236.
Myriophyllum 1193. 1194.

N.

Najas 1204.
Nardus 70.
Nepeta 710.
Nymphaea 655. 656.

O.

Oenanthe 350 — 352.
Oenothera 456.
Ononis 872. 873.
Onopordum 988.
Ophioglossum 1233.
Ophrys 1092 — 1104. 1076.
Orchis 1072 — 1087.
Origanum 736.
Ornithogalum 399 — 402.
Ornithopus 899.
Orobanche 773.
Orobus 877 — 879.
Osmunda 1274 — 1278.
Othonna 1026.
Oxalis 569. 570.

P.

Panicum 74 — 78.
Papaver 651 — 654.
Parietaria 1254.
Paris 485.
Parnassia 379.
Pastinaca 366.
Pasellaria 1494. 1517. 1524.
 1534. 1536. 1538. 1544.
 1546.
Pedicularis 754 — 761.
Peltigera 1603. 1607. 1608.
 1611.
Peplis 439.

Peziza

- Peziza 1959—1982. 1668.
 Phaca 901.
 Phalaris 72. 73.
 Phallus 1920. 1921. 1933.
 1934.
 Phascum 1314—1316.
 Pheilandrium 353.
 Phleum 79—82.
 Physalis 277.
 Phyteuma 266.
 Picris 939. 940.
 Pilularia 1303.
 Pimpinella 368. 369.
 Pinguicula 33—34.
 Pinus 1201. 1202.
 Pisum 875. 876.
 Plantago 191—197.
 Plazifma 1584. 1587.
 Poa 109—122.
 Polemonium 254.
 Polygala 863—865.
 Polygonum 476—484.
 Polypodium 1288—1304.
 Polytrichum 1331—1334. 1353.
 Populus 1239—1241.
 Potamogeton 210—218.
 Potentilla 631—642.
 Poterium 1196.
 Prenanthes 953.
 Primula 238—240.
 Prunella 743.
 Prunus 598—608.
 Pfora 1569.
 Pteris 1282.
 Pulehoharia 229—231. 1579.
 1598. 1599.
 Pyrola 501—505.
 Pyrus 611. 612. 604.
- Q.
- Quercus 1197.
- R.
- Ranunculus 680—700.
 Raphanus 834.
 Refeda 590.
 Rhamnus 280. 281.
 Rheum 489.
 Rhinanthus 745.
 Rhodiola 1242.
 Ribes 283—288.
 Riccia 1478—1482.
 Rosa 616—623.
 Rortböllia 162.
 Rubus 624—629.
 Rumex 440—448.
 Ruppia 219.
- S.
- Sagina 220. 221.
 Sagittaria 1195.
 Salicornia 1. 2.
 Salix 1208—1232.
 Salfola 306. 307.
 Salvia 248.
 Sambucus 373. 374.
 Samolus 267.
 Sanguiforba 199.
 Sanicula 223.
 Saponaria 523.
 Satyrium 1088—1091.
 Saxifraga 508—523.
 Scabiosa 175—177.
 Scandix 356—360.
 Scheuchzeria 449.
 Schoenus 49—53. 64.
 Scilla 403. 404.
 Scirpus 56—66.
 Scleranthus 524. 525.
 Scorzonera 937. 938.
 Scrophellaria 767—769.
 Scutellaria 741. 742.
- Sedum

Sedum 559—568.
Selinum 332. 333.
Sempervivum 597.
Senecio 1017—1022.
Serapias 1105—1109.
Serratula 976—978.
Seseli 365.
Sherardia 178.
Sibbaldia 387.
Silene 538—542.
Sinapis 832. 833. 834.
Sison 349.
Sisymbrium 811—817.
Sium 344—348.
Solanum 278. 279.
Solidago 1025.
Sonchus 943—949.
Sorbus 608. 609.
Sparganium 1128—1130.
Spartium 866.
Spergula 583—586.
Sphagnum 1313—1321.
Sphaeria 1674. 1999.
Spiraea 613—615.
Splachnum 1324—1330.
Squamaria 1557. 1558. 1565.
 1566. 1586.
Stachys 726—730.
Statice 380. 381.
Stellaria 543—552.
Stipa 148.
Stratiotes 1246.
Subularia 776.
Swertia 309. 310.
Symphytum 234.

T.

Tamarix 375.
Tanacetum 996.
Targionia 1472.
Taxus 1248.

Teucrium 708. 709.
Thalictrum 673—677.
Thaëlaephora 1919.
Thesium 292. 293.
Thlaspi 788—790.
Thymus 737. 738.
Tilia 657.
Tillaea 222.
Tordylium 328.
Tormentilla 643.
Tragopogon 935. 936.
Trapa 203.
Tremella 1660—1675. 1590.
 1751. 1981. 1999.
Trichomanes 1302.
Trientalis 455.
Trifolium 909—923.
Triglochin 450. 451.
Triticum 167. 168. 147.
Trollius 701.
Tubulifera 2015.
Tulipa 398.
Turritis 827—829.
Tussilago 1012—1016.
Typha 1126. 1127.

U.

Ulex 871.
Ulmus 308.
Ulua 1737—1751. 1710.
 1714. 1720.
Umbilicaria 1615. 1616.
Urtica 1187. 1188.
Usnea 1656. 1657.
Utricularia 36. 37.

V.

Vaccinium 468—471.
Valantia 1253.

Vale.

Valeriana 42—44.

Vallisneria 1205.

Veratrum 1249.

Verbascum 271—273.

Verbena 38.

Veronica 10—31.

Verrucaria 1487. 1489. 1492.

1501. 1504. 1513. 1514.

Viburnum 372.

Vicia 886—893.

Viola 1063—1070.

Viscum 1234.

X.

Xanthium 1189.

Z.

Zannichellia 1117.

Zostera 1115. 1116.





